Natural Resources for Human Health
eISSN: 2583-1194

Original Research

Received 15 Oct 2025
Revised 20 Nov 2025
Accepted 28 Dec 2025

Natural Resources for Human Health 2025, Vol. 5 (4): 998–1006

KEYWORDS:
Nursing intervention, constipation, bedridden patients, hydration, fiber intake

Nursing Interventions for Preventing Constipation in Bedridden Hospitalized Patients

Ahmed Saad Hadaj Alhadi[1], Aleuayduh, Bandar Khilifuh E[2], Alanouod Talal Alsuliman[3], Farhan Nazal Fulayyih Alruwaili[4], Nazal Owaynana Alruwaili[5], Alruwaili, Rasmiah Rajaa M[6], Abdullah Saud Alshammari[7], Ali Saeed Jarw Alruwaili[8], Afra Hamoud Fatil Aldaghmani[9], Fahad Saeed Jaro Alrawoaili[10]

[1] Nursing Technician – Warehouses Department, Supply Executive Vice Presidency – Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia
Email: ahmed.alhadaj@hotmail.com
[2] Nursing Technician – Eradah and Mental Health Hospital, Al Jouf – Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia
Email: balawidah@moh.gov.sa
[3] Nursing Technician – Internal Communication Department – Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia
Email: aalsaluman@moh.gov.sa
[4] Nursing Technician – Eradah and Mental Health Hospital, Al Jouf – Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia
Email: fal-rwele@moh.gov.sa
[5] Nursing Technician – Khowa Health Center – Al Jouf Health Cluster – Ministry of Health – Al Jouf Region – Saudi Arabia
Email: nazaloa@moh.gov.sa
[6] Nursing Technician – Al-Tuwair Primary Health Care Center – Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia
Email: rralrawily@gmail.com
[7] Nursing Technician – Warehouse Management Department, Deputy Chief Executive Officer for Supply – Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia
Email: zaaeed2011@hotmail.com
[8] Nursing Technician – Mental Health Hospital – Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia
Email: aloosh32@gmail.com
[9] Nursing Technician – Cardiac Center Al-Jouf – Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia
Email: aloosh32@gmail.com
[10] Nursing Technician – Al Mukhattat Primary Healthcare Center – Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia
Email: aloosh32@gmail.com

ABSTRACT: Constipation is a common issue faced by bedridden hospitalized patients due to factors such as immobility, poor dietary intake, and inadequate fluid consumption. Nurses play a crucial role in implementing interventions that can prevent this condition. Key nursing interventions include assessing the patient's bowel habits and history to

establish a baseline, encouraging fluid intake by offering frequent sips of water or clear fluids, and promoting a high-fiber diet when appropriate. Additionally, nurses can facilitate physical activity, even within the constraints of bed rest, by encouraging passive range-of-motion exercises or repositioning to stimulate bowel function. Regularly scheduled toileting routines can also be beneficial, as they create a habit that may prompt bowel movements. Moreover, patient education about the importance of diet and hydration can empower patients and caregivers to partake in managing constipation. Utilizing laxatives or stool softeners may be warranted as a preventive strategy based on the individual's health status and physician recommendations. Monitoring bowel movements and documenting any changes is essential to evaluate the effectiveness of the interventions implemented. It is important for nurses to communicate with the healthcare team regarding any persistent issues related to constipation, ensuring a collaborative approach to care that addresses the underlying causes and improves patient comfort.

INTRODUCTION

Constipation is one of the most frequent yet under-recognized complications of hospitalization, particularly among patients whose mobility is restricted by illness, surgery, trauma, or critical care. For the bedridden patient, the combination of prolonged immobility, altered dietary intake, insufficient hydration, unfamiliar toileting conditions, and the use of constipating medications creates a cumulative risk that is often neither systematically assessed nor proactively managed [1]. Nevertheless, constipation in bedridden patients continues to be treated reactively rather than prevented, and the consequences range from discomfort and abdominal distension to fecal impaction, stercoral ulceration, colonic perforation, and prolonged hospital stay [2].

Constipation in the hospital setting is not a trivial symptom. Among acutely hospitalized older adults, being bedridden for more than two weeks has been associated with a six-fold increase in the risk of dissatisfaction with bowel emptying, and cerebrovascular events independently elevate that risk [3]. The Merck Manual explicitly notes that when people stay in bed or are less active, stool moves more slowly through the intestine and rectum, and that hospital medications, particularly certain pain relievers, further compound the problem [4]. For the bedridden patient, the physiological consequences of immobility are superimposed on the iatrogenic effects of opioids, anticholinergics, calcium channel blockers, and iron supplementation, as well as on the psychological discomfort of using a bedpan without privacy [5]. The complications of unrelieved constipation are clinically serious. Fecal impaction may present with overflow diarrhea, nausea, vomiting, or abdominal pain, and in frail bedridden patients it can progress to stercoral ulceration and perforation [2]. Stercoral colitis, a rare but life-threatening complication of chronic constipation, results from increased colonic intraluminal pressure leading to mucosal edema, inflammation, and ulceration, and it has been described in bedbound patients requiring total care [6]. Constipation also prolongs intensive care unit and hospital length of stay and contributes to intra-abdominal infections [7]. Prevention, therefore, is not merely a comfort measure but a patient safety imperative.

Defining and Recognizing Constipation in the Bedridden Patient

The definition of constipation in hospitalized patients requires both objective and subjective elements. The Rome IV criteria for functional constipation require the presence of two or more of the following: straining during more than one quarter of defecations, lumpy or hard stools in more than one quarter of defecations, a sensation of incomplete evacuation in more than one quarter of defecations, a sensation of anorectal obstruction or blockage in more than one quarter of defecations, manual maneuvers to facilitate defecation in more than one quarter of defecations, and fewer than three spontaneous bowel movements per week [8]. Although the Rome IV framework was developed for chronic constipation, its components are clinically useful for structuring nursing assessment in the hospital setting, particularly when a premorbid bowel pattern can be established from the patient or family. Stool form should be documented using the Bristol Stool Form Scale, which classifies stool into seven types; types 1 and 2 represent hard, lumpy stool consistent with constipation, types 3 to 4 represent soft, formed stool, and types 5 to 7 represent increasingly loose stool [9]. In bedridden patients, the Bristol scale provides a practical, reproducible means of documenting stool consistency at the bedside and can be incorporated into a simple bowel diary that records frequency, consistency, volume, straining, and the use of aids such as suppositories or enemas.

Natural Resources for Human Health
eISSN: 2583-1194

Risk Factors and Pathophysiology

The pathophysiology of constipation in bedridden patients is multifactorial, but reduced physical activity is the central mechanism. Immobility decreases colonic transit, reduces the mechanical stimulation of the colon by abdominal musculature and gravity, and impairs the defecation reflex that is normally facilitated by upright posture and ambulation [4]. Prolonged bed rest also predisposes to dehydration because patients may limit oral intake to reduce the need for toileting assistance, and inadequate fluid intake directly hardens stool and slows transit. Dietary fiber intake is frequently low in hospitalized patients because of anorexia, nil-by-mouth status, dysphagia, or the monotony of hospital meals, and low fiber reduces stool bulk and the frequency of propulsive colonic contractions. Opioid analgesics are among the most potent constipating drugs because they inhibit gastrointestinal motility and secretion through mu-opioid receptors in the enteric nervous system, and prophylactic laxatives are generally recommended when opioids are initiated, particularly in patients who are bedbound or already constipated [10]. Anticholinergic medications, calcium channel blockers, diuretics, iron supplements, and some antipsychotics also contribute. Psychological and environmental factors are equally important: lack of privacy, dependence on staff for bedpan placement, unfamiliar toilet facilities, and the indignity of defecating in bed can lead to voluntary suppression of the urge to defecate, which in turn leads to increased water absorption from retained stool and worsening constipation [5]. Table 1 summarizes the principal risk factors and their clinical implications.

Table 1. Risk Factors for Constipation in Bedridden Hospitalized Patients

Risk factor	Mechanism	Nursing implication
Prolonged immobility	Reduced colonic transit; loss of gravity-assisted defecation	Prioritize passive and active range-of-motion exercise; avoid unnecessary bed rest
Inadequate fluid intake	Hardened stool; reduced stool water content	Offer fluids regularly; target 1,500–2,000 mL/day when not contraindicated
Low dietary fiber	Reduced stool bulk and propulsive contractions	Collaborate with dietitian; introduce fiber gradually if tolerated
Opioid analgesics	Inhibition of motility and secretion	Anticipate constipation; advocate for prophylactic laxatives
Anticholinergics, calcium channel blockers, iron	Slowed transit or hardened stool	Review medication list daily; flag constipating agents
Lack of privacy and bedpan use	Voluntary suppression of defecation urge	Provide screens, curtains, and uninterrupted time; simulate normal posture
Unfamiliar environment and routine	Disruption of conditioned defecation reflex	Establish a consistent toileting schedule based on the patient's premorbid pattern
Dehydration from fever, drains, or restricted intake	Reduced stool water	Monitor intake and output; assess mucous membranes and skin turgor

Systematic Nursing Assessment and Early Identification

Prevention begins with assessment, yet constipation is frequently overlooked because nurses and physicians prioritize the presenting medical or surgical problem. A structured bowel assessment should be completed on admission and at regular intervals thereafter, and it should include the patient's premorbid bowel pattern, current medications, fluid and fiber intake, mobility status, abdominal symptoms, and any history of laxative use. The use of laxatives at home has been identified as the only independent risk factor for objective constipation during hospitalization in acutely hospitalized older patients, which underscores the importance of taking a careful laxative history [3]. A three-day bowel record is a practical nursing tool: it documents the date and time of each bowel movement, Bristol stool type, volume, degree of straining, and any assistance required, and it provides the baseline against which the effectiveness of interventions can be judged. Physical assessment should include abdominal inspection and auscultation, palpation for fecal loading, and, where appropriate and within scope of practice, a digital rectal examination to exclude impaction. However, in a patient who is bedbound or has severe dementia, an abdominal

radiograph or digital rectal examination may be necessary to diagnose impaction, and bulk-forming laxatives should be avoided in patients with limited mobility because of the risk of worsening impaction or obstruction [10]. Table 2 outlines the core components of a nursing bowel assessment for the bedridden patient.

Table 2. Components of a Structured Nursing Bowel Assessment

Domain	Key elements	Rationale
History	Premorbid frequency, consistency, straining, laxative use, fiber and fluid habits	Establishes the patient's normal pattern and identifies pre-existing risk
Medication review	Opioids, anticholinergics, calcium channel blockers, iron, diuretics	Identifies iatrogenic contributors that may be modified
Physical assessment	Abdominal distension, bowel sounds, palpable fecal mass, rectal examination if indicated	Detects fecal loading or impaction
Stool documentation	Bristol Stool Form Scale, frequency, volume, straining	Provides objective, reproducible data for monitoring
Fluid and dietary intake	Oral intake records, intravenous fluids, fiber content of diet	Identifies modifiable deficits
Mobility and function	Bed mobility, ability to use bedpan or commode, transfer status	Determines the feasible level of activity and toileting assistance
Environmental and psychological factors	Privacy, embarrassment, dependence on staff, anxiety	Guides dignity-preserving care and bowel training

Non-Pharmacological Nursing Interventions

Non-pharmacological interventions are the foundation of constipation prevention in bedridden patients, and they are particularly important because pharmacological agents carry risks of interaction, overuse, and dependence. A research-based interdisciplinary protocol that combined dietary fiber, increased fluid intake, and hygiene measures in hospitalized immobile vascular surgery patients reduced the incidence of constipation from 59% to approximately 9% over a three-year period, demonstrating that a multicomponent non-pharmacological approach can be highly effective [11]. The scoping review of nurse-led strategies similarly found that abdominal massage, reflexology, acupressure, auricular therapy, dietary fiber solutions, and structured bowel management protocols were the most frequently evaluated interventions, with most studies reporting reductions in constipation severity scores, improved stool consistency, and increased bowel movement frequency [1].

Dietary Fiber and Fluid Management

Adequate fluid intake is essential for maintaining stool softness, and a daily intake of 1,500 to 2,000 mL is commonly recommended for adults who are expected to be able to manage such a volume and who have no cardiac or renal contraindication [12]. In bedridden patients, fluids should be offered at regular intervals rather than only with meals, and nurses should document actual intake rather than assuming that a full water pitcher has been consumed. Fiber intake should be increased gradually to approximately 25 to 30 grams per day, using foods such as prunes, bran, oats, fruits, vegetables, and legumes, but fiber supplementation must be accompanied by adequate fluid intake because fiber without water can worsen constipation and increase the risk of impaction [10]. In patients with poor oral intake or dysphagia, a dietitian should be consulted to identify palatable, texture-appropriate fiber sources and to consider oral nutritional supplements with added fiber. Prune juice and warm fluids are traditional nursing measures that may help stimulate the gastrocolic reflex, particularly when offered in the morning.

Physical Activity and Passive Range-of-Motion Exercise

Although bedridden patients cannot ambulate, they can still receive structured movement interventions that stimulate intestinal motility. Passive range-of-motion exercises of the lower extremities and trunk increase parasympathetic activity and have been shown to improve intestinal motility in patients who must remain in bed [13]. A randomized controlled trial in enterally fed

palliative care patients found that in-bed passive exercise significantly reduced constipation, nausea, vomiting, and abdominal distension, and that both abdominal massage and in-bed exercise improved comfort and reduced pain compared with a control group [14]. Nurses can incorporate passive leg and trunk movements into routine care, perform them twice daily for approximately 15 minutes, and coordinate with physiotherapists to ensure that the exercises are safe for the patient's clinic al condition. Even small amounts of bed mobility, such as turning, sitting up at the edge of the bed, and performing ankle pumps, can contribute to maintaining some level of physical stimulation and should be encouraged whenever clinically permissible.

Abdominal Massage

Abdominal massage is a safe, inexpensive, nurse-delivered intervention with a growing evidence base. Swedish abdominal massage has been shown to be an effective and non-invasive alternative to laxatives for the management of secondary constipation in hospitalized patients [15]. Effleurage abdominal massage, a gentle rhythmic technique, has demonstrated promise in reducing the time to first postoperative bowel movement and improving stool consistency in patients with lower limb fractures [16]. A meta-analysis cited in the palliative care trial demonstrated that abdominal massage is an effective intervention for chronic constipation, significantly improving straining and reducing gastrointestinal complications in bedridden patients [14]. The technique typically involves superficial and deep effleurage, petrissage, vibration, and repeated superficial effleurage in a clockwise direction following the course of the colon, and it is usually performed for 10 to 15 minutes once or twice daily, ideally before breakfast or before the planned time of defecation. Contraindications include suspected bowel obstruction, recent abdominal surgery where massage is not permitted, abdominal aortic aneurysm, and unstable abdominal conditions, and a medical order or nursing protocol should specify the parameters of practice.

Toileting Positioning and Bowel Training

Defecation is facilitated by an upright or squatting posture that aligns the rectum and anal canal and allows the abdominal muscles and pelvic floor to relax. For patients who cannot use a toilet, the squat position can be simulated in bed by placing the patient in a left-side-lying position with the knees bent and the legs drawn toward the abdomen, which facilitates the passage of feces through the transverse and descending colon [17]. When the patient's condition allows sitting, a bedside commode with the feet supported and the trunk leaning slightly forward is preferable to a bedpan because it approximates the normal defecation posture and preserves dignity. A individualized toileting schedule should be based on the patient's premorbid bowel pattern, and a common recommendation is to offer toileting or bedpan placement 20 to 30 minutes after breakfast, when the gastrocolic reflex is most active. The schedule should be consistent, unhurried, and protected from interruption, and the patient should be encouraged not to rush or strain excessively. Bowel training also involves teaching the patient to respond to the urge to defecate rather than suppressing it, and to use abdominal breathing and gentle bearing-down efforts that coordinate with the abdominal muscles rather than prolonged Valsalva straining.

Privacy, Dignity, and Psychological Support

Lack of privacy is a recognized contributor to hospital-acquired constipation, and immobile patients in particular may feel ashamed and embarrassed when defecating in bed, leading to voluntary retention [5]. Every effort should be made to provide a private environment using curtains, screens, and closed doors, and care should be organized so that the patient is not interrupted during defecation. Nurses should communicate sensitively, avoid comments that may cause embarrassment, and allow sufficient time for the patient to defecate without feeling rushed. Offering a warm drink, adjusting the bed to a comfortable position, and providing a bedpan or commode that is warm and clean are simple measures that promote relaxation and reduce the psychological barrier to defecation. In patients with cognitive impairment or dementia, a calm and familiar approach, together with consistent toileting cues, may be more effective than verbal instruction alone.

Table 3. Summary of Non-Pharmacological Nursing Interventions

Intervention	Recommended approach	Key evidence or rationale
Fluid optimization	Offer 1,500–2,000 mL/day in small, frequent amounts unless contraindicated	Adequate hydration softens stool and supports fiber action [12]
Fiber-rich diet	Gradually increase to 25–30 g/day; use prunes, bran, fruits, vegetables, legumes	Fiber increases stool bulk and stimulates propulsion [10]

Passive range-of-motion exercise	Twice daily, 15 minutes, lower limbs and trunk, as tolerated	Improves intestinal motility and reduces constipation in bedridden patients [13,14]
Abdominal massage	Clockwise effleurage and petrissage for 10–15 minutes once or twice daily	Reduces constipation severity and improves stool consistency [14,15,16]
Positioning	Left lateral position with knees flexed for bedbound patients; bedside commode with feet supported when possible	Simulates squat posture and facilitates fecal passage [17]
Toileting schedule	Individualized, consistent, typically after breakfast; unhurried and private	Uses gastrocolic reflex and re-establishes conditioned defecation pattern
Privacy and dignity	Screens, curtains, uninterrupted time, sensitive communication	Reduces voluntary suppression and promotes relaxation [5]

Pharmacological Prophylaxis and Nursing Considerations

Although non-pharmacological measures are the first line of prevention, pharmacological prophylaxis is appropriate for high-risk bedridden patients, particularly those receiving opioids or those who have not responded to lifestyle measures. The choice of agent should be guided by the patient's clinical status, the likely mechanism of constipation, and the adverse effect profile. Stimulant laxatives such as senna and bisacodyl increase colonic peristalsis and secretion and are commonly used in opioid-induced constipation because they address the motility component [18]. Osmotic laxatives, including polyethylene glycol and lactulose, retain water in the bowel and soften stool, and polyethylene glycol is often better tolerated and more effective than lactulose in hospitalized patients [12]. Stool softeners such as docusate are widely used, although evidence for their efficacy as monotherapy is limited, and they are often combined with a stimulant. Bulk-forming laxatives should be avoided in patients with limited mobility or fluid restriction because they may increase the risk of fecal impaction and obstruction [10]. Suppositories and enemas are generally reserved for patients who have not responded to oral laxatives or who require rapid relief, and they should be used cautiously in neutropenic, thrombocytopenic, or cardiac patients. Nurses play a critical role in administering these agents at the correct time, monitoring the response, documenting bowel movements, and recognizing adverse effects such as diarrhea, electrolyte imbalance, and rectal irritation [12]. Table 4 summarizes the main laxative classes and their nursing implications.

Table 4. Pharmacological Agents for Constipation Prevention and Nursing Considerations

Class	**Examples**	**Nursing considerations**
Stimulant laxatives	Senna, bisacodyl	Effective in opioid-induced constipation; anticipate abdominal cramping; avoid in suspected obstruction
Osmotic laxatives	Polyethylene glycol, lactulose	Ensure adequate fluid intake; monitor for diarrhea and electrolyte disturbance; polyethylene glycol often better tolerated
Stool softeners	Docusate sodium, docusate calcium	Limited efficacy alone; often combined with stimulant; ensure adequate fluid intake
Bulk-forming laxatives	Psyllium, ispaghula	Avoid in bedbound or fluid-restricted patients because of impaction risk [10]
Suppositories	Bisacodyl, glycerin	Reserve for rescue; administer per protocol; assess for rectal bleeding or discomfort
Enemas	Phosphate, sodium chloride	Reserve for refractory cases; monitor electrolytes; avoid in renal impairment and cardiac disease

Structured Bowel Management Protocols

The heterogeneity of constipation prevention practices in hospitals has prompted interest in standardized, nurse-driven bowel protocols that specify assessment frequency, non-pharmacological measures, and escalation to pharmacological therapy. A standardized nurse-driven bowel regimen in critically ill patients did not reduce the incidence of constipation lasting more than 72 hours, but it significantly reduced the likelihood of prolonged constipation lasting more than six days, from 44% to 18%, after adjustment for opioid exposure, illness severity, and age [7]. This finding suggests that protocolized care may be particularly valuable in preventing the progression from acute to prolonged constipation, even if it does not eliminate all early cases. The scoping review of nurse-led strategies similarly identified structured bowel management pathways and standardized assessment protocols as important components of nursing practice, although the heterogeneity of interventions and outcome measures precluded definitive conclusions about comparative effectiveness [1]. A structured protocol typically includes a bowel assessment on admission, a daily bowel diary, a non-pharmacological bundle (fluids, fiber, mobility, massage, positioning), a stepped pharmacological algorithm, and clear criteria for escalation to medical review. Table 5 outlines the core components of such a protocol.

Table 5. Components of a Structured Nurse-Driven Bowel Management Protocol

Component	Specification
Admission assessment	Bowel history, medication review, risk stratification, baseline Bristol stool type
Daily monitoring	Bowel diary documenting frequency, consistency, straining, and assistance required
Non-pharmacological bundle	Fluid target, fiber target, passive exercise, abdominal massage, positioning, toileting schedule, privacy
Pharmacological step 1	Stool softener or osmotic laxative for patients at moderate risk or receiving constipating medications
Pharmacological step 2	Add stimulant laxative if no bowel movement within 48–72 hours
Rescue therapy	Suppository or enema if no response to oral measures, per protocol
Escalation	Medical review for suspected impaction, obstruction, or persistent constipation beyond 72 hours
Documentation and audit	Regular audit of protocol adherence and constipation outcomes

Patient Education and Discharge Planning

Prevention of constipation does not end at discharge, particularly for patients who will remain bedridden or have limited mobility at home. Patient and caregiver education should cover the importance of adequate fluid and fiber intake, the use of abdominal massage and passive exercises, the recognition of warning signs such as absence of bowel movement for more than three days, abdominal pain, vomiting, or overflow diarrhea, and the safe use of laxatives as prescribed. Discharge planning should include a written bowel management plan, a supply of prescribed laxatives, and clear instructions about when to seek medical advice. The use of laxatives at home before admission has been identified as a risk factor for constipation during hospitalization, which suggests that patients with a history of laxative use require particular attention during transitions of care [3]. Nurses should also educate family caregivers about the importance of privacy and dignity during defecation, and about the correct technique for assisting with bedpan use or commode transfers [17].

Barriers to Implementation and Strategies for Improvement

Despite strong evidence for non-pharmacological interventions, several barriers impede their consistent implementation in busy hospital wards. Constipation is often viewed as a low-priority issue compared with the patient's primary medical or surgical condition, and bowel care may be delegated to the least experienced staff or omitted when the ward is short-staffed. Nurses may lack knowledge about the Rome IV criteria, the Bristol Stool Form Scale, or the contraindications to certain laxatives, and they may not have access to structured assessment tools or bowel protocols. Patients may be reluctant to discuss bowel habits because of embarrassment, and cultural or language barriers may further complicate communication. Organizational barriers

include the absence of a standardized bowel protocol, insufficient documentation systems, and a lack of audit and feedback on constipation outcomes. Strategies to overcome these barriers include incorporating bowel assessment into admission and shift-handover checklists, providing education and competency training for nurses, appointing bowel-care champions or link nurses, using electronic health record prompts to document bowel movements and trigger interventions, and auditing constipation prevalence and protocol adherence as quality indicators. Interprofessional collaboration with dietitians, physiotherapists, and pharmacists is essential, because fiber optimization, safe physical activity, and medication review cannot be achieved by nursing alone [18].

CONCLUSION

Constipation in bedridden hospitalized patients is a predictable, multifactorial, and largely preventable complication. The evidence reviewed here supports a proactive, multicomponent nursing approach that begins with systematic risk assessment and bowel documentation, proceeds through a bundle of non-pharmacological interventions including fluid and fiber optimization, passive exercise, abdominal massage, positioning, toileting schedules, and privacy protection, and escalates to pharmacological prophylaxis when indicated. Structured nurse-driven bowel protocols can reduce prolonged constipation and provide a framework for consistent, auditable care. However, implementation requires organizational commitment, interprofessional collaboration, and sustained nursing education. By treating bowel care as an essential component of patient safety and dignity rather than an optional comfort measure, nurses can substantially reduce the burden of constipation and its serious complications in this vulnerable population.

REFERENCES

[1] Pinto CF, Oliveira PD, Fernandes OM, Padilha JM, Machado PA, Ribeiro AL, Ramos JL. Nonpharmacological clinical effective interventions in constipation: a systematic review. Journal of Nursing Scholarship. 2020 May;52(3):261-9.

[2] Salvi F, Petrino R, Conroy SP, Liperoti R, Paoletti L, Beccacece A, dell'Aquila G, Fedecostante M, Cherubini A. Constipation: a neglected condition in older emergency department patients. Internal and Emergency Medicine. 2024 Oct;19(7):1977-86.

[3] Cardin F, Minicuci N, Droghi AT, Inelmen EM, Sergi G, Terranova O. Constipation in the acutely hospitalized older patients. Archives of gerontology and geriatrics. 2010 May 1;50(3):277-81.

[4] Chernova I, Krishnan N. Resistant hypertension updated guidelines. Current cardiology reports. 2019 Oct;21(10):117.

[5] Kelly AM. Public toilets and their potential impact on an individual's health. British Journal of Nursing. 2024 Jun 20;33(12):538-44.

[6] Laubli D, Eapen D, El Hachem P. SEVERE STERCORAL COLITIS: A CASE TREATED IN THE HOSPICE SETTING. InJOURNAL OF GENERAL INTERNAL MEDICINE 2022 Jun 1 (Vol. 37, pp. S518-S519). ONE NEW YORK PLAZA, SUITE 4600, NEW YORK, NY, UNITED STATES: SPRINGER.

[7] Krenz Z, Bennett A, Hall K, Phillips T, Henry K. Impact of a standardised nurse-driven bowel regimen on bowel frequency in critically ill patients. Nursing in Critical Care. 2025;30(6):e70207.

[8] Eltorki M, Bhattacharjee A, Khan M, Martin E, Shyleyko R, Freedman SB. Treatment failure in children diagnosed with constipation in a paediatric emergency department in relation to Rome III criteria. Paediatrics & Child Health. 2019 May 15;24(3):185-92.

[9] Harrison WN, Kumar A, Quinonez R, Stephens JR. Things We Do for No Reason™: Routinely hospitalizing children with fecal impaction for inpatient cleanouts. Journal of Hospital Medicine. 2023 May;18(5):439-43.

[10] Luo J, To WL, Xu Q, Zhang J, Ma Y, Chow S, Yu DJ, Yuan CS, Bian Z. Clinical practice guidelines for the diagnosis of constipation-predominant irritable bowel syndrome and functional constipation in adults: a scoping review. BMC gastroenterology. 2025 9;25(1):234.

[11] Hall GR, Karstens M, Rakel B, Swanson E, Davidson A. Managing constipation using a research-based protocol. Medsurg nursing: official journal of the Academy of Medical-Surgical Nurses. 1995 Feb 1;4(1):11-8.

[12] Andrews A, St Aubyn B. Assessment, diagnosis and management of constipation. Nurs Stand. 2020 Aug 10;35(9):59-65.

[13] Turan M, Cengiz Z. The effect of abdominal massage and in-bed ROM exercise on gastrointestinal complications and comfort in intensive care unit patients receiving enteral nutrition: A randomized controlled trial. Japan Journal of Nursing Science. 2024 Jul;21(3):e12602.

[14] Turan M, Cengiz Z. The effect of abdominal massage and in-bed ROM exercise on gastrointestinal complications and comfort in intensive care unit patients receiving enteral nutrition: A randomized controlled trial. Japan Journal of Nursing Science. 2024 Jul;21(3):e12602.

[15] Nouhi E, Mansour-Ghanaei R, Hojati SA, Chaboki BG. The effect of abdominal massage on the severity of constipation in elderly patients hospitalized with fractures: A randomized clinical trial. International journal of orthopaedic and trauma nursing. 2022 Nov 1;47:100936.

[16] Hung E, Juha G, Davenport TE, Williams J. Physical Therapy Management of Myasthenia Gravis: A Case Study. Internet Journal of Allied Health Sciences and Practice. 2024;22(4):6.

[17] Bookbinder M, McHugh ME. Symptom management in palliative care and end of life care. Nursing Clinics. 2010 Sep 1;45(3):271-327.

[18] Damani A, Salins N, Coyle S. Pain management and palliative care in the head and neck cancer setting. InStell & Maran's Head and Neck Surgery and Oncology 2025 Jan 15 (pp. 368-390). CRC Press.