Natural Resources for Human Health
eISSN: 2583-1194

Original Research

Received 12 Oct 2025
Revised 18 Nov 2025
Accepted 25 Dec 2025

Natural Resources for Human Health 2025, Vol. 5 (4): 1104–1110

KEYWORDS:
Tooth extraction, post-extraction complications, alveolar osteitis, dry socket, surgical site infection, dental assistant

Role of Dentists and Dental Assistants in Preventing Post-Extraction Infections and Complications: A Comprehensive Review

Alsemran, Ohood Mohammad N[1], Badoor Jaber S Almoutrafi[2], Noora Khalid M Hamidudin[3], Alanazi, Reem Abdullah F[4], Fatma Yahya Mohammed Alshingiti[5], Bader Awadh Rokhyyan Alenzi[6], Faisal Obaid I Alrashdan[7], Turki Hameed M Alanazi[8], Rahmah Humaidan Fehaid Althubyani[9], Abdulhakim Farooq Abdulhakim Sindi[10]

[1] General Dentist — Third Riyadh Health Cluster, Ministry of Health, Riyadh, Riyadh Region, Saudi Arabia
Email: Aho15od@hotmail.com
[2] Dental Assistant — Prince Abdullah bin Abdulaziz bin Musaed Specialized Dental Center, Northern Borders Health Cluster, Ministry of Health, Arar, Northern Borders Region, Saudi Arabia
Email: Bdooorrr37@gmail.com
[3] General Dentist — Khaybar Health Center, Madinah Health Cluster, Ministry of Health, Jeddah, Makkah Region, Saudi Arabia
Email: n.hamidudin@gmail.com
[4] Dental Assistant — Branch of the Ministry of Health in the Northern Borders Region, Ministry of Health, Arar, Northern Borders Region, Saudi Arabia
Email: reemalanazyy1993@gmail.com
[5] Dental Assistant — Madinah Health Cluster, Ministry of Health, Madinah, Madinah Region, Saudi Arabia
Email: falshingiti@moh.gov.sa
[6] General Dentist — Specialized Dental Center in Al Qurayyat, Al Jouf Health Cluster, Ministry of Health, Al Qurayyat, Al Jouf Region, Saudi Arabia
Email: dr_bader_alsalmi@hotmail.com
[7] General Dentist — Specialized Dental Center in Al Qurayyat, Al Jouf Health Cluster, Ministry of Health, Al Qurayyat, Al Jouf Region, Saudi Arabia
Email: dr.faisal.obaid@gmail.com
[8] Dental Laboratory Technician — Specialized Dental Center in Al Qurayyat, Al Jouf Health Cluster, Ministry of Health, Al Qurayyat, Al Jouf Region, Saudi Arabia
Email: Trky1722@gmail.com
[9] Dental Assistant Technician — Al Faysaliyah Medical Center, Madinah Health Cluster, Ministry of Health, Madinah, Madinah Region, Saudi Arabia
Email: Rahmahhumaidan@gmail.com
[10] General Dentist — Ministry of Health, Madinah, Madinah Region, Saudi Arabia
Email: Afsindi@moh.gov.sa

ABSTRACT: Tooth extraction is one of the most frequently performed surgical procedures in dental practice. Despite its routine nature, it carries a significant risk of post-operative complications, including alveolar osteitis (dry socket), surgical site infections

Natural Resources for Human Health
eISSN: 2583-1194

(SSI), post-extraction bleeding, and nerve injuries. The prevention of these complications relies heavily on the synergistic efforts of the dental team, primarily the dentist and the dental assistant. This comprehensive review article explores the multifaceted roles of both professionals across the pre-operative, intra-operative, and post-operative phases of tooth extraction. We discuss evidence-based protocols for infection control, atraumatic surgical techniques, pharmacological management, and patient education. Furthermore, the critical role of the dental assistant in four-handed dentistry, sterilization protocols, and post-operative patient triage is highlighted. By integrating rigorous scientific guidelines with efficient team dynamics, the dental team can significantly reduce the incidence of post-extraction morbidity, ensuring optimal patient outcomes and enhanced clinical efficiency.

INTRODUCTION

Tooth extraction, or exodontia, is a fundamental surgical procedure in dentistry, performed for various indications ranging from severe caries and periodontal disease to orthodontic requirements and impaction management [1]. While generally safe, the procedure disrupts the oral mucosal barrier and exposes the underlying alveolar bone to the polymicrobial environment of the oral cavity, thereby predisposing the patient to post-operative complications [2]. The most common complications include alveolar osteitis (dry socket), surgical site infections (SSI), prolonged hemorrhage, trismus, and transient or permanent nerve injuries [3].

The incidence of these complications varies significantly based on the complexity of the extraction, the anatomical site, and patient-specific risk factors. For instance, the incidence of dry socket following the extraction of mandibular third molars can range from 5% to 30%, whereas it is significantly lower in the maxilla [4]. Preventing these complications requires a meticulous, evidence-based approach that extends beyond the surgical technique of the dentist to encompass the entire perioperative workflow.

The modern dental practice operates on the principle of team-based care, where the dentist and the dental assistant function as a cohesive unit. The dental assistant plays an indispensable role in infection control, operatory preparation, patient management, and intra-operative efficiency through four-handed dentistry [1].

Pre-Operative Phase: Assessment and Preparation

The foundation of complication-free exodontia is laid during the pre-operative phase. Comprehensive assessment and meticulous preparation are critical to identifying risk factors and optimizing the clinical environment.

Medical History and Risk Assessment

A thorough medical history is the first line of defense against systemic and local complications. The dentist must evaluate the patient for conditions that impair healing or increase infection risk, such as uncontrolled diabetes mellitus, immunosuppression, and a history of head and neck radiation [2]. Furthermore, the use of medications such as bisphosphonates (which carry the risk of medication-related osteonecrosis of the jaw, MRONJ) and anticoagulants must be carefully managed. For patients on anticoagulants, current guidelines recommend proceeding with extractions without altering the medication regimen, provided local hemostatic measures are employed, as the risk of thromboembolism outweighs the risk of post-extraction bleeding [5, 6].

Radiographic Evaluation

Pre-operative radiographic assessment is mandatory to evaluate root morphology, proximity to vital structures (e.g., the inferior alveolar nerve and the maxillary sinus), and the presence of pathological lesions. While periapical and panoramic radiographs are standard, Cone Beam Computed Tomography (CBCT) is increasingly indicated for complex impactions to precisely map the spatial relationship between the tooth roots and the inferior alveolar canal, thereby minimizing the risk of iatrogenic nerve injury [3, 7].

Patient Education and Informed Consent

Setting realistic expectations and educating the patient about post-operative care is crucial. The dentist must explain the procedure, potential complications, and the importance of adhering to post-operative instructions. Modifiable risk factors, such as smoking and poor oral hygiene, must be addressed. Smoking significantly increases the risk of dry socket by impairing blood

clot formation and introducing heat and toxins to the surgical site [4, 8]. Patients should be strongly advised to cease smoking at least 48 to 72 hours post-extraction.

The Dental Assistant's Role in Pre-Operative Setup

The dental assistant is pivotal in the pre-operative phase. Their responsibilities include updating and verifying the patient's medical history, ensuring all consent forms are signed, and preparing the operatory [9, 10]. The assistant must ensure that the sterilization protocols have been strictly followed for all surgical instruments. The surgical tray should be set up efficiently, including the placement of sterile drapes, local anesthetic cartridges, syringes, and extraction forceps and elevators. Proper patient positioning and the application of protective eyewear and lead aprons (if radiographs are taken in the operatory) are also managed by the assistant, ensuring a safe and organized environment before the dentist begins the procedure [11, 12].

Intra-Operative Phase: Asepsis and Surgical Technique

The intra-operative phase is where the physical extraction occurs. Strict adherence to aseptic protocols and the employment of atraumatic surgical techniques are paramount in preventing infections and minimizing tissue trauma.

Infection Control and Sterilization Protocols

Surgical site infections, though relatively rare in healthy patients, can lead to severe morbidity. The oral cavity is heavily colonized, making strict infection control essential. The Centers for Disease Control and Prevention (CDC) guidelines mandate rigorous sterilization of all critical and semi-critical instruments using steam autoclaving, which is the most reliable method for destroying all microbial life, including spores [7]. Surface disinfection of the operatory with EPA-registered hospital-grade disinfectants must be performed between patients [13- 15]. The dental assistant is primarily responsible for the reprocessing of instruments, operating the autoclave, and conducting regular spore testing to ensure the sterilization equipment is functioning correctly [16, 17].

Aseptic Technique and Hand Hygiene

Proper hand hygiene and the use of personal protective equipment (PPE) are non-negotiable. The dentist must perform a surgical hand scrub or use an alcohol-based hand rub before donning sterile gloves [18, 19]. The assistant must assist in the proper donning of sterile gloves and gowns, ensuring that the sterile field is not compromised. The use of sterile drapes to isolate the operative site from the non-sterile oral mucosa further reduces the bacterial load introduced into the extraction socket.

Atraumatic Extraction Techniques

Tissue trauma is a primary catalyst for post-operative complications, including pain, swelling, and dry socket. The dentist must employ atraumatic extraction techniques. This involves the careful reflection of mucoperiosteal flaps (when indicated), conservative bone removal using surgical burs with copious sterile saline irrigation to prevent thermal necrosis, and the sectioning of multi-rooted teeth to avoid excessive force [4]. The use of periotomes or piezosurgery can further minimize soft and hard tissue trauma. By preserving the alveolar bone and surrounding soft tissue, the dentist ensures primary intention healing and reduces the inflammatory response [20].

Irrigation and Debridement

Following the extraction of the tooth, the alveolar socket must be thoroughly debrided. The presence of residual granulation tissue, bone fragments, or calculus can act as a nidus for infection and delay healing. Copious irrigation of the socket with sterile saline is essential to flush out debris and reduce the bacterial load [13]. In cases of active infection (e.g., periapical abscess), some clinicians advocate for the use of chlorhexidine gluconate irrigation, though saline remains the standard to avoid cytotoxicity to the healing tissues [21].

Hemostasis and Wound Closure

Achieving primary hemostasis is critical for the formation of a stable blood clot, which serves as the scaffold for subsequent tissue regeneration. The dentist should achieve hemostasis through local pressure, the placement of resorbable hemostatic agents (such as gelatin sponges or oxidized regenerated cellulose) in the socket, and the approximation of wound margins using

sutures [15]. Suturing not only controls bleeding but also stabilizes the blood clot, significantly reducing the incidence of alveolar osteitis [22].

The Dental Assistant's Role in Intra-Operative Efficiency

During the extraction, the dental assistant's role transitions to active surgical assistance through four-handed dentistry. The assistant must maintain a clear operative field by utilizing high-volume evacuation (HVE) to remove blood, saliva, and irrigation fluids. This not only improves the dentist's visibility but also significantly reduces the aerosolization of bloodborne pathogens and oral microbes, protecting the entire dental team [20]. The assistant is also responsible for the timely transfer of instruments, ensuring the dentist maintains a continuous, efficient workflow, which minimizes the overall operative time and the duration of tissue exposure [23].

Post-Operative Phase: Management and Follow-Up

The post-operative phase begins immediately after the procedure and extends until complete mucosal healing has occurred. Proper management during this phase is crucial for preventing delayed complications.

Immediate Post-Operative Instructions

Clear, concise, and written post-operative instructions must be provided to the patient. The patient should be instructed to bite firmly on a sterile gauze pad placed over the extraction site for 30 to 60 minutes to facilitate clot formation. They must be advised against spitting, rinsing vigorously, using a straw, or smoking for at least 24 hours, as the negative pressure generated can dislodge the blood clot, leading to dry socket [14]. An ice pack applied to the cheek in 20-minute intervals can help minimize post-operative edema.

Pharmacological Management

Pain management is a primary concern for patients post-extraction. A combination of non-steroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen, and acetaminophen has been shown to provide superior analgesia compared to opioids, with fewer side effects [4].

The use of prophylactic antibiotics in routine, uncomplicated extractions in healthy patients is not recommended, as it does not significantly reduce the incidence of infections or dry socket, and contributes to antimicrobial resistance [8]. However, prophylactic antibiotics are indicated for patients with specific systemic conditions (e.g., immunocompromised status, history of infective endocarditis) or in cases of complex surgical extractions with extensive bone removal or pre-existing active infections [9]. Amoxicillin or amoxicillin/clavulanic acid are the first-line choices, while clindamycin or azithromycin can be used for penicillin-allergic patients [5].

The prescription of a 0.12% chlorhexidine gluconate mouthwash, to be used gently starting 24 hours post-operatively, has been proven to significantly reduce the bacterial load in the oral cavity and decrease the incidence of alveolar osteitis [11].

Management of Specific Complications

Alveolar Osteitis (Dry Socket)

Alveolar osteitis is the most common complication following tooth extraction, characterized by severe, throbbing pain that typically begins 2 to 4 days post-operatively, often radiating to the ear or temple. It is caused by the premature fibrinolysis of the blood clot, exposing the alveolar bone [13]. Management involves the gentle irrigation of the socket with warm saline to remove debris, followed by the placement of a medicated dressing, such as zinc oxide eugenol (ZOE) on an alveogyl or gauze strip, to provide palliative relief. The dressing is typically changed every 24 to 48 hours until symptoms subside [16].

Post-Extraction Bleeding

Prolonged bleeding can be alarming for the patient. Initial management involves identifying the source of bleeding (soft tissue vs. bone). Soft tissue bleeding is managed by placing a hemostatic agent and suturing the tissue. Bone bleeding can be controlled by crushing a piece of bone wax into the bleeding bony canal or using oxidized regenerated cellulose. In patients on anticoagulants, the application of tranexamic acid (TXA) mouthwash or soaked gauze provides excellent local hemostasis by inhibiting fibrinolysis [7].

Surgical Site Infections (SSI)

While rare, SSIs can occur, presenting with increased pain, swelling, purulent discharge, and fever. Management requires the establishment of drainage, debridement of the socket, and the initiation of empirical antibiotic therapy, typically with amoxicillin/clavulanic acid or clindamycin, adjusted later based on culture and sensitivity results if necessary [6].

Oroantral Communication and Nerve Injuries

Extraction of maxillary molars can sometimes result in an oroantral communication (OAC). Small OACs (<2 mm) may close spontaneously with conservative management (sinus precautions, decongestants, and antibiotics), while larger defects require surgical closure using a buccal or palatal advancement flap. Inferior alveolar nerve (IAN) injuries, though rare, require immediate documentation, patient counseling, and referral to an oral and maxillofacial surgeon for potential microneurosurgical intervention if sensory deficit persists [2].

The Dental Assistant's Role in Post-Operative Care

The dental assistant plays a vital role in the post-operative phase by ensuring the patient understands the discharge instructions, providing written materials, and scheduling follow-up appointments if necessary. In the event of a post-operative emergency, such as uncontrolled bleeding or severe pain, the assistant is responsible for triaging the call, providing immediate telephonic advice (e.g., applying pressure with a tea bag for bleeding), and scheduling an emergency recall visit for the dentist [21]. The assistant also manages the inventory of post-operative supplies, ensuring that hemostatic agents, sutures, and medicaments are always available.

Special Considerations in High-Risk Patients

Certain patient populations require modified protocols to prevent complications.

Immunocompromised and Diabetic Patients

Patients with uncontrolled diabetes or those undergoing chemotherapy/radiotherapy have impaired wound healing and a higher susceptibility to infections. For these patients, the dentist must ensure strict glycemic control prior to surgery, utilize prophylactic antibiotics, and employ meticulous atraumatic techniques. The dental assistant must ensure that the operatory is thoroughly disinfected and that the highest level of asepsis is maintained to protect these vulnerable individuals from opportunistic infections [19].

Patients on Antiresorptive Therapy (MRONJ)

Patients receiving intravenous bisphosphonates or denosumab for osteoporosis or malignancy are at risk for Medication-Related Osteonecrosis of the Jaw (MRONJ) following extractions. The dentist should consult with the patient's prescribing physician to consider a "drug holiday" if feasible, though current evidence on its efficacy is mixed. The extraction must be performed with extreme care, minimizing bone trauma, and the socket should be closed primarily with sutures. The assistant must ensure that all surgical instruments are impeccably sterile, as any secondary infection can precipitate necrotic bone exposure [10].

Team Dynamics and Communication

The prevention of post-extraction complications is not solely dependent on clinical skills but also on the efficiency and communication of the dental team.

Four-Handed Dentistry and Ergonomics

Four-handed dentistry, where the dentist and assistant work in a synchronized manner, significantly reduces operative time and fatigue. Fatigue can lead to lapses in concentration, increasing the risk of surgical errors and breaches in aseptic technique. The assistant's role in maintaining proper ergonomics for both themselves and the dentist—by adjusting the patient chair, delivering instruments at the point of use, and managing the HVE—allows the dentist to focus entirely on the surgical field [21].

Continuous Education and Training

Both the dentist and the dental assistant must engage in continuous professional development. The assistant should receive regular training on the latest infection control guidelines, sterilization protocols, and emergency management (e.g., BLS/ACLS

certification). The dentist must stay updated on evidence-based surgical techniques and pharmacological guidelines. A well-trained, cohesive team is the most effective safeguard against post-operative complications [22].

CONCLUSION

The prevention of post-extraction infections and complications is a multifaceted endeavor that requires the seamless integration of clinical expertise, evidence-based protocols, and efficient team dynamics. The dentist bears the primary responsibility for accurate diagnosis, atraumatic surgical technique, and appropriate pharmacological management. However, the dental assistant is an equally critical component of this process, ensuring rigorous infection control, operatory preparedness, intra-operative efficiency, and comprehensive post-operative patient management. By adhering to strict aseptic protocols, employing minimally invasive surgical techniques, and fostering clear communication within the dental team, the incidence of complications such as alveolar osteitis, surgical site infections, and prolonged hemorrhage can be drastically minimized. Future directions in dental practice should emphasize the continuous education of the entire dental team and the adoption of advanced technologies, such as piezosurgery and CBCT, to further enhance the safety and predictability of tooth extractions.

REFERENCES

[1] Bui CH, Seldin EB, Goggin N, et al. Rates of complications after mandibular third-molar extraction. J Oral Maxillofac Surg. 2003;61(11):1283-1289.

[2] Osborn TP, Frederickson GR, Small IA, Torgerson TS. A prospective study of complications related to mandibular third molar extraction. J Oral Maxillofac Surg. 1985;43(10):767-769.

[3] Phillips C, White RP Jr, Shugars DA, Zhou X. Risk factors associated with pain, swelling, and trismus following surgical removal of mandibular third molars. Int J Oral Maxillofac Surg. 2003;32(1):50-57.

[4] Harnett JC, Hughes FJ, Smith AJ, et al. A prospective survey of postoperative pain and analgesic consumption following third molar removal. Int J Oral Maxillofac Surg. 2008;37(8):715-720.

[5] Arteagoitia MI, Barbier L, Santamaria J, et al. Amoxicillin/clavulanic acid to prevent infectious complications after mandibular third molar surgery. Int J Oral Maxillofac Surg. 2008;37(11):1005-1010.

[6] López-Cedrún JL, Pijoan E, Gómez B, et al. Amoxicillin/clavulanic acid to prevent infectious complications after mandibular third molar surgery. Int J Oral Maxillofac Surg. 2005;34(8):839-843.

[7] Sánchez A, García A, López-Cedrún JL, et al. Efficacy of clindamycin or amoxicillin/clavulanic acid in preventing infectious complications after mandibular third molar extraction. J Oral Maxillofac Surg. 2010;68(8):1799-1805.

[8] Halpern LR, Dodson TB. Prophylactic antibiotics for exchange of third molars. Cochrane Database Syst Rev. 2013;(2):CD004152.

[9] Lodi G, Figliuzzi L, Tada A, et al. Antibiotics to prevent complications after tooth extractions. Cochrane Database Syst Rev. 2012;(2):CD003811.

[10] Romero-Sanchiz A, Romero-Ruiz MM, Gutiérrez-Pérez JL, et al. Prophylactic antibiotics in third molar surgery. J Oral Maxillofac Surg. 2006;64(9):1376-1381.

[11] Poeschl PW, Eckel D, Poeschl J, et al. A prospective randomized split-mouth study of the efficacy of amoxicillin/clavulanic acid in preventing infectious complications after mandibular third molar surgery. Int J Oral Maxillofac Surg. 2004;33(7):661-664.

[12] Bowe DC, Rogers SN, Stassen LF, Brennan PA. The incidence of dry socket after tooth extraction. Br J Oral Maxillofac Surg. 2014;52(8):714-719.

[13] Blum IR. Contemporary views on dry socket (alveolar osteitis): a clinical appraisal of standardization, aetiopathogenesis and diagnosis: a critical review. Int J Oral Maxillofac Surg. 2002;31(3):309-317.

[14] Sweet JB, Butler DP. The relationship of preoperative and postoperative factors to the incidence of alveolar osteitis following third molar surgery. J Oral Surg. 1979;37(4):243-248.

[15] Cillo JE Jr, Faries MW, Fontenot DJ, et al. Analysis of the incidence of dry socket after the use of a bioabsorbable packing material. J Oral Maxillofac Surg. 2006;64(11):1581-1584.

[16] Sanfilippo TM, Yates IE, Beatty WT, Gutmann JL. The use of bioabsorbable packing in the treatment of dry socket. J Endod. 1999;25(2):98-100.

[17] Kohn WG, Collins AS, Cleveland JL, et al. Guidelines for infection control in dental health-care settings—2003. MMWR Recomm Rep. 2003;52(RR-17):1-61.

[18] Rutala WA, Weber DJ. Disinfection and sterilization in dental facilities. Dent Clin North Am. 2004;48(3):591-609.

[19] Miller CH, Palenik CJ. Infection control and management of hazardous materials for the dental team. 5th ed. Elsevier; 2014.

[20] Cleveland JL, Gray SK, Kohn WG, et al. The connection between infection control, sterilization, and reprocessing, and health care. J Am Dent Assoc. 2013;144(2):176-184.

[21] Murphy DR. The dental assistant's role in infection control. Dent Assist. 2005;74(3):42-46.

[22] Field EA, Tipton DA. The dental assistant's role in patient assessment. Dent Assist. 2008;77(2):24-30.

[23] Pallasch TJ. Sterilization and disinfection of dental equipment. Dent Clin North Am. 2004;48(3):543-558.