Natural Resources for Human Health
eISSN: 2583-1194

Original Research

Received 12 Oct 2025
Revised 18 Nov 2025
Accepted 25 Dec 2025

Natural Resources for Human Health 2025, Vol. 5 (4): 1327–1346

KEYWORDS:
Orthostatic hypotension, nurse-led interventions, patient mobilization, nursing assessment, orthostatic vital signs, fall prevention

Effectiveness of Nurse-Led Interventions in Preventing Orthostatic Hypotension During Patient Mobilization

Nasser Shabab Alharbi[1], Ghurub Naif Huraysh Alruwaili[2], Albandri Madhi Sameer Alanazi[3], Reem Ghazi Alanazi[4], Shuaa Awadh Mohmed Alanazi[5], Ayed Mahdi Saleh Alanazi[6], Samah Obid Omar Altraici[7], Haya Mater Alanazi[8], Moniah Ibrahim Alrefay[9], Dalal Saud D Al Shammari[10]

[1] Nursing Technician — Madinah Health Cluster – Ministry of Health – Madinah – Madinah Region – Saudi Arabia. Email: nalmdwy035@gmail.com
[2] Nursing Technician — Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia. Email: G.k7@outlook.sa
[3] Nursing Technician — Mental Health Hospital – Al Jouf Health Cluster – Ministry of Health – Al Qurayyat – Al Jouf Region – Saudi Arabia. Email: aalanize@moh.gov.sa
[4] Staff Nurse — Branch of the Ministry of Health in Riyadh Region – Ministry of Health – Riyadh – Riyadh Region – Saudi Arabia. Email: trrki-1@hotmail.com
[5] Staff Nurse — Branch of the Ministry of Health in Riyadh Region – Ministry of Health – Riyadh – Riyadh Region – Saudi Arabia. Email: Shoooshawad@gmail.com
[6] Nursing Technician — Forensic Medicine Department, Northern Borders Forensic Medical Services Center, Branch of the Ministry of Health in the Northern Borders Region – Ministry of Health – Arar – Northern Borders Region – Saudi Arabia. Email: amalenzy@moh.gov.sa
[7] Nursing Technician — Al Malqa Primary Health Care Center – Third Riyadh Health Cluster – Ministry of Health – Riyadh – Riyadh Region – Saudi Arabia. Email: saltraici@moh.gov.sa
[8] Nursing Specialist — Al-Qurayyat General Hospital – Al Jouf Health Cluster – Ministry of Health – Al Qurayyat – Al Jouf Region – Saudi Arabia. Email: Hhaya_39@hotmail.com
[9] Nursing Specialist — Al-Suraif Primary Health Care Center – Madinah Health Cluster – Ministry of Health – Yanbu – Madinah Region – Saudi Arabia. Email: Malrefay@moh.gov.sa
[10] Nursing — Al Jouf Health Cluster – Ministry of Health – Sakaka – Al Jouf Region – Saudi Arabia. Email: Dalshamrry@moh.gov.sa

ABSTRACT: Orthostatic hypotension (OH) is a clinically significant cardiovascular condition characterized by an abnormal reduction in blood pressure following a transition from a lying or sitting position to an upright posture. It represents an important patient safety concern during mobilization, particularly among hospitalized individuals, older adults, critically ill patients, and patients recovering from surgical procedures or prolonged immobilization. The occurrence of orthostatic hypotension during mobilization may result in dizziness, weakness, blurred vision, impaired balance, syncope, falls, and interruption of essential rehabilitation activities. These complications can negatively affect functional recovery, prolong hospitalization, increase healthcare costs, and compromise patient safety. Nurses play a central role in preventing orthostatic hypotension through early identification

of susceptible patients, systematic assessment of orthostatic vital signs, evaluation of hydration status, medication monitoring, gradual positional transitions, appropriate mobilization techniques, and continuous observation of physiological responses. Nurse-led interventions also include patient education, application of prescribed compression garments, encouragement of appropriate fluid intake, implementation of physical counter-maneuvers, individualized activity planning, and collaboration with multidisciplinary healthcare teams. The effectiveness of these interventions depends on their appropriate selection, timing, consistency, and adaptation to individual patient characteristics. Evidence suggests that combining non-pharmacological interventions with standardized nursing assessment and supervised mobilization can improve orthostatic tolerance and facilitate safer early mobilization. However, variations in patient populations, underlying medical conditions, intervention protocols, and outcome measurements highlight the need for individualized evidence-based nursing practices.

INTRODUCTION

Patient mobilization is an essential component of contemporary nursing practice and represents a fundamental intervention for maintaining physiological function, preventing complications associated with immobility, and promoting recovery among hospitalized individuals. Early mobilization involves the progressive transition of patients from bed rest to sitting, standing, transferring, and walking according to their clinical condition and functional abilities. It is particularly important in intensive care units, postoperative recovery settings, medical-surgical wards, and rehabilitation departments, where prolonged immobilization may contribute to muscle weakness, impaired cardiovascular regulation, venous thromboembolism, pulmonary complications, pressure injuries, and functional decline [1]. Although mobilization provides substantial clinical benefits, it also introduces physiological challenges, particularly in patients whose cardiovascular systems cannot adequately compensate for the redistribution of blood that occurs when changing body position. Among these challenges, orthostatic hypotension is one of the most important conditions that may interfere with safe mobilization and increase the likelihood of adverse events during nursing care [1,2].

Orthostatic hypotension is generally defined as a sustained decrease in systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg within three minutes of standing or assuming an upright position following a period of recumbency. This definition has been established through international consensus recommendations and is widely applied in clinical assessment and research. The condition may occur independently or in association with several underlying disorders, including autonomic dysfunction, cardiovascular diseases, diabetes mellitus, neurological disorders, dehydration, anemia, and medication-related adverse effects. Importantly, orthostatic hypotension is not exclusively a disease affecting older adults; it may also occur in younger hospitalized patients, particularly following surgery, prolonged bed rest, acute illness, or significant fluid loss. Its clinical importance becomes especially evident during mobilization because the transition to an upright position imposes an immediate hemodynamic challenge that requires coordinated cardiovascular and autonomic responses [1,3].

Under normal physiological conditions, standing causes gravitational redistribution of blood toward the lower extremities and splanchnic circulation, temporarily reducing venous return to the heart and consequently decreasing ventricular filling and cardiac output. Baroreceptors located in the carotid sinus and aortic arch detect the reduction in arterial pressure and initiate compensatory responses through increased sympathetic activity and reduced parasympathetic activity. These responses promote peripheral vasoconstriction, increased heart rate, and enhanced cardiac contractility, thereby maintaining adequate cerebral and systemic perfusion. In individuals with impaired autonomic regulation, reduced circulating blood volume, or diminished cardiovascular reserve, these compensatory mechanisms may be insufficient. Consequently, blood pressure decreases excessively during postural changes, potentially producing cerebral hypoperfusion and characteristic symptoms such as dizziness, visual disturbances, weakness, nausea, and fainting. The severity of these manifestations varies according to the magnitude and duration of the blood pressure reduction, the patient's underlying physiological condition, and the effectiveness of compensatory cardiovascular mechanisms [2,4].

The relationship between orthostatic hypotension and patient mobilization is particularly important in acute healthcare environments, where patients frequently experience several simultaneous risk factors. Surgical patients may develop orthostatic intolerance because of perioperative fluid shifts, blood loss, anesthetic effects, postoperative pain, inflammatory responses, and

medication exposure. Similarly, critically ill patients may experience cardiovascular instability because of systemic infection, vasopressor withdrawal, sedation, mechanical ventilation, reduced physical activity, and prolonged recumbency. Older hospitalized individuals may be especially vulnerable because of age-related changes in baroreceptor sensitivity, reduced vascular elasticity, impaired autonomic responses, multiple chronic diseases, and polypharmacy. These factors can interact, making the physiological response to mobilization unpredictable and increasing the importance of individualized nursing assessment before initiating upright activity [5,6].

Clinical evidence demonstrates that orthostatic intolerance is a frequent barrier to postoperative mobilization. In a prospective study involving patients undergoing total hip arthroplasty, approximately 22% experienced orthostatic intolerance that prevented successful mobilization on the first postoperative day. Female sex, postoperative gabapentin administration, and higher pain levels during mobilization were identified as important associated factors. Similarly, research involving patients recovering from abdominal surgery demonstrated that orthostatic intolerance occurred frequently during very early postoperative mobilization, particularly within the first hours after surgery. These findings illustrate that apparently stable patients may develop clinically important symptoms when transitioning from bed rest to an upright position, emphasizing the need for structured nursing supervision rather than relying exclusively on resting vital signs [5,7].

Orthostatic hypotension is also associated with important consequences beyond transient discomfort. Affected patients may experience falls, traumatic injuries, syncope, reduced confidence in movement, delayed rehabilitation, and decreased participation in activities of daily living. In older hospitalized populations, orthostatic blood pressure abnormalities are frequently identified during clinical assessment, although the relationship between documented orthostatic hypotension and inpatient falls is influenced by multiple additional factors. Moreover, recurrent episodes may contribute to reduced mobility because patients become reluctant to stand or walk after experiencing dizziness or near-syncope. This creates a potentially harmful cycle in which fear of symptoms leads to reduced activity, prolonged inactivity contributes to physical deconditioning, and deconditioning further reduces cardiovascular tolerance during subsequent mobilization [8,9].

In this context, nursing care represents a central component of prevention because nurses are continuously involved in monitoring patients, assisting with positional changes, evaluating symptoms, administering prescribed treatments, and coordinating daily mobilization activities. Unlike interventions that depend primarily on medical prescribing, several preventive strategies can be incorporated directly into routine nursing practice. These include assessing blood pressure and heart rate before and after positional changes, identifying patients with previous episodes of orthostatic symptoms, ensuring that patients are adequately hydrated when clinically appropriate, assisting with gradual transitions from supine to sitting and standing, and observing for early manifestations of hemodynamic instability. Nurses can also reinforce the use of prescribed compression garments, encourage appropriate lower-limb muscle activation, and provide education regarding safe movement techniques [10,11].

The importance of nurse-led prevention extends beyond the performance of individual interventions. Effective prevention requires a coordinated approach in which patient assessment, clinical decision-making, mobilization planning, monitoring, documentation, and reassessment are integrated into a consistent nursing care process. For example, a patient with a history of orthostatic hypotension, recent diuretic administration, and reduced oral intake requires a different mobilization strategy from a clinically stable patient without recognized risk factors. Similarly, patients recovering from major surgery may require more frequent assessment and closer supervision during their initial attempts to stand and walk. Therefore, nursing interventions should be individualized according to physiological stability, underlying disease, medication exposure, functional capacity, and the patient's response to previous mobilization attempts [10,12].

Non-pharmacological strategies have received considerable attention in the management of orthostatic hypotension because they can be incorporated into daily care and may improve upright blood pressure without necessarily requiring additional medication. These strategies include gradual positional transitions, adequate hydration, physical counter-maneuvers, abdominal compression, lower-limb compression, avoidance of prolonged immobility, and appropriate adjustment of activity schedules. A systematic review of non-pharmacological interventions identified potentially beneficial effects associated with abdominal compression, physical counter-maneuvers, and water intake, although the strength of evidence varied across interventions and patient populations. This evidence supports the integration of appropriate non-pharmacological measures into individualized nursing care plans while recognizing that not every intervention is suitable for every patient [13,14].

Another important consideration is the distinction between preventing orthostatic hypotension and responding to an episode after it has developed. Preventive nursing interventions aim to identify physiological vulnerability before mobilization and reduce the likelihood of a significant blood pressure decrease. In contrast, responsive interventions focus on recognizing symptoms promptly, stopping unsafe activity, assisting the patient into a safe position, reassessing hemodynamic status, and escalating care when necessary. Both approaches are essential because even carefully selected preventive measures cannot eliminate all episodes, particularly in patients with severe autonomic dysfunction or acute cardiovascular instability. The effectiveness of nursing care therefore depends on the combination of prevention, early recognition, immediate management, and continuous evaluation [3,15].

Despite the growing recognition of orthostatic hypotension as a patient safety concern, its prevention during mobilization remains challenging in many clinical settings. Differences in institutional protocols, inconsistent measurement of orthostatic vital signs, limited staff awareness, insufficient documentation, and competing clinical priorities may interfere with the systematic implementation of preventive interventions. Furthermore, evidence derived from outpatient populations cannot always be directly generalized to critically ill or postoperative patients because the underlying causes and physiological responses may differ substantially. These considerations emphasize the need for evidence-based nursing protocols that address the specific circumstances of hospitalized patients and integrate standardized assessment with safe, progressive mobilization [11,16].

The effectiveness of nurse-led interventions should consequently be considered in relation to several complementary outcomes, including the frequency and severity of orthostatic blood pressure reductions, the occurrence of dizziness and presyncope, the ability to complete planned mobilization, the incidence of falls and related injuries, patient tolerance of upright activity, and the achievement of rehabilitation milestones. An intervention that produces a modest improvement in standing blood pressure may still be clinically valuable if it allows a patient to participate safely in essential rehabilitation activities. Conversely, successful mobilization should not be evaluated solely by the ability to stand or walk, but also by the patient's physiological stability, symptom burden, functional independence, and safety throughout the process [6,13].

Accordingly, nurse-led prevention represents an important component of comprehensive patient safety and rehabilitation-oriented care. Through systematic risk assessment, standardized orthostatic monitoring, gradual mobilization, hydration support, medication surveillance, physical counter-maneuvers, compression strategies, and patient education, nurses can reduce preventable hemodynamic challenges and support safer transitions to upright activity. The integration of these interventions into routine nursing practice may improve patient tolerance of mobilization, minimize interruptions to rehabilitation, and contribute to better clinical outcomes across different healthcare environments [10,14,16].

Understanding Orthostatic Hypotension

Definition and Clinical Classification

Orthostatic hypotension is a disorder of blood pressure regulation characterized by an excessive and sustained reduction in arterial pressure following a change from a recumbent or seated position to an upright posture. The internationally accepted diagnostic definition is a reduction in systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg within three minutes of standing or head-up tilt. In patients with significant supine hypertension, a larger systolic reduction may be considered when interpreting the clinical findings. The definition provides a standardized framework for identifying abnormal postural blood pressure responses, although clinical interpretation should also consider the presence of symptoms, baseline blood pressure, timing of measurements, and underlying medical conditions [1,3].

Orthostatic hypotension can be classified according to the timing and underlying mechanism of the blood pressure reduction. Classic orthostatic hypotension develops within three minutes of standing, whereas delayed orthostatic hypotension occurs after more than three minutes in the upright position. Initial orthostatic hypotension involves a transient blood pressure decrease immediately after standing, generally occurring within the first 15 seconds, and may not be detected by conventional intermittent blood pressure measurements. Neurogenic orthostatic hypotension results from impaired autonomic regulation, whereas non-neurogenic forms are frequently associated with volume depletion, medication effects, cardiovascular abnormalities, or acute illness. Understanding these distinctions is important for nurses because a patient may develop symptoms even when routine measurements obtained at a single time point appear relatively normal [3,4].

Pathophysiological Mechanisms

The development of orthostatic hypotension involves an imbalance between gravitational blood redistribution and the body's compensatory cardiovascular responses. During standing, approximately 500–1,000 mL of blood may accumulate in the lower body and abdominal circulation, resulting in reduced venous return and cardiac filling. Under normal circumstances, sympathetic activation increases vascular resistance and heart rate, helping to maintain cerebral perfusion. When these mechanisms are impaired, arterial pressure may decrease excessively, reducing blood flow to the brain and producing symptoms associated with orthostatic intolerance. The degree of impairment may vary according to autonomic function, circulating blood volume, vascular responsiveness, cardiac performance, and the patient's physical condition [4,17].

Several mechanisms can contribute to this abnormal response. Autonomic failure may impair vasoconstriction and appropriate heart rate regulation, while dehydration and blood loss reduce circulating volume and venous return. Prolonged bed rest can decrease plasma volume, reduce lower-limb muscle strength, and impair cardiovascular conditioning. In addition, medications such as diuretics, vasodilators, alpha-adrenergic blockers, and certain antidepressants may aggravate postural blood pressure reductions. These mechanisms frequently coexist in hospitalized patients, explaining why orthostatic hypotension may develop even in individuals without a previous history of the condition [10,18].

Clinical Manifestations

The clinical manifestations of orthostatic hypotension range from mild and transient symptoms to severe episodes associated with syncope and falls. Common symptoms include dizziness, lightheadedness, blurred vision, weakness, fatigue, nausea, palpitations, headache, and difficulty maintaining balance after standing. Some patients experience cognitive slowing, impaired concentration, or a sensation of impending fainting, while others may demonstrate a measurable blood pressure decrease without reporting symptoms. Symptom recognition is particularly important during mobilization because patients may not immediately communicate discomfort, especially when they are elderly, cognitively impaired, sedated, or recovering from anesthesia [3,4].

The timing of symptoms provides important clinical information. Symptoms that develop immediately after standing may suggest initial orthostatic hypotension, whereas symptoms occurring within the first three minutes may indicate classic orthostatic hypotension. Delayed symptoms may develop after prolonged standing or walking, highlighting the importance of continued observation beyond the initial transition. Nurses should therefore assess both objective blood pressure changes and subjective patient experiences throughout mobilization rather than relying exclusively on a single blood pressure measurement [3,19].

Epidemiology and Clinical Significance of Orthostatic Hypotension During Mobilization

Orthostatic hypotension is a relatively common clinical finding among hospitalized patients, although its prevalence varies considerably according to age, underlying disease, clinical setting, diagnostic criteria, and timing of assessment. Older adults are particularly susceptible because of age-related autonomic changes, reduced cardiovascular reserve, impaired baroreceptor sensitivity, and the frequent coexistence of chronic illnesses. In acute-care settings, the condition may also develop transiently in patients who have experienced surgery, infection, dehydration, blood loss, or prolonged immobilization. Its identification is therefore important not only in patients with previously diagnosed autonomic disorders but also in individuals undergoing routine nursing-assisted mobilization [8,9,20].

A retrospective investigation involving 7,745 hospitalized older adults who underwent orthostatic vital sign assessment identified orthostatic hypotension in 39.2% of patients during the first recorded assessment. Although this investigation did not establish a statistically significant association between documented orthostatic hypotension and inpatient falls, the high prevalence demonstrates the importance of systematic assessment in older hospitalized populations. These findings also indicate that orthostatic hypotension should be interpreted within the broader context of individual fall risk, including mobility impairment, cognitive status, medication exposure, environmental hazards, and previous falls [8].

Postoperative patients constitute another important population because early mobilization is increasingly incorporated into enhanced recovery pathways. However, postoperative orthostatic intolerance may interfere with the ability to sit, stand, or walk during the initial recovery period. In abdominal surgery patients, orthostatic intolerance was particularly frequent approximately three hours after surgery, with a substantial reduction in its occurrence by the following day. This temporal pattern suggests that the first mobilization attempt represents a particularly important opportunity for nursing assessment and preventive

intervention. It also emphasizes the importance of considering the patient's recovery stage rather than applying identical mobilization expectations throughout the postoperative period [5,7].

Risk Factors Associated with Orthostatic Hypotension During Patient Mobilization

Advanced Age and Physiological Changes

Advanced age is an important risk factor because the cardiovascular and autonomic systems undergo progressive physiological changes that may reduce the ability to maintain arterial pressure during postural transitions. Reduced baroreceptor sensitivity, impaired vascular responsiveness, diminished thirst perception, and decreased renal concentrating ability can contribute to abnormal blood pressure regulation. Furthermore, older adults frequently have multiple chronic conditions and receive several medications that may influence cardiovascular stability. Consequently, even relatively minor changes in hydration, medication administration, or physical activity may precipitate symptomatic orthostatic hypotension during mobilization [9,20].

Prolonged Bed Rest and Physical Deconditioning

Prolonged bed rest contributes to cardiovascular deconditioning through reductions in circulating plasma volume, lower-limb muscle strength, and orthostatic tolerance. The skeletal muscles of the lower extremities normally contribute to venous return through rhythmic contractions that facilitate blood movement toward the heart. When patients remain inactive for extended periods, this muscle-pump mechanism becomes less effective, increasing venous pooling when they eventually assume an upright position. Reduced physical conditioning may also impair the ability to tolerate sustained standing and walking. Therefore, prolonged immobilization represents an important modifiable risk factor that should be considered when planning progressive mobilization [9,17].

Dehydration and Reduced Circulating Blood Volume

Inadequate fluid intake, excessive fluid losses, vomiting, diarrhea, fever, bleeding, and diuretic administration may reduce intravascular volume and increase susceptibility to orthostatic hypotension. A reduced circulating volume limits venous return during standing and may prevent the cardiovascular system from maintaining sufficient cardiac output. Hospitalized patients may be particularly vulnerable because their oral intake can be restricted by diagnostic procedures, postoperative nausea, swallowing difficulties, altered consciousness, or prescribed fluid restrictions. Nurses should therefore monitor fluid balance, assess clinical indicators of dehydration, and identify patients whose fluid requirements require medical review before mobilization [10,18].

Medication-Related Factors

Medication exposure represents a major contributor to orthostatic hypotension among hospitalized patients, particularly those receiving multiple cardiovascular or centrally acting medications. Diuretics may reduce circulating blood volume, whereas vasodilators and alpha-adrenergic blockers can reduce vascular resistance. Certain antidepressants, antipsychotic medications, sedatives, and medications used for neurological disorders may also interfere with autonomic regulation or cardiovascular compensation. Nurses should review medication administration records, identify recent changes in therapy, monitor blood pressure in relation to medication timing, and communicate suspected adverse effects to the prescribing healthcare professional. Medication modification must remain a collaborative clinical decision rather than an independent nursing decision [10,11,18].

Cardiovascular and Neurological Disorders

Patients with heart failure, arrhythmias, ischemic heart disease, autonomic neuropathy, Parkinsonian disorders, and other neurological conditions may have reduced cardiovascular reserve or impaired autonomic regulation. Diabetes mellitus may contribute to autonomic neuropathy, while neurodegenerative diseases can interfere with the normal sympathetic responses required to maintain blood pressure during standing. In such patients, orthostatic hypotension may occur repeatedly despite adequate hydration and careful mobilization. Nurses should therefore consider underlying disease mechanisms when evaluating recurrent episodes and communicate persistent abnormalities for further medical investigation [4,20].

Postoperative and Critical Illness-Related Factors

Postoperative patients may experience orthostatic hypotension because of anesthesia-related vasodilation, perioperative fluid shifts, blood loss, inflammatory responses, postoperative pain, and exposure to analgesic medications. Pain may further complicate mobilization by increasing sympathetic stimulation and limiting the patient's ability to perform gradual movements.

In critically ill patients, systemic inflammation, sepsis, prolonged sedation, vasopressor therapy, and impaired cardiovascular regulation may produce additional instability. These factors require close coordination between nursing assessment and the broader clinical plan before initiating or advancing mobilization [5,6,7].

Nutritional and Hematological Factors

Anemia may contribute to exercise intolerance, fatigue, weakness, and reduced oxygen delivery to tissues, potentially aggravating the clinical consequences of an orthostatic blood pressure decrease. Nutritional deficiencies and inadequate caloric intake may also contribute to generalized weakness and impaired functional recovery. Although anemia does not independently establish the diagnosis of orthostatic hypotension, it is an important contributing factor when evaluating patients who develop dizziness, weakness, or intolerance during mobilization. Nurses should monitor relevant laboratory results, observe for clinical manifestations of anemia, and communicate abnormalities that may affect rehabilitation tolerance [9,20].

Table 1. Major risk factors for orthostatic hypotension during patient mobilization and their nursing implications [5,9,10,11,18,20].

Risk factor	Potential physiological mechanism	Nursing implication
Advanced age	Reduced baroreceptor sensitivity and autonomic responsiveness	Increased monitoring and supervised mobilization
Prolonged bed rest	Reduced plasma volume and muscle-pump activity	Gradual progressive mobilization
Dehydration	Reduced circulating blood volume	Fluid balance assessment
Blood loss and anemia	Reduced oxygen delivery and cardiovascular reserve	Monitor laboratory findings and clinical symptoms
Diuretic administration	Reduced intravascular volume	Monitor fluid status and medication timing
Vasodilator therapy	Reduced peripheral vascular resistance	Monitor blood pressure and report symptomatic reductions
Neurological disorders	Impaired autonomic regulation	Individualized mobilization and symptom surveillance
Postoperative recovery	Anesthesia effects, fluid shifts, and pain	Supervised first mobilization
Sedation and analgesics	Altered autonomic responses and impaired awareness	Assess alertness and hemodynamic stability
Heart failure	Limited cardiac reserve and possible fluid-management complexity	Individualized fluid and mobilization plans

Comprehensive Nursing Assessment Before Patient Mobilization

Initial Patient Evaluation

Comprehensive nursing assessment is the foundation of an effective prevention strategy because it enables the identification of patients who may experience an abnormal cardiovascular response during positional changes. Before mobilization, nurses should review the patient's medical history, current diagnosis, previous episodes of dizziness or syncope, cardiovascular and neurological conditions, medication profile, recent surgical procedures, fluid balance, and functional capacity. Particular attention should be directed toward patients with previous falls, prolonged immobilization, recent blood loss, reduced oral intake, or documented orthostatic hypotension. The assessment should also establish whether the patient can understand instructions, maintain balance, and communicate symptoms during the mobilization process [10,11].

Natural Resources for Human Health
eISSN: 2583-1194

Assessment of Baseline Vital Signs

Baseline blood pressure and heart rate measurements provide important information about cardiovascular stability before mobilization. When clinically appropriate, blood pressure should be measured after the patient has rested in the supine position for approximately five minutes. Subsequent measurements should be obtained after standing, generally at one and three minutes, while documenting associated symptoms. If standing is unsafe, an individualized alternative involving gradual head elevation or supported sitting may be considered according to clinical judgment. The same arm, appropriate cuff size, and consistent measurement technique should be used whenever possible to improve reliability and comparability [3,16].

A standardized measurement approach is particularly important because isolated readings may fail to identify certain patterns of orthostatic hypotension. For example, initial orthostatic hypotension may develop within seconds of standing and resolve before a conventional blood pressure reading is obtained. Delayed orthostatic hypotension may occur after the initial three-minute observation period. Therefore, nurses should interpret measurements alongside the patient's symptoms and clinical history, extending observation when the clinical presentation suggests an abnormal postural response [3,4].

Assessment of Orthostatic Symptoms

Symptom assessment should be performed before, during, and after mobilization. Nurses should ask patients about dizziness, visual changes, nausea, weakness, palpitations, headache, and sensations of impending fainting. Observation is equally important in patients who cannot communicate effectively or who have cognitive impairment. Changes in facial expression, reduced responsiveness, unsteadiness, pallor, sweating, or sudden weakness may indicate intolerance to upright activity. Symptoms should be documented with their onset, duration, severity, associated position, and response to returning to a safe posture [3,11].

Assessment of Hydration and Fluid Balance

Evaluation of hydration status should include oral intake, intravenous fluid administration, urine output, ongoing fluid losses, and prescribed fluid restrictions. Nurses should consider whether the patient has experienced vomiting, diarrhea, fever, excessive sweating, or bleeding. Clinical findings such as dry mucous membranes, reduced urine output, and postural symptoms may support the suspicion of volume depletion, although these findings should not be interpreted in isolation. In patients with heart failure, renal impairment, or other conditions associated with fluid overload, hydration interventions must be individualized and coordinated with the treating team [10,18].

Medication Assessment

Medication reconciliation and ongoing medication surveillance are essential because pharmacological agents may contribute to abnormal postural blood pressure responses. Nurses should identify medications that may reduce circulating volume, decrease vascular resistance, or interfere with autonomic compensation. The timing of administration should be considered in relation to mobilization attempts, especially when patients report recurrent symptoms after receiving specific medications. Any suspected medication-related problem should be communicated to the prescribing clinician, with appropriate documentation of the associated blood pressure changes and clinical manifestations [10,11].

Table 2. Standardized nursing assessment of patients at risk of orthostatic hypotension [3,4,10,11,16].

Assessment component	Nursing assessment	Clinical significance
Medical history	Cardiovascular, neurological, and metabolic disorders	Identifies underlying susceptibility
Previous episodes	History of dizziness, syncope, or falls	Identifies recurrent postural intolerance
Baseline blood pressure	Supine blood pressure after rest	Establishes baseline hemodynamic status
Standing blood pressure	Measurements at one and three minutes	Identifies postural blood pressure reduction
Heart rate	Supine and upright heart rate	Helps evaluate compensatory response
Symptoms	Dizziness, blurred vision, weakness, nausea	Identifies symptomatic intolerance

Hydration	Intake, output, and fluid losses	Identifies possible volume depletion
Medications	Diuretics, vasodilators, and other relevant agents	Identifies potentially modifiable contributors
Functional status	Balance, strength, and ability to follow instructions	Determines the appropriate level of assistance
Laboratory findings	Hemoglobin, electrolytes, and renal function when available	Identifies associated clinical abnormalities

Nurse-Led Interventions for Preventing Orthostatic Hypotension During Mobilization

Gradual and Staged Positional Changes

Gradual positional transition is one of the most fundamental nursing interventions for preventing orthostatic hypotension during mobilization. Abrupt movement from a prolonged supine position to standing can produce rapid gravitational redistribution of blood before compensatory cardiovascular responses have become adequate. Nurses should therefore assist patients through progressive stages, beginning with appropriate positioning in bed, followed by sitting upright, dangling the legs over the bedside, standing with assistance, and eventually walking when tolerated. Each stage should provide sufficient time for the patient to demonstrate hemodynamic stability before proceeding to the next stage [10,17].

During the transition from lying to sitting, nurses should observe the patient's facial expression, level of alertness, balance, and reported symptoms. If the patient tolerates sitting without dizziness or other concerning manifestations, standing may be attempted with appropriate assistance and safety equipment. Patients who develop symptoms should not be encouraged to continue standing merely to complete a planned activity. Instead, mobilization should be paused, the patient should be assisted into a safe position, and the clinical assessment should be repeated. This individualized approach allows mobilization to progress according to physiological tolerance rather than predetermined time schedules alone [10,11].

Standardized Orthostatic Blood Pressure Monitoring

Routine assessment of orthostatic vital signs is an important preventive intervention, particularly among patients with recognized risk factors or previous episodes of postural intolerance. Nurses should establish baseline blood pressure and heart rate measurements and repeat them after positional changes according to institutional protocols and the patient's clinical condition. A reduction meeting the diagnostic threshold should prompt clinical evaluation, while symptoms such as severe dizziness, near-syncope, or sudden weakness require immediate attention even if the measured blood pressure reduction does not reach the formal diagnostic threshold [1,3].

The effectiveness of standardized monitoring depends on the quality and consistency of the assessment process. Accurate measurements require appropriate cuff placement, adequate rest before baseline assessment, correct positioning, and documentation of the timing of each reading. Nurses should also recognize that normal blood pressure measurements obtained while supine do not necessarily indicate that the patient can safely tolerate standing. Incorporating orthostatic assessment into the pre-mobilization nursing routine may therefore help identify vulnerable patients before an adverse event occurs [3,16].

Hydration Support and Fluid Management

Adequate hydration can help maintain circulating blood volume and improve the cardiovascular response to standing. Nurses can contribute to prevention by monitoring fluid intake and output, identifying patients with inadequate oral intake, encouraging permitted fluid consumption, and reporting signs of volume depletion. In selected patients with orthostatic hypotension, rapid ingestion of approximately 480–500 mL of water may temporarily improve standing blood pressure through a physiological pressor response. However, this strategy should not be applied universally because patients with heart failure, advanced kidney disease, dysphagia, or prescribed fluid restrictions may require different fluid-management approaches [10,13].

Hydration support should be individualized and integrated with the overall treatment plan. Nurses should assess whether the patient can safely swallow fluids, determine whether oral intake is appropriate, and coordinate intravenous fluid administration with medical orders. Fluid administration should not be used as a substitute for investigating ongoing bleeding, infection,

excessive fluid loss, or other underlying causes. Its effectiveness is greatest when incorporated into a broader strategy that includes medication review, symptom assessment, and progressive mobilization [10,18].

Compression Garments and Abdominal Binders

Compression garments are used to reduce venous pooling in the lower extremities and improve venous return during upright activity. Graduated compression stockings may be prescribed for selected patients, while abdominal binders can reduce blood pooling within the splanchnic circulation. The effectiveness of compression depends on the type of garment, appropriate fitting, correct application, patient tolerance, and the underlying mechanism of orthostatic hypotension. Evidence suggests that abdominal compression may provide a meaningful improvement in standing blood pressure in selected patients, although its long-term effectiveness and acceptability vary [13,14].

Nurses are responsible for checking prescribed compression devices, ensuring that they are applied correctly, assessing skin integrity, and monitoring for discomfort or complications. Garments should not be applied indiscriminately, particularly in patients with significant peripheral arterial disease, fragile skin, or other contraindications. The choice between lower-limb compression and abdominal compression should be guided by the clinical plan and individual patient characteristics. Compression should be considered an adjunct to, rather than a replacement for, careful mobilization and hemodynamic monitoring [13,14].

Physical Counter-Maneuvers

Physical counter-maneuvers involve deliberate contraction of selected skeletal muscle groups to improve venous return and temporarily support arterial pressure. Examples include lower-limb muscle tensing, leg crossing, buttock contraction, and other appropriate isometric movements. These techniques can be particularly useful in patients who experience early warning symptoms during standing and who have sufficient strength, balance, and cognitive ability to perform them safely. Their effectiveness is related to increased muscle-pump activity and reduced venous pooling [13,17].

Nurses can teach appropriate counter-maneuvers during routine patient education and reinforce their use before standing or when early symptoms develop. However, the selected maneuver should be appropriate for the patient's functional condition. Patients with severe weakness, impaired balance, postoperative movement restrictions, or significant neurological impairment may require alternative strategies. Counter-maneuvers should never delay assistance when a patient is experiencing severe presyncope or is at immediate risk of falling [11,13].

Avoidance of Prolonged Immobilization

Avoiding unnecessary prolonged bed rest is important because physical inactivity contributes to cardiovascular deconditioning and reduced orthostatic tolerance. Nurses should collaborate with the multidisciplinary team to establish an individualized mobility schedule that progresses from bed exercises to sitting, standing, transfers, and walking according to clinical stability. Regular changes in position and appropriate activity may help maintain lower-limb muscle function and support cardiovascular adaptation. Nevertheless, mobilization should be postponed or modified when the patient demonstrates acute instability or has a clinical contraindication [9,10].

Progressive mobilization is particularly important for patients recovering from surgery and critical illness. Initial activities may include active or assisted range-of-motion exercises, supported sitting, and brief standing periods before longer walking sessions are introduced. Nurses should evaluate the response to each activity and modify subsequent sessions according to blood pressure, heart rate, symptoms, fatigue, and functional performance. This approach reduces the risk of overwhelming patients who have become physically deconditioned during hospitalization [6,7].

Medication Surveillance and Interprofessional Collaboration

Medication surveillance is an essential component of nurse-led prevention because medication-related orthostatic hypotension may be modifiable through appropriate clinical review. Nurses should identify medications that may contribute to reduced blood pressure or circulating volume and report clinically significant findings to physicians and pharmacists. Particular attention should be given to recent medication initiation, dose escalation, administration before mobilization, and combinations of medications with additive hypotensive effects. Medication adjustment must be individualized because unnecessary discontinuation of prescribed cardiovascular therapy may create additional clinical risks [10,11].

Interprofessional collaboration is especially important when orthostatic hypotension persists despite routine nursing interventions. Physicians may need to investigate underlying cardiovascular or neurological causes, while pharmacists can support medication review and physiotherapists can assist with functional assessment and progressive rehabilitation. Nurses contribute by providing continuous clinical observations and communicating changes in patient tolerance. This coordinated approach helps ensure that preventive interventions address both immediate mobilization-related risks and underlying medical contributors [11,15].

Patient and Family Education

Patient education is an important preventive intervention because patients who understand the symptoms and triggers of orthostatic hypotension can participate more safely in mobilization. Nurses should explain the importance of changing positions gradually, requesting assistance before standing, recognizing early warning symptoms, and avoiding sudden movements after prolonged recumbency. Patients should also be instructed to report dizziness, blurred vision, weakness, nausea, or palpitations immediately rather than attempting to continue walking independently. Family members and caregivers should receive similar information when they are involved in assisting with mobility [10,11].

Education should be tailored to the patient's cognitive status, language, health literacy, and functional ability. Demonstration and return demonstration may be more effective than verbal instructions alone, particularly in patients with limited previous experience of safe mobilization. Educational reinforcement should continue throughout hospitalization and be incorporated into discharge planning when orthostatic symptoms are likely to persist after discharge [11,20].

Table 3. Major nurse-led interventions and their expected clinical effects [10,11,13,14,17,18].

Nursing intervention	Proposed mechanism	Expected benefit	Important considerations
Gradual positional transitions	Allows cardiovascular adaptation	Reduces abrupt postural stress	Observe symptoms at every stage
Orthostatic vital sign monitoring	Early detection of abnormal BP responses	Identifies patients requiring modified mobilization	Use standardized measurement technique
Hydration support	Maintains circulating blood volume	May improve upright BP	Consider cardiac and renal restrictions
Compression garments	Reduces peripheral venous pooling	Supports venous return	Ensure correct fitting and skin assessment
Abdominal binders	Reduces splanchnic pooling	May improve standing BP	Assess comfort and contraindications
Physical counter-maneuvers	Activates skeletal muscle pumps	May reduce postural symptoms	Requires adequate strength and balance
Progressive mobilization	Reduces physical deconditioning	Improves functional tolerance	Individualize activity progression
Medication surveillance	Identifies potentially modifiable causes	Reduces medication-related risk	Collaborate with prescribers
Patient education	Improves symptom recognition and safe behavior	Promotes safer independent mobility	Adapt to health literacy
Multidisciplinary coordination	Integrates medical and rehabilitation management	Supports continuity of prevention	Maintain clear documentation

Natural Resources for Human Health
eISSN: 2583-1194

Effectiveness of Nurse-Led Interventions in Preventing Orthostatic Hypotension

Effectiveness of Gradual Mobilization Protocols

Gradual mobilization protocols are clinically important because they allow the cardiovascular system to adapt progressively to changes in body position. By introducing sitting, standing, and walking in sequential stages, nurses can identify early manifestations of orthostatic intolerance before patients are exposed to more demanding physical activity. This strategy is particularly relevant during the first mobilization attempt after surgery or prolonged bed rest, when physiological adaptation may be impaired. Its effectiveness is reflected not only in the prevention of symptomatic blood pressure reductions but also in the ability to complete planned mobility activities without interruption [5,7,17].

The value of staged mobilization is also related to its ability to support functional recovery without exposing patients to unnecessary physiological stress. Patients who tolerate sitting may progress to supported standing, whereas those who develop dizziness or weakness can receive additional assessment and a modified activity plan. This individualized approach is more appropriate than expecting all patients to achieve the same mobilization milestones within identical timeframes. Nevertheless, direct comparative evidence demonstrating the independent effect of gradual mobilization protocols on orthostatic hypotension incidence remains limited, and their effectiveness should be evaluated alongside other preventive interventions [6,7].

Effectiveness of Orthostatic Vital Sign Monitoring

Standardized orthostatic blood pressure assessment is an important intervention because it provides objective information about the cardiovascular response to postural changes. Its primary benefit is early identification of patients who may require additional supervision, modified activity progression, or medical evaluation before mobilization. Measuring blood pressure in the supine and upright positions, together with heart rate and symptom assessment, enables nurses to identify clinically important abnormalities that may not be apparent from resting measurements alone [3,4].

However, the effectiveness of orthostatic monitoring depends on its integration into clinical decision-making. Detecting an abnormal blood pressure response does not independently prevent complications unless appropriate actions follow the assessment. For example, a patient with a significant postural reduction may require additional observation, medication review, hydration assessment, or an alternative mobilization strategy. Therefore, orthostatic vital sign monitoring should be considered part of a comprehensive prevention protocol rather than an isolated diagnostic activity [3,16].

Effectiveness of Hydration Interventions

Hydration interventions can improve orthostatic tolerance by supporting intravascular volume and increasing the physiological reserve available during postural changes. Evidence from clinical investigations indicates that rapid water ingestion may produce a temporary improvement in standing systolic blood pressure in some patients. In one investigation involving older adults, water bolus administration produced a clinically meaningful response in a proportion of participants, supporting its potential role as an adjunctive non-pharmacological intervention [13].

Nevertheless, hydration should not be considered universally effective or appropriate. Patients with heart failure, renal impairment, fluid overload, or prescribed fluid restrictions may experience complications if additional fluids are administered without proper assessment. Furthermore, dehydration may be only one component of a complex clinical problem involving autonomic dysfunction or medication-related hypotension. Consequently, nurse-led hydration interventions are most effective when guided by individualized fluid assessment and integrated with the patient's overall treatment plan [10,13].

Effectiveness of Compression Therapy

Compression therapy may improve upright blood pressure by reducing venous pooling and supporting central blood volume. Abdominal binders are particularly relevant because compression of the abdominal region may reduce blood accumulation in the splanchnic circulation, which represents an important reservoir during standing. Clinical evidence has demonstrated improvements in standing blood pressure with abdominal compression in selected patients, although the magnitude of response varies among individuals [13,14].

The effectiveness of compression garments is influenced by several practical factors, including correct sizing, application technique, duration of use, comfort, and patient adherence. In addition, lower-limb compression stockings may not provide equivalent benefits to abdominal compression in every patient. Nurses therefore need to assess whether the prescribed device

is being used correctly and whether the patient tolerates it during mobilization. Compression therapy should be selected according to clinical indications and should not replace standard hemodynamic assessment [13,14].

Effectiveness of Physical Counter-Maneuvers

Physical counter-maneuvers may improve orthostatic tolerance through active contraction of the lower-extremity and abdominal muscles. These contractions increase venous return and can temporarily support arterial pressure, particularly when performed before or during early manifestations of orthostatic symptoms. Systematic reviews have identified potentially beneficial effects associated with leg crossing, muscle pumping, and abdominal compression, although the available studies vary in methodology and clinical populations [13,14].

A practical advantage of these maneuvers is that they can be taught as part of routine nursing education without requiring specialized equipment. However, their effectiveness depends on the patient's physical and cognitive abilities. Patients with significant weakness, impaired balance, severe neurological impairment, or postoperative movement restrictions may be unable to perform them safely. Therefore, nurses should assess functional suitability before incorporating these techniques into an individualized mobilization plan [11,13].

Effectiveness of Multicomponent Nursing Interventions

Multicomponent interventions may provide greater clinical value than isolated preventive measures because orthostatic hypotension frequently results from several interacting physiological and treatment-related factors. A combined approach may include pre-mobilization risk assessment, standardized orthostatic vital signs, appropriate fluid management, medication surveillance, gradual positional changes, compression therapy when indicated, and continuous symptom monitoring. Such an approach addresses different mechanisms contributing to postural blood pressure reduction and allows preventive care to be adapted to changing clinical circumstances [10,11,18].

The importance of multicomponent care is particularly evident in patients with multiple comorbidities or complex postoperative recovery. For example, a patient who has experienced blood loss, received diuretics, and remained on bed rest may require several simultaneous interventions rather than a single strategy. Nurses can coordinate these measures through individualized care plans and continuous reassessment. Nevertheless, the contribution of each individual intervention should be distinguished whenever possible because the combined effect of a protocol does not automatically establish the independent effectiveness of every component [13,15].

Effectiveness in Preventing Falls and Improving Mobilization Outcomes

The prevention of orthostatic hypotension may contribute to safer mobilization by reducing episodes of dizziness, weakness, and presyncope that can interfere with standing and walking. However, falls are multifactorial events, and not every fall is directly attributable to an orthostatic blood pressure reduction. Effective nursing prevention must therefore integrate orthostatic assessment with other fall-prevention measures, including assistance during transfers, appropriate footwear, environmental safety, assessment of cognitive status, and individualized supervision [8,16].

Successful prevention should also be evaluated in relation to rehabilitation participation. Patients who tolerate mobilization more effectively may be better able to participate in physiotherapy, regain functional independence, and progress toward discharge readiness. This is particularly important after major surgery, when delayed mobilization may interfere with recovery pathways. However, the available evidence does not justify attributing improvements in length of stay or functional outcomes exclusively to nurse-led orthostatic hypotension interventions, because these outcomes are also influenced by surgical complexity, disease severity, pain management, and overall quality of care [5,6,7].

Comparison of Evidence Supporting Different Preventive Interventions

The available literature demonstrates that different interventions target different mechanisms contributing to orthostatic hypotension. Some strategies primarily improve venous return, whereas others support circulating blood volume or reduce the physiological stress associated with postural changes. Their relative effectiveness varies according to patient characteristics, clinical setting, and the underlying cause of orthostatic hypotension.

Table 4. Comparative evidence regarding non-pharmacological interventions for orthostatic hypotension [3,8,13,14,17].

Intervention	Main physiological effect	Evidence-based interpretation	Nursing applicability
Water bolus	Temporary increase in blood pressure through a pressor response	Beneficial response demonstrated in selected older adults	Appropriate after fluid suitability assessment
Abdominal compression	Reduces splanchnic venous pooling	Demonstrated improvement in standing systolic BP	Useful when prescribed and tolerated
Compression stockings	Reduces lower-extremity venous pooling	Effects vary and may be less pronounced than abdominal compression	Requires correct fitting
Physical counter-maneuvers	Enhances muscle-pump activity	May improve postural blood pressure responses	Requires adequate physical ability
Gradual mobilization	Reduces abrupt postural stress	Important clinical safety strategy	Easily incorporated into routine nursing care
Orthostatic BP monitoring	Detects abnormal postural responses	Strong clinical value for recognition and assessment	Requires standardized measurement
Avoidance of prolonged bed rest	Reduces physical deconditioning	Supports longer-term orthostatic tolerance	Requires individualized activity planning
Patient education	Improves recognition and response to symptoms	Important supportive component of prevention	Suitable across clinical settings

Nurse-Led Mobilization Protocol for Patients at Risk of Orthostatic Hypotension

A standardized mobilization protocol provides a practical framework for translating clinical evidence into routine nursing practice. Such a protocol should begin with an assessment of the patient's readiness for activity, followed by appropriate preparation, progressive positional changes, continuous observation, and post-mobilization reassessment. The protocol must remain flexible enough to accommodate individual clinical circumstances, including postoperative restrictions, cardiovascular instability, neurological impairment, and prescribed fluid limitations [10,11].

Pre-Mobilization Phase

Before initiating mobilization, the nurse should review the patient's medical history, recent vital signs, medication administration, fluid balance, and previous response to positional changes. Patients with previous orthostatic symptoms, significant blood loss, dehydration, prolonged bed rest, or medications associated with hypotension should receive particular attention. The nurse should ensure that the environment is safe, necessary mobility aids are available, and adequate assistance is provided. When indicated, baseline supine blood pressure and heart rate should be recorded before proceeding [3,10].

Initial Positioning Phase

The patient should be assisted into a supported sitting position, allowing sufficient time for physiological adaptation. During this stage, the nurse should assess dizziness, weakness, visual changes, nausea, and other symptoms associated with reduced cerebral perfusion. Blood pressure and heart rate should be reassessed according to the patient's risk profile and institutional protocol. Patients who demonstrate adequate tolerance may progress toward sitting at the bedside, while those experiencing symptoms should remain supported and undergo further assessment [3,11].

Standing Phase

Standing should be attempted only when the patient demonstrates sufficient stability and can safely tolerate the preceding positional change. Nurses should provide appropriate physical support and ensure that the patient does not stand independently when there is a recognized risk of falling. The patient should be observed for unsteadiness, pallor, sweating, dizziness, or sudden

weakness. Orthostatic blood pressure measurements should be obtained at the appropriate intervals, and mobilization should be interrupted if clinically significant symptoms or instability develop [3,16].

Ambulation Phase

Patients who tolerate standing may progress to short-distance walking with appropriate assistance. The initial walking distance should be individualized according to functional capacity, hemodynamic response, and the clinical indication for mobilization. Nurses should observe gait stability, fatigue, respiratory response, and the development of delayed orthostatic symptoms. The duration and intensity of subsequent activity should be adjusted according to the patient's response rather than based exclusively on predetermined mobility targets [6,7].

Post-Mobilization Phase

Following mobilization, the nurse should assist the patient into a safe resting position and reassess symptoms and relevant vital signs. Any episode of dizziness, presyncope, abnormal blood pressure reduction, or inability to complete the planned activity should be documented. The findings should be communicated to the healthcare team when further evaluation or modification of the treatment plan is required. Documentation should also indicate the level of assistance needed, the distance or duration of activity, and the patient's tolerance [3,11].

Table 5. Suggested nursing protocol for safe patient mobilization [3,10,11,16].

Stage	Nursing action	Monitoring parameters	Expected outcome
Before mobilization	Review history, medications, hydration, and previous symptoms	Baseline BP, HR, clinical stability	Identification of risk
Supine preparation	Allow adequate rest and obtain baseline measurements	BP and HR	Establish baseline
Supported sitting	Assist patient into sitting position	Dizziness, nausea, BP, HR	Tolerance of initial change
Bedside sitting	Allow gradual adaptation before standing	Symptoms and balance	Stable sitting
Supported standing	Assist with appropriate mobility support	BP, HR, pallor, weakness	Safe upright posture
Initial walking	Begin short-distance ambulation	Gait, fatigue, symptoms	Tolerated mobilization
Recovery	Assist patient to a safe resting position	BP, HR, symptom resolution	Hemodynamic recovery
Documentation	Record response and communicate abnormalities	Mobilization outcome	Continuity of care

Nursing Management of Patients Who Develop Orthostatic Symptoms During Mobilization

Immediate Recognition and Response

Immediate recognition of orthostatic symptoms is essential because continuing mobilization despite significant dizziness, weakness, or presyncope may increase the risk of falling and injury. Nurses should remain attentive to both patient-reported symptoms and observable changes, including pallor, sweating, unsteadiness, reduced responsiveness, and sudden inability to maintain an upright posture. The patient's activity should be stopped immediately when significant symptoms develop, and appropriate assistance should be provided to prevent injury [3,11].

Safe Repositioning and Hemodynamic Reassessment

The patient should be assisted into a safe recumbent position when clinically appropriate, with reassessment of blood pressure, heart rate, symptoms, and level of consciousness. Lower-extremity elevation may be considered when appropriate and not

contraindicated. Nurses should avoid allowing symptomatic patients to remain standing or walk without assistance. Repeated measurements can help establish whether blood pressure is recovering and whether the clinical response is consistent with an orthostatic episode [3,16].

Identification of Underlying Causes

Following stabilization, nurses should assess potential contributing factors, including dehydration, blood loss, medication exposure, infection, cardiac abnormalities, and prolonged immobility. Laboratory findings and relevant clinical observations should be reviewed when available. Persistent, recurrent, or severe episodes require medical assessment because they may reflect an underlying disorder that cannot be adequately managed through routine nursing interventions alone [4,10].

Escalation of Care

Urgent escalation is necessary when orthostatic symptoms are accompanied by syncope, chest pain, severe dyspnea, new neurological deficits, persistent hypotension, significant arrhythmia, or altered consciousness. These manifestations may indicate a serious cardiovascular or neurological condition rather than an uncomplicated postural blood pressure reduction. Nurses should follow institutional emergency response procedures and ensure that the patient remains protected from further injury while definitive assessment is arranged [3,16].

Factors Influencing the Effectiveness of Nurse-Led Interventions

Patient-Related Factors

The effectiveness of preventive interventions is strongly influenced by individual patient characteristics, including age, disease severity, autonomic function, hydration status, medication exposure, and physical ability. Patients with severe autonomic dysfunction may continue to experience orthostatic hypotension despite adequate hydration and gradual mobilization. In contrast, patients whose symptoms are primarily associated with temporary volume depletion or prolonged bed rest may respond more favorably to targeted interventions. Therefore, the selection of preventive measures should be based on clinical assessment rather than a uniform approach [4,9,18].

Nursing Knowledge and Clinical Competence

Nurses' knowledge of orthostatic hypotension, its risk factors, diagnostic criteria, and preventive interventions is an important determinant of care quality. Inadequate knowledge may result in missed symptoms, inaccurate blood pressure measurements, inappropriate mobilization progression, or failure to communicate clinically significant findings. Continuing education should therefore emphasize the recognition of orthostatic intolerance, standardized measurement techniques, appropriate use of mobility aids, and the implementation of individualized preventive strategies [11,16].

Staffing and Workload

Nursing workload and staffing levels may influence the consistency of preventive interventions. Safe mobilization frequently requires direct supervision, particularly among older adults, postoperative patients, and individuals with impaired balance. Limited staffing may reduce opportunities for repeated assessment or delay mobilization when assistance is unavailable. Healthcare organizations should therefore consider mobilization safety when planning staffing assignments and allocating resources to high-risk patients [8,16].

Availability of Equipment

Access to accurate blood pressure monitors, appropriately sized cuffs, mobility aids, compression devices, and safe patient-handling equipment supports effective preventive care. Inadequate or poorly maintained equipment may compromise assessment accuracy and increase the physical demands of mobilization. Standardized equipment availability and routine maintenance should therefore be incorporated into institutional patient safety programs [3,11].

Institutional Policies and Clinical Protocols

The presence of standardized institutional policies can improve consistency in orthostatic assessment, mobilization planning, documentation, and escalation of care. Protocols should define when orthostatic measurements are indicated, how measurements should be performed, which symptoms require interruption of activity, and when medical review is necessary.

However, protocols should not eliminate clinical judgment because individual patients may require modified approaches according to their condition and treatment goals [3,16].

Challenges and Barriers to Implementing Nurse-Led Preventive Interventions

Despite the availability of several non-pharmacological interventions, their consistent implementation remains challenging in clinical practice. One major barrier is the variability in assessment practices among healthcare professionals. Orthostatic vital signs may not be routinely measured before the first mobilization attempt, particularly when patients appear clinically stable while resting. This can lead to the under-recognition of postural intolerance and missed opportunities for prevention. Standardized nursing assessment procedures may help reduce this inconsistency [8,16].

Another challenge involves the complexity of patients' medical conditions. Patients with heart failure, kidney disease, or other fluid-sensitive disorders may not tolerate routine hydration strategies, while those with severe neurological impairment may be unable to perform physical counter-maneuvers. Similarly, compression garments may be uncomfortable or difficult to apply in patients with limited mobility or fragile skin. These limitations demonstrate that preventive interventions should be selected according to individual clinical needs and should not be implemented as universal requirements [10,13,14].

Patient-related barriers may also affect adherence. Some individuals may refuse mobilization because of fear of falling, previous experiences of dizziness, postoperative pain, fatigue, or lack of confidence. Others may attempt to stand independently despite instructions to request assistance. Nurses should address these barriers through reassurance, appropriate pain assessment, gradual exposure to movement, and repeated education regarding the importance of safe mobilization [5,11].

Organizational barriers include insufficient staffing, competing nursing responsibilities, lack of standardized protocols, inadequate staff education, and limited availability of equipment. In some settings, the absence of clear documentation requirements may make it difficult to identify recurrent episodes or evaluate whether preventive interventions are effective. Institutional support is therefore necessary to transform individual nursing actions into a coordinated and sustainable prevention strategy [11,16].

Evaluation of Nursing Outcomes and Indicators of Intervention Effectiveness

The evaluation of nurse-led interventions should involve objective and subjective indicators that reflect both physiological stability and functional performance. Blood pressure measurements provide objective evidence of postural changes, whereas symptom assessment captures the patient's experience of mobilization. Combining these measures offers a more comprehensive evaluation than relying on either measurement alone. Nurses should also document whether patients were able to complete planned activities and whether additional assistance was required [3,11].

One important indicator is the frequency of symptomatic orthostatic episodes during mobilization. A reduction in episodes following implementation of a structured nursing protocol may suggest improved preventive care, although differences in patient characteristics and clinical severity must be considered when interpreting the results. Other relevant indicators include the magnitude of systolic and diastolic blood pressure reductions, the frequency of dizziness and presyncope, and the proportion of patients who successfully complete planned sitting, standing, and walking activities [3,6].

Patient safety outcomes should also be included in the evaluation process. These may include falls during mobilization, near-fall events, injuries, episodes requiring urgent clinical intervention, and unplanned interruptions of rehabilitation. Although these outcomes are influenced by multiple factors, their monitoring can help healthcare organizations identify opportunities to improve the quality of mobilization practices. Functional outcomes, including walking distance, transfer independence, and participation in rehabilitation, provide additional information regarding the broader clinical value of preventive interventions [6,8].

Table 6. Suggested indicators for evaluating nurse-led orthostatic hypotension prevention [3,6,8,11].

Outcome domain	Suggested indicator	Measurement approach
Hemodynamic stability	Frequency of orthostatic BP reductions	Standardized orthostatic BP measurements
Symptom control	Episodes of dizziness and presyncope	Patient report and nursing observation

Mobilization tolerance	Successful completion of planned mobilization	Nursing documentation
Fall prevention	Falls and near-fall incidents	Institutional incident reports
Functional recovery	Ability to stand, transfer, and walk	Functional mobility assessment
Nursing practice	Compliance with assessment protocols	Nursing documentation audit
Patient education	Understanding of safe mobilization techniques	Teach-back and return demonstration
Clinical utilization	Mobilization interruptions and missed rehabilitation sessions	Clinical records
Patient experience	Confidence and perceived safety during mobilization	Patient feedback

Quality Improvement Strategies for Nursing Practice

Improving the prevention of orthostatic hypotension requires the integration of evidence-based nursing interventions into institutional quality improvement programs. Healthcare organizations should establish standardized protocols that identify high-risk patients, define appropriate orthostatic assessment procedures, and provide clear instructions for safe mobilization. These protocols should also specify the documentation requirements and clinical circumstances that necessitate interruption of activity or escalation of care. Regular review of institutional practices can help identify gaps between recommended interventions and actual nursing performance [11,16,20].

Continuing professional development is another important quality improvement strategy. Educational programs should address the physiological mechanisms of orthostatic hypotension, recognition of early symptoms, correct blood pressure measurement techniques, safe patient handling, and appropriate use of non-pharmacological interventions. Simulation-based education may be particularly valuable because it allows nurses to practice responding to symptomatic patients during mobilization without exposing actual patients to unnecessary risk. Periodic competency assessments can further support the consistent application of clinical protocols [11,15].

Healthcare organizations should also establish monitoring systems to evaluate adherence to mobilization protocols and track relevant clinical outcomes. Nursing audits may assess the completeness of orthostatic vital sign documentation, identification of high-risk patients, and appropriate responses to abnormal findings. Reviewing falls, near-falls, symptomatic episodes, and interrupted rehabilitation activities can provide additional information about the effectiveness of preventive practices. These findings can guide modifications to nursing protocols and support continuous improvement in patient safety [8,16].

Implications for Nursing Practice

The prevention of orthostatic hypotension has important implications for nursing practice because mobilization is a routine activity that requires continuous assessment of patient readiness and physiological tolerance. Nurses should recognize that apparently stable resting blood pressure does not necessarily indicate that a patient can safely tolerate standing or walking. Incorporating risk assessment and appropriate orthostatic monitoring into mobilization procedures can improve early recognition of hemodynamic instability and support safer clinical decision-making [3,10].

Another important implication is the need to move beyond isolated nursing interventions toward individualized, multicomponent care. Hydration support, gradual positional changes, compression therapy, medication surveillance, and physical counter-maneuvers address different mechanisms contributing to orthostatic hypotension. Their selection should be based on the patient's underlying condition, functional ability, and treatment restrictions. This approach reinforces the importance of clinical reasoning and patient-centered care rather than the indiscriminate application of standardized interventions [10,13,18].

Nursing practice should also emphasize the relationship between hemodynamic stability and functional recovery. Preventing orthostatic symptoms may allow patients to participate more consistently in rehabilitation and regain independence in activities of daily living. However, mobilization should always be balanced against immediate clinical safety. Nurses should avoid prioritizing activity targets over physiological stability and should ensure that every mobilization attempt is appropriate for the patient's current condition [5,6,7].

Recommendations for Future Research

Further research is required to establish the independent and combined effectiveness of nurse-led interventions in preventing orthostatic hypotension during mobilization. Although evidence supports several non-pharmacological strategies, many studies have focused on chronic orthostatic hypotension or neurological populations rather than hospitalized patients undergoing early mobilization. Consequently, additional clinical investigations are needed to determine which interventions provide the greatest benefit in intensive care, postoperative, medical-surgical, and rehabilitation settings [13,14].

Future randomized controlled trials should evaluate standardized nurse-led mobilization protocols against usual care, using clearly defined clinical and functional outcomes. Important outcomes should include the incidence of orthostatic hypotension, severity of postural blood pressure reductions, frequency of dizziness and presyncope, successful completion of mobilization, falls, and participation in rehabilitation. Studies should also consider differences in age, underlying diseases, medication exposure, hydration status, and duration of immobilization to identify patient populations most likely to benefit from particular interventions [3,5,7].

Research should additionally investigate the feasibility and sustainability of multicomponent nursing protocols in real-world healthcare environments. Factors such as nursing workload, staffing, equipment availability, staff education, and institutional support may influence intervention adherence and clinical effectiveness. Implementation studies could provide valuable information about how evidence-based preventive practices can be integrated into routine nursing workflows and maintained over time [11,16].

CONCLUSION

Orthostatic hypotension is an important clinical challenge during patient mobilization because it can interfere with cardiovascular stability, reduce tolerance of upright activity, increase the risk of falls, and delay functional recovery. Hospitalized patients are particularly vulnerable because of the combined effects of advanced age, prolonged immobilization, dehydration, medication exposure, neurological disorders, cardiovascular diseases, and postoperative physiological changes. Early recognition and prevention are therefore essential components of safe mobilization and high-quality nursing care.

Nurse-led interventions provide a practical and clinically important approach to reducing the risks associated with orthostatic hypotension. These interventions include comprehensive pre-mobilization assessment, standardized orthostatic vital sign monitoring, gradual positional transitions, individualized hydration support, prescribed compression therapy, physical counter-maneuvers, medication surveillance, patient education, and progressive mobilization. Each intervention addresses specific physiological or behavioral factors that may contribute to postural blood pressure reductions, and their appropriate combination can improve the safety and tolerability of mobilization.

The effectiveness of these interventions depends on accurate assessment, appropriate patient selection, consistent implementation, and continuous evaluation. Nurses must recognize that no single intervention is universally effective and that patients with complex medical conditions may require individualized multidisciplinary management. Furthermore, the identification of orthostatic hypotension should always lead to an appropriate clinical response, rather than being treated as an isolated abnormal measurement.

REFERENCES

[1] Lambert E, Lambert GW. Sympathetic dysfunction in vasovagal syncope and the postural orthostatic tachycardia syndrome. Frontiers in physiology. 2014 Jul 28;5:280.

[2] Freeman R, Wieling W, Axelrod FB, et al. Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome. Clinical Autonomic Research. 2011;21(2):69–72. doi:10.1007/s10286-011-0119-5.

[3] Wieling W, Kaufmann H, Claydon VE, et al. Diagnosis and treatment of orthostatic hypotension. The Lancet Neurology. 2022;21(8):735–746. doi:10.1016/S1474-4422(22)00169-7.

[4] Park JW, Okamoto LE, Shibao CA, Biaggioni I. Pharmacologic treatment of orthostatic hypotension. Autonomic Neuroscience. 2020 Dec 1;229:102721.

[5] Kurkis GM, Dennis DA, Johnson RM, Mejia M, Yazdani-Farsad Y, Jennings JM. Incidence and risk factors of orthostasis after primary hip and knee arthroplasty. The Journal of Arthroplasty. 2022 Jun 1;37(6):S70-5.

[6] Jans Ø, et al. Orthostatic intolerance during early mobilization after fast-track hip arthroplasty. British Journal of Anaesthesia. 2012;108(3):436–443.
[7] The impact of orthostatic intolerance on early ambulation following abdominal surgery in an enhanced recovery programme. Acta Anaesthesiologica Scandinavica. 2022.
[8] Orthostatic hypotension and falls in hospitalized older adults. Clinical Nursing Research. 2021;30(5):699–706. doi:10.1177/1054773820986682.
[9] Polin R, Ditmar MF. Pediatric Secrets-E-Book: Pediatric Secrets-E-Book. Elsevier Health Sciences; 2020 Jul 2.
[10] Orthostatic hypotension – a practical approach to investigation and management. Canadian Journal of Cardiology. 2017;33(12):1725–1731.
[11] Role of nurses and nurse practitioners in the recognition, diagnosis, and management of neurogenic orthostatic hypotension: A narrative review. International Journal of General Medicine. 2019.
[12] Orthostatic hypotension: management of a complex, but common, medical problem. Journal of the American Heart Association. 2022.
[13] Nonpharmacologic management of orthostatic hypotension: a systematic review. The American Journal of Medicine. 2015.
[14] Effectiveness of non-pharmacological interventions to treat orthostatic hypotension in elderly people and people with a neurological condition: A systematic review. Journal of Clinical Nursing. 2020.
[15] Gibbons CH, Schmidt P, Biaggioni I, et al. The recommendations of a consensus panel for the screening, diagnosis, and treatment of neurogenic orthostatic hypotension and associated supine hypertension. Journal of Neurology. 2017;264:1567–1582.
[16] Shen WK, Sheldon RS, Benditt DG, et al. 2017 ACC/AHA/HRS guideline for the evaluation and management of patients with syncope. Circulation. 2017;136:e60–e122.
[17] Shibao C, Lipsitz LA, Biaggioni I. Evaluation and treatment of orthostatic hypotension. Journal of the American Society of Hypertension. 2013;7(4):317–324.
[18] Ricci F, De Caterina R, Fedorowski A. Orthostatic hypotension: epidemiology, prognosis, and treatment. Journal of the American College of Cardiology. 2015;66(7):848–860.
[19] Robinson LJ, Pearce RM, Frith J. Acceptability of non-drug therapies in older people with orthostatic hypotension: a qualitative study. BMC geriatrics. 2018 Dec 17;18(1):315.
[20] Mancia G, Kreutz R, Brunström M, et al. 2023 ESH guidelines for the management of arterial hypertension. Journal of Hypertension. 2023;41(12):1874–2071.
[21] Freeman R. Clinical practice. Neurogenic orthostatic hypotension. New England Journal of Medicine. 2008;358:615–624.
[22] Low PA, Tomalia VA, Park KJ. Autonomic function tests: some clinical applications. Journal of Clinical Neurology. 2013;9(1):1–8.