

Original Research

View Article Online



Received 02 November 2021

Revised 04 January 2022

Accepted 05 January 2022

Available online 13 January 2022

Edited by Gokhan Zengin

KEYWORDS:

Muntingia calabura
biological activity
phytochemicals
bioactive compounds
leaf extract
root extract

Natr Resour Human Health 2022; (0): 1-10
<https://doi.org/10.53365/nrfhh/145564>
ISSN: xxxx-xxxx
Copyright © 2022 Visaga Publishing House

Gas Chromatographic analysis of potentially bioactive compounds in leaf and root extracts of *Muntingia calabura* and their expected antibacterial activities

Suvarchala Vankudoth¹, Pavani Chirumamilla¹, Spoorthi Veera¹, Shasthree Taduri^{1,*}

¹Department of Biotechnology, Kakatiya University, Warangal – 506009, Telangana, India

ABSTRACT: The study aimed to identify bioactive compounds in *Muntingia calabura* leaf and root methanolic extracts. The Gas Chromatography and Mass Spectroscopy (GC-MS) technique were used to identify bioactive compounds. GC-MS analysis revealed 38 compounds in the leaf and 15 compounds in the root methanolic extracts of *M. calabura*. The prime potent compound found in leaf extract is 2-{3-[(E)-2-(1H-indol-3-yl)ethenyl]-1,2,4-oxadiazol-5-yl}phenol with 5.78% peak area and cholest-4-en-6-on-3-ol is found in root extracts, has the highest 63.7% peak area and another potent compound Lupeol has 7.3% peak area. The bioactive compounds identified in *M. calabura* have antibacterial activity against various bacterial strains such as gram-positive and gram-negative bacteria, which showed the efficacy of *in vivo* plant extracts. These findings validate the therapeutic potentiality of *M. calabura* leaf and root samples. Furthermore, these screened potential bioactive compounds can be used effectively for biomedical and therapeutic applications.

1. INTRODUCTION

Medicinal plants are an essential source of several bioactive compounds, which have been used to treat several diseases caused in human beings. The World Health Organization (WHO) estimates that 80% of developing countries rely on therapeutic drugs (Vital & Rivera, 2009). Their low cost, easy accessibility and the search on medicinal plants have led to the discovery of novel therapeutic drugs against diseases.

Gas chromatography is an exciting tool used nowadays to determine new bioactive compounds from critical medicinal plants, which interest is discovering novel drugs. The identified compounds are used in cosmetics, drugs, pharmaceuticals, the food industry and forensic applications (Uma et al., 2009). Identifying bioactive compounds from different plant parts leads to an additional investigation of pharmacological and biological studies (Farid et al., 2015; Momin et al., 2014). Different potent active compounds in the various plant extracts can treat various human diseases (Konappa et al., 2020). Most of the world's population depends on plant-derived medicines, which has drawn the interest of researchers to invent new drugs. (Iwu et al., 1999). The infections caused by microbes threaten a major health hazard that is around 25% of all the deaths among the countries (Priyanka et al., 2015). Natural products have been mainly a rich source of anti-infective agents.

Many researchers have screened the *in vitro* crude extracts from plants with a history of use in folk medicine have been screened for antibacterial activity by many researchers (Cushnie & Lamb, 2005).

According to these reports, several pharmacological studies have been demonstrated using various plant parts of *M. calabura*. The *M. calabura* is a beautiful flowering shrub that belongs to the Muntingaceae family. The secondary metabolites of plants such as alkaloids, flavonoids, phenols, tannins and saponins were reported in the leaf extracts of *M. calabura*. (Buhian et al., 2016). These active compounds are essential in the food and research industry (Leema & Prakash, 2019). The leaf and root parts of *M. calabura* have high anti-inflammatory, antiulcer, antibacterial, antipyretic, and antioxidant activities (Krishnaveni et al., 2015; Sufian et al., 2013; Zakaria et al., 2007, 2011, 2014).

Thus, the present study evaluates the presence of bioactive compounds present in methanolic extracts of *M. calabura* leaf and root parts using GC-MS analysis and evaluations of their antibacterial activity by using the agar well diffusion method.

* Corresponding author.

E-mail address: tadurishasthree@kakatiya.ac.in (Shasthree Taduri)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2. MATERIALS AND METHOD

2.1. Plant material collection and sample preparation

Fresh *M. calabura* leaves and roots were collected from the greenhouse at (18°01'35.1"N 79°33'31.7"E). The collected plant materials were thoroughly washed under running tap water to remove impurities. These were dried for 12 days in the shade at room temperature (27 ± 2 °C) and ground into a fine powder. Five gram of plant materials were mixed with 50 mL of methanol and incubated in an orbital shaker for 48 hours at 130 rpm. Whatman No. 1 filter paper was used to filter the supernatant. The supernatants were evaporated at room temperature to obtain a crude extract, then subjected to GC-MS analysis and anti-microbial activity testing.

2.2. GC-MS instrumentation

M. calabura leaf and root methanol extracts were GC-MS analysed using a Perkin-Elmer GC system coupled with a silica capillary column 30m x 0.25mm ID x 0.25m df equipped with Elite-5MS (5 % diphenyl and 95 % dimethyl polysiloxane). Helium was used as a gaseous carrier in a 1:1 mg/minute ratio, with an injector volume of 2µl and a temperature of 250 °C and ionization at 70-eV. The oven temperature began at 110 °C and ended at 280 °C. The extraction time was 29 minutes for *M. calabura* leaf extracts and 13 minutes for root methanolic extracts. The total peak area of *M. calabura* was used to calculate the percentage of each bioactive compound in leaf and root extract. The Sophisticated Analytical Instrument Facility, IIT Bombay, performed the GC-MS analysis.

2.3. Microbial cultures and culture conditions

Muntingia calabura methanolic leaf and root extracts were tested against gram-negative and gram-positive bacteria. *Escherichia coli*, *Proteus vulgaris*, *Bacillus sphericus*, and *Pseudomonas fluorescens* were used as test organisms. Preparation of 24 hr old bacterial strains: inoculate nutrient broth with original cultures and incubate overnight at 37 °C. The antibacterial activity was tested using the agar well diffusion method with streptomycin (10 µg/mL) as the standard. The media was prepared and poured 15 mL into a petri dish for 5 minutes to solidify. The inoculums were swabbed uniformly over the media and dried. The standard (10 µg/mL) and three different leaf and root extract concentrations (45, 60, and 75 µg/mL) were incubated overnight at 37 °C in each well. After incubation, zones emerged, and the proportion of inhibition was scaled in millimetres.

3. RESULTS AND DISCUSSION

M. calabura phytochemical components were detected in methanolic leaf and root extracts. The percentage of the compounds was estimated using the retention time (RT) and peak area.

3.1. GC-MS analysis of leaf methanolic extract

The bioactive molecules found in methanolic leaf extracts were thirty-eight compounds. Among the thirty-eight compounds, the first compound identified with less retention time (4.85 min) was 1,3-dihydroxypropan-2-one and the peak area (0.24%), whereas the last compound with the longest retention time (33.68 min) was 3-Hydropregn-5-en-20-one hydrazone and the peak area (0.33%). Many of these compounds possess various pharmacological activities. The remaining phytochemical compounds are as follows: 1-(6-oxabicyclo[3.1.0]hexan-1-yl)ethan-1-one (or) 6-oxabicyclo(3.1.0)hexan-3-one, a phytoconstituent of methanolic leaf and root extracts of pili and safedshatavar (*Asparagus racemosus*), known to possess antioxidant activity (Banakar & Jayaraj, 2017). The 3,5-Dihydroxy-6-methyl-2,3-dihydro-4H-pyran-4-one reported antimicrobial, automatic nerve, and antioxidant properties in ethanolic stem extracts of an important medicinal plant *Waltheria indica* L (Ashwathanarayana & Naika, 2018). 2,3-dihydro-1-benzofuran, a bioactive compound, possesses anti-inflammatory activity in crude extracts of leaf and flower of *Pavetta crassicaulis* Bremek (Correa et al., 2017). Decan-3-yl acetate, reported to possess activities like asthma, anticancer, anti-inflammatory activities and Histamine H3 and H4 receptors, have been discovered drugs for Parkinson's and Alzheimer's disease (Tatipamula et al., 2019). The 1,2,3-benzenetriol (Pyrogallol), identified as a phytochemical compound that possesses anticancer activity in ethanolic extract of *Clathria baltica* (Sukprasert et al., 2020). The compound (1R,2S,3R,4S)-cyclopentane-1,2,3,4-tetrol, known to possess antiviral activity in *Lysiphyllum strychnifolium* plant extracts that treat against influenza virus (Rubaye et al., 2018). The compound, 4-[(1E)-3-hydroxyprop-1-en-1-yl]-2-methoxyphenol have multiple biological properties like anti-hyperglycemic, hepatoprotective, anti-obesity and antibiotic activities in root extracts of *Eclipta alba* (L) (Naik et al., 2019). More recently, it was reported that the compound (2E)-3,7,11,15-tetramethyl-2-hexadecen-1-ol has antifungal and antibacterial activity in leaf extracts *Rhizophora apiculata* (Lakshmanan et al., 2019).

Hexadecanoic acid, a phytochemical compound, has properties like strong antimicrobial, anti-inflammatory activities and acts against arthritis in medicinal plants like *Albizia adianthifolia*, *Preocarpus angolensis* and in some plants against food-borne pathogens (Abubakar & Majinda, 2016; Preethi et al., 2010). In *Cycas revolute*, a medicinal plant is known to have antioxidant properties in the 1-hentetracontanol bioactive molecule (Olabisi & Olubunmi, 2019). The (9Z,12E)-octadeca-9,12-dienoic acid and (9Z)-octadec-9-enoic acid have phytochemical properties like hypercholesterolemia, anti-acne, anti-cancer, anti-bacterial, and diabetes mellitus in methanolic leaf extract *Crateva adansonii* DC and root extract of *Jatropha pelargonifolia* Courb (Christiana et al., 2014). Another bioactive molecule, 3-[(4-ter-butylcyclohexyl)oxy]-3-phenyl-2-benzofuran-1(3H)-one is identified to synthesize green synthetic particles using ferrous and nickel nanoparticles by leaf extract of eucalyptus (Weng et al., 2017).

Table 1GC-MS of bioactive compounds present in the methanolic extracts of leaves derived from *in vivo* grown plants of *Muntingia calabura* L

S No	RT	Compound name	Formulae	Mwt	Area	Biological activity
1	4.85	1,3-dihydroxypropan-2-one	C ₃ H ₆ O ₃	90	0.24%	Ant oxidative (Leema & Prakash, 2019)
2	8.49	1-(6-oxabicyclo[3.1.0]hexan-1-yl)ethan-1-one (or) 6-oxabicyclo(3.1.0)hexan-3-one	C ₇ H ₁₀ O ₂	126	0.25%	Antimicrobial, automatic nerve activity and antioxidant (Banakar & Jayaraj, 2017)
3	9.68	3,5-Dihydroxy-6-methyl-2,3-dihydro-4H-pyran-4-one	C ₆ H ₈ O ₄	144	0.16%	Anti-inflammatory (Ashwathanarayana & Naika, 2018)
4	11.20	2,3-dihydro-1-benzofuran	C ₈ H ₈ O	120	0.3%	Asthma, Anticancer, Parkinson's and Alzheimer's disease (Correa et al., 2017)
5	11.56	decan-3-yl acetate	C ₁₄ H ₂₈ O ₂	228	0.07%	Anticancer (Tatipamula et al., 2019)
6	14.20	benzene-1,2,3-triol (Pyrogallol)	C ₆ H ₆ O ₃	126	0.17%	Antiviral (Sukprasert et al., 2020)
7	15.22	-D-Glucopyranosyl-(1->3)-β-D-fructofuranosyl β-D-glucopyranoside (or) D-(+)-Melezitose	C ₁₈ H ₃₂ O ₁₆	504	0.03%	Nutritional Anemia (Wei et al., 2020)
8	15.81	(1R,2S,3R,4S)-cyclopentane-1,2,3,4-tetrol (or))-cyclopentane-1,2,3,4-tetrol, 4b, 3b, 2a(1	C ₅ H ₁₀ O ₄	134	0.44%	Antibacterial and antifungal (Rubaye et al., 2018)
9	18.02	4-(ethoxymethyl)-2-methoxyphenol	C ₁₀ H ₁₄ O ₃	183	0.10%	No activity
10	19.30	4-[(1E)-3-hydroxyprop-1-en-1-yl]-2-methoxyphenol	C ₁₀ H ₁₂ O ₃	180	0.29%	Antihyperglycemic, hepatoprotective, Immunodulatory, Antiviral, Antifungal, Analgesic (Naik et al., 2019)
11	21.14	(2E)-3,7,11,15-tetramethyl-2-hexadecen-1-ol	C ₂₀ H ₄₀ O	296	0.13%	Antibacterial and antifungal (Lakshmanan et al., 2019)
12	22.35	hexadecanoic acid	C ₁₆ H ₃₂ O ₂	256	0.35%	Anti-inflammatory and Arthritis (Abubakar & Majinda, 2016)
13	24.35	1-hentetracontanol	C ₄₁ H ₈₄ O	592	0.07%	Antioxidant (Olabisi & Olubunmi, 2019)
14	24.75	(9Z,12E)-octadeca-9,12-dienoic acid	C ₁₈ H ₃₂ O ₂	280	0.74%	Hypocholesterolemic, Anti-acne (Christiana et al., 2019)
15	24.97	(9Z)-octadec-9-enoic acid Or Oleic acid	C ₁₈ H ₃₄ O ₂	282	0.29%	Diabetes mellitus (Aati et al., 2019)
16	26.77	N'-(4-tert-butylcyclohexylidene)-4-methylbenzene-1-sulfonohydrazide	C ₁₇ H ₂₆ N ₂ O ₂ S	322	0.08%	No activity

Continued on next page

Table 1 continued

17	27.22	4,8,12,16, tetramethylheptadecan-4-olide or 5-Methyl-5-(4,8,12-trimethyltridecyl)dihydro2(3H)-furanone	$C_{21}H_{40}O_2$	324	0.16%	No activity
18	27.45	2H-1-Benzopyran-7 Ol,3,4-dihydro-5-methoxy-6-methyl 2-phenyl	$C_{17}H_{18}O_3$	270	0.15%	No activity
19	27.61	3-[(4-ter-butylcyclohexyl)oxy]-3-phenyl-2-benzofuran-1(3H)-one	$C_{24}H_{28}O_3$	364	0.06%	Green synthetic particles (Weng et al., 2017)
20	27.92	5- hydroxy 2 - phenyl - 4 – chromanone	$C_{15}H_{12}O_3$	240	1.24%	Helicobacter pylori and Gastric antisecretory activity (Das et al., 2010)
21	28.46	(2Z)-2-[[5-(methoxycarbonyl)-3,4-dimethyl-1H-pyrrol-2-yl]methylidene]-3,4,5-trimethyl2H-pyrrol-1-iumbromid	$C_{17}H_{23}BrN_2O_2$	366	0.47%	No activity
22	28.64	1-tetradecene,2-decyl	$C_{24}H_{48}$	336	0.02%	Anti-inflammatory, Arthritis and antioxidant (Fernandes & Krishnan, 2019)
23	28.92	4H-1-benzopyran-4-one,2,3-dihydro-5,7-dihydroxy-2-phenyl(3)	$C_{15}H_{12}O_4$	256	2.11%	No activity
24	29.12	1,2-bis-[2-[4-cyclohepta-2,4,6-trienyl-phenoxy]-ethoxy]-ethane	$C_{32}H_{34}O_4$	482	1.20%	No activity
25	29.24	(2E)-1,3-diphenylprop-2-en-1-one	$C_{15}H_{12}O_3$	240	2.48%	Anti-proliferative (Landim et al., 2019)
26	29.56	2-{3 -[(E) - 2 -(1H -indol - 3 -yl)ethenyl]-1,2,4-oxadiazol-5-yl}phenol -	$C_{18}H_{13}N_3O_2$	303	5.78%	Antioxidant and Antitumor (Kopoytkoba & Caxho, 2019).
27	29.92	5 -hydroxy - 7 -methoxy - 2 -phenyl - 4H-1-benzopyran-4-one	$C_{16}H_{12}O_4$	268	0.63%	Anticancer, anti-prostate cancer and anti-inflammatory activities (Wang et al., 2019).
28	30.01	(2S)-5,7-dihydroxy-2-(4-hydroxyphenyl)-2,3-dihydro-4H-1-benzopyran-4-one	$C_{15}H_{12}O_5$	272	2.08%	Anti-tuberculosis, Anticancer (Sahu et al., 2020).
29	30.32	7-hydroxy-3-methoxy-2-(4-methoxyphenyl)-4 H-1-benzopyran-4-one	$C_{17}H_{14}O_5$	298	2.57%	Antioxidant (Suttiarporn et al., 2016)
30	30.44	4(1,1-Dimethylaaly)9-methoxy7H-furo (3,2-g)(1) benzopyrn-7-one	$C_{17}H_{16}O_4$	284	0.52%	No activity
31	30.83	2,2' -((1E,1'E) -(1,4 -phenylenebis(azanylylidene))bis(m ethanylylidene))diphenol	$C_{20}H_{16}N_2O_2$	316	1.10%	Anti-tumour and Cancer phototherapy (Kopoytkoba & Caxho, 2019)

Continued on next page

Table 1 continued

32	31.29	5,7-dihydroxy - 2 -(4 - hydroxyphenyl)-6-methoxy-4Hchromen-4-one	$C_{16}H_{12}O_6$	300	2.46%	Antioxidant (Karker et al., 2019; Mu et al., 2009; Yausheva et al., 2019)
33	31.67	2,3,7,8-tetramethoxy-5,11-dihydro10H-dibenzo[a,d][7]annulen-10-one	$C_{19}H_{20}O_5$	328	1.08%	Arthritis, Anti-inflammatory (Fernandes & Krishnan, 2019)
34	32.17	1-(4-methylphenyl)anthracene - 9,10-dione	$C_{21}H_{14}O_2$	298	2.15%	Anticancer (Chu et al., 2019)
35	32.78	8,9-dihydro-aflatoxin B1	$C_{17}H_{14}O_6$	314	0.66%	Antimycotoxin (Pok et al., 2020)
36	33.15	1-Docosene	$C_{22}H_{44}$	308	0.74%	Anticancer (Swantara et al., 2019)
37	33.45	3,6,8-trimethoxy-5,7-dimethyl-2-phenyl-4H-1-benzopyran-4-one	$C_{18}H_{16}O_7$	344	0.09%	No activity
38	33.68	3-Hydropregn-5-en-20-one hydrazone	$C_{21}H_{34}N_2O$	330	0.33%	No activity

5-hydroxy-2-phenyl-4-chromanone is one compound that possesses *Helicobacter pylori* and Gastric antisecretory activity (Das et al., 2010). 1-tetradecene,2-decyl reported to have phytochemical properties like an anti-inflammatory in leaf extracts of *Azima tetracantha* Lam (Jose & Panneerselvam, 2019) and antioxidant activity in leaf and stem of *Strobilanthes* species (Fernandes & Krishnan, 2019). Recently, the anti-inflammatory activity of (2E)-1,3-diphenylprop-2-en-1-one has been reported by Landim et al. (2019) in *Lonchocarpus cultratus*.

The compound, 5-hydroxy-7-methoxy-2-phenyl-4H-1-benzopyran-4-one, possesses anti-cancer, anti-prostate and anti-inflammatory activities (Marques et al., 2019; Wang et al., 2019). Similarly, a compound is known as (Naringenin) (2S)-5,7-dihydroxy-2-(4-hydroxyphenyl)-2,3-dihydro-4H-1-benzopyran-4-one is observed to have two biological properties like anti-tuberculosis and anti-cancer activities (Sahu et al., 2020). The major compound identified in the *M. calabura* leaf extract was found to be 2-{3-[(E)-2-(1H-indol-3-yl)ethenyl]-1,2,4-oxadiazol-5-yl}phenol 5.78% peak area, and have antioxidant and antitumor activity (Figure 1). These bioactive compounds are synthetic aromatic C-nucleoside derivatives developed by Sadek et al. (2014). 2,2'-((1E,1'E)-(1,4-phenylenebis(azanylylidene))bis(methanylylidene)diphenol (Nitrilo methyl dyne) has been reported to have an anti-tumor activity (Kopoytkoba & Caxho, 2019). The compound 7-hydroxy-3-methoxy-2-(4-methoxyphenyl)-4H-1-benzopyran-4-one that is identified to possess antioxidant activity from the bran of Thai black rice (Suttiarporn et al., 2016) and 5,7-dihydroxy-2-(4-hydroxyphenyl)-6-methoxy-4H-chromen-4-one reported to have strong antioxidant activity (Karker et al., 2019; Mu et al., 2009; Yausheva et al., 2019). 2,3,7,8-tetramethoxy-5,11-dihydro-10H-dibenzo[a,d][7]annulen-10-one has anti-inflammatory activity (Fernandes & Krishnan, 2019). The compound, Anthraquinone, 1-(4-methylphenyl)anthracene-9,10-dione is identified to act against the cancer cell and has antibacterial activities (Chu et al., 2019). 8,9-dihydro-aflatoxin B1 has the mycotoxin activity (Aparna et al., 2012; Pok et al., 2020), and 1-Docosene has the anticancer activity of *Xestospongia testudinaria*. Some compounds have been identified, but their biological activity has not yet been discovered (Table 1).

3.2. GC-MS analysis of root methanolic extract

Of all the bioactive compounds exhibited, cholest-4-en-6-on-3-ol has the highest 63.7% retention time (24.58 min) and has anti-obesity activity (Shaheed et al., 2019). At the same time, another compound, Lupeol, was observed during prolonged retention (35.9 min), and the maximum area percentage was 7.36% (Figure 2). Similarly, a compound is known, 17-octadecynoic acid has phytochemical properties like Hypocholesterolaemia, anti-inflammatory, anti-acne, and anti-coronary activities from various plant extracts (Benzidia et al., 2019; Mishra & Patnaik, 2020; Revathi & Dhanaraj, 2019). (2R)-2,5,8-trimethyl-2-[(4R,8R)-4,8,12-trimethyltridecyl]-3,4-dihydro-2H-1-benzopyran-6-ol (β -Tocopherol) is one of




Figure 1. GC-MS chromatogram of leaf methanolic extracts of *Muntingia calabura* and potent bioactive compound 2-{3-[(E)-2-(1H-indol-3-yl)ethenyl]-1,2,4-oxadiazol-5-yl}phenol (antioxidant and antitumor 5.78%).

the compounds that have anti-spasmodic activity (Lanuzza et al., 2017). The compounds like cholestan-3-one-4,4-dimethyl-(5 α); 9-octadecenoic acid(Z)-phenyl methyl ester; (9Z)-octadec-9-enoic acid (Oleic acid); androst-1-en-3-one,4,4-dimethyl-(5 α) identified to possess anti-cancer, anti-inflammatory, antioxidant, antiulcerogenic, antipyretic activities in various plant extracts of medicinally important plants (Safwat et al., 2018; Shelke & Bhot, 2019; Yamuna et al., 2017; Yue et al., 2020).




Figure 2. GC-MS chromatogram of root methanolic extracts of *Muntingia calabura*, potent anti-obesity compound cholest-4-en-6-on-3-ol(63.7%) and anti-cancerous compound Lupeol (7.3%).

Cholest-4-en-6-on-3-ol and Lupeol are the two identified compounds with highest percent of compounds among all the compounds in root methanolic extracts of *M. calabura* and reported to possess anticancer, antioxidant, anti-prostate, anti-inflammatory activities in different plant extracts of medicinal plants (Geetha & Varalakshmi, 2001; Shahaby et al., 2019; Shaheed et al., 2019; Yang et al., 2020). There are a few identified compounds, 1,12-Dibromododecane; 5-(2-bromopropan-2-yl)-2-methylcyclohexan-1-ol; 2H-1-benzopyran-6-ol,3,4-dihydro-2,8-dimethyl-(2-4,8,12-trimethyl tridecyl); A'-neogammacer-22(29)-en-3-one; 4,4,6a,6b,8a,11,11,14b-octamethyl-1,4,4a,5,6 yet to be

Table 2GC-MS of bioactive compounds present in the methanolic extracts of root-derived from *in vivo* grown plants of *Muntingia calabura* L

S No	RT	Compound name	Formulae	MWt	Area	Biological activity
1	22.36	hexadecanoic acid	C ₁₆ H ₃₂ O ₂	256	1.15%	Anti-inflammatory and arthritis (Abubakar & Majinda, 2016)
2	24.58	cholest-4-en-6-on-3-ol	C ₂₇ H ₄₆ O ₂	400	63.7%	Anti-obesity activity (Shaheed et al., 2019).
3	26.22	1,12-Dibromododecane	C ₁₂ H ₂₄ Br ₂	326	0.40%	No activity
4	26.66	17-octadecynoic acid	C ₁₈ H ₃₂ O ₂	280	0.66%	Hypocholesterolemic, Anti-inflammatory, Anti-acne, Anti-coronary (Revathi & Dhanaraj, 2019)
5	27.22	(2R)-2,5,8-Trimethyl-2-[(4R,8R)-4,8,12-trimethyltridecyl]-3,4-dihydro-2H-1-benzopyran-6-ol	C ₂₈ H ₄₈ O ₂	416	1.34%	Anti-spasmodic (Lanuzza et al., 2017)
6	28.2	5-(2-bromopropan-2-yl)-2-methylcyclohexan-1-ol	C ₁₆ H ₃₂ O ₂	256	0.94%	No activity
7	28.71	cholestan-3-one-4,4-dimethyl-(5 α)	C ₂₉ H ₅₀ O	414	0.25%	Anti-cancer and anti-inflammatory (Shaheed et al., 2019)
8	29.41	9-octadecenoic acid(Z)-phenyl methyl ester	C ₂₅ H ₄₀ O ₂	372	0.48%	Anti-ulcer genic, anti-androgenic, anti-inflammatory, anti-cancer (Shelke & Bhot, 2019).
9	29.87	(9Z)-octadec-9-enoic acid (Or)Oleic acid	C ₁₈ H ₃₄ O ₂	282	0.48%	Anti-pyretic, anti-nociceptive, anti-inflammatory (Aati et al., 2019)
10	30.19	(2R)-2,5,7,8-Tetramethyl-2-[(4R,8R)-4,8,12-trimethyltridecyl]-3,4-dihydrochromen-6-ol (or) Vitamin E	C ₂₉ H ₅₀ O ₂	430	8.26%	Anti-cancer and anti-oxidant (Shahaby et al., 2019)
11	33.73	2H-1-benzopyran-6-ol,3,4-dihydro-2,8-dimethyl(-2-4,8,12-trimethyl tridecyl)	C ₂₇ H ₄₆ O ₂	402	0.34%	No activity
12	34.7	androst 1-en-3-one,4,4-dimethyl-(5 α)	C ₂₁ H ₃₂ O	300	0.87%	Breast and cancer cell activity (Yue et al., 2020)
13	35.1	A'-neogammacer-22(29)-en-3-one	C ₃₀ H ₄₈ O	424	5.82%	No activity
14	35.4	4,4,6a,6b,8a,11,11,14b-Octamethyl-1,4,4a,5,6	C ₃₀ H ₄₈ O	424	6.78%	No activity
15	35.9	1,2,5,14,18,18-hexamethyl-8-(prop-1-en-2-yl)pentacyclo[11.8.0.0 ^{2,1'} .0 ^{5,9} .0 ^{1,19}]henicosan-17-ol (or) Lupeol	C ₃₀ H ₅₀ O	426	7.3%	Anti-prostate and anti-cancer (Yang et al., 2020).

identified (Figure 2 and Table 2).




Figure 3. Effect of *Muntingia calabura* leaf methanolic extract on various bacterial cultures.




Figure 4. Effect of *Muntingia calabura* root methanolic extract on various bacterial cultures.

3.3. Antibacterial activity for leaf and root methanolic extracts

The antibacterial activity of leaf methanolic extracts (45 µg/mL, 60 µg/mL, 75 µg/mL) and the control Streptomycin (10 µg/mL) was evaluated. The maximum

antibacterial activity was shown towards *E. coli* at 75 µg/mL concentration, followed by *B. sphericus*, *P. vulgaris* and *P. fluorescens*. As the concentration of the plant extract increased, the inhibition zone was also found to increase. (Figure 3 and 4). The leaf methanolic extract of *M. calabura* was considered the most effective extract with high anti-bacterial activity (Zakaria et al., 2010). In *M. calabura*, the ethanolic leaf extract has shown strong antibacterial activity against *E. coli* (Gurning et al., 2021). The root extracts of *Carica papaya* have shown the highest antibacterial activity against *Salmonella typhi* among tested gram-negative and gram-positive bacterial strains (Doughari et al., 2007).

4. CONCLUSION

This study revealed the presence of various bioactive molecules in the leaf and root methanolic extracts of *M. calabura*. Leaf methanolic extracts yielded thirty-eight compounds, while root extracts yielded a total of fifteen compounds based on their molecular weight, retention time, and peak area. The leaves of *M. calabura* contain the highest concentration of the antioxidant and antitumor compound is 2-{3-[(E)-2-(1H-indol-3-yl)ethenyl]-1,2,4-oxadiazol-5-yl}phenol, followed by roots of cholest-4-en-6-on-3-ol and Lupeol. These bioactive substances have been shown to have antibacterial effects against gram-positive and gram-negative bacteria. According to this study, the abundance of phytochemicals and bioactive substances in *M. calabura* makes it a potential source of medicines.

CONFLICTS OF INTEREST

Conflict of interest the authors declare that there are no conflicts of interest in this study.

ACKNOWLEDGMENTS

The first author gratefully acknowledges the National fellowship for Higher Education (NFST-2015-17-ST-TEL-2837) for providing financial assistance. The authors also thank Professor Nagavelli Vasudava Reddy, Department of Chemistry, Kakatiya University, Warangal for his guidance in correcting the Chemical names according to the IUPAC method.

ORCID

Suvarchala Vankudoth 0000-0001-5292-3999
 Pavani Chirumamilla 0000-0001-7645-025X
 Spoorthi Veera 0000-0002-0380-3250
 Shasthree Taduri 0000-0002-2372-6889

AUTHOR CONTRIBUTIONS

SV conducted the experimental work, writing the article, PC analyzed the data and designed the manuscript helped in experimental work, and SV designing Chromatogram figures and interpretation the data. ST extended overall guidance and finalized the manuscript.

REFERENCES

- Aati, H., El-Gamal, A., Kayser, O., 2019. Chemical composition and biological activity of the essential oil from the root of *Jatropha pelargonifolia* Courb native to Saudi Arabia. *Saudi Pharmaceutical Journal*. 27, 88–95. <https://doi.org/10.1016/j.jpsps.2018.09.001>
- Abubakar, M.N., Majinda, R.R., 2016. GC-MS analysis and preliminary antimicrobial activity of *Albizia adianthifolia* (Schumach) and *Pterocarpus angolensis* (DC). *Medicines*. 3(3), 1–9. <https://doi.org/10.3390/medicines3010003>
- Aparna, V., Dileep, K.V., Mandal, P.K., Karthe, P., Sadasivan, C., Haridas, M., 2012. Anti-inflammatory property of n-hexadecanoic acid Structural evidence and kinetic assessment. *Chemical Biology & Drug Design*. 80(3), 434–439. <https://doi.org/10.1111/j.1747-0285.2012.01418.x>
- Ashwathanarayana, R., Naika, R., 2018. Anti-Inflammatory properties of *Pavetta crassicaulis* Bremek. leaf and flower crude extracts and its pure compounds collected from Western Ghats. *Asian Journal of Pharmaceutical and Clinical Research*. 11(9), 72–90. <https://doi.org/10.22159/ajpcr.2018.v11i9.21885>
- Banakar, P., Jayaraj, M., 2017. Pharmacognosy, Phytochemistry and GC-MS Analysis of Ethanolic Stem Extract of *Waltheria indica* L-A Potent Medicinal Plant. *Journal of Biologically Active Products from Nature*. 7(5), 369–378. <https://doi.org/10.1080/22311866.2017.1400923>
- Benzidia, B., Barbouchi, M., Hammouch, H., Belahbib, N., Zouarhi, M., Erramli, H., Daoud, N.A., Badrane, N., Hajjaji, N., 2019. Chemical composition and antioxidant activity of tannins extract from green rind of *Aloe vera* (L) Burm.F. *Journal of King Saud University Science*. 31(4), 1175–1181. <https://doi.org/10.1016/j.jksus.2018.05.022>
- Buhian, W., Rubio, R., Valle, J.L., Puzon, J., 2016. Bioactive metabolite profiles and antimicrobial activity of ethanolic extracts from *Muntingia calabura* L. leaves and stems. *Asian Pacific Journal of Tropical Biomedicine*. 6(8), 682–685. <https://doi.org/10.1016/j.apjtb.2016.06.006>
- Christiana, O.A., Bull, O.E.J., Raphael, C.M., Joseph, O.O., Paul, M.O., Emmanuel, J. G., 2019. Gas Chromatographic Study of Bioactive Compounds in Methanolic Extract of Leaf of *Cratogeomys adansonii* DC. *Journal of Physics: conference series*. 1299(1), 1–9. <https://doi.org/10.1088/1742-6596/1299/1/012014>
- Chu, M.J., Wang, W., Ren, Z.L., Liu, H., Cheng, X., Mo, K., Wang, L., Tang, F., Lv, X.H., 2019. Discovery of novel triazole-containing pyrazole ester derivatives as potential anti-bacterial agents. *Molecules*. 1311(24), 1–11. <https://doi.org/10.3390/molecules24071311>
- Correa, M.F., Barbosa, A.J., Teixeira, L.B., Duarte, D.A., Simoes, S.C., Parreiras-E-Silva, L.T., Balbino, A.M., Landgraf, R.G., Bouvier, M., Costa-Neto, C.M., Fernandes, J.P., 2017. Pharmacological Characterization of 5-Substituted 1- [(2, 3-dihydro-1-benzofuran-2-yl) methyl] piperazines: Novel Antagonists for the Histamine H3 and H4 Receptors with Anti-inflammatory Potential. *Frontiers in Pharmacology*. 8, 1–12. <https://doi.org/10.3389/fphar.2017.00825>
- Cushnie, T.P.T., Lamb, A.J., 2005. Antimicrobial activity of flavonoids. *International journal of Antimicrobial Agents*. 26(5), 343–356. <https://doi.org/10.1016/j.ijantimicag.2005.09.002>
- Das, K., Tiwari, R.K.S., Shrivastava, D.K., 2010. Techniques for evaluation of medicinal plant products as antimicrobial agents: current methods and future trends. *Journal of Medicinal Plants*. 4(2), 104–111.
- Doughari, J.H., Elmahmood, A.M., Manzara, S., 2007. Studies on the antibacterial activity of root extracts of *Carica papaya* L. *African Journal of Microbiology Research*. 1(3), 37–041.
- Farid, M.M., Hussein, S.R., Ibrahim, L.F., Desouky, M.A., Elsayed, A.M., Oqlah, A.A., Saker, M.M., 2015. Cytotoxic activity and phyto-

- chemical analysis of *Arum palaestinum* Boiss. *Asian Pacific Journal of Tropical Biomedicine*. 5(11), 944–947. <https://doi.org/10.1016/j.apjtb.2015.07.019>
- Fernandes, M.C., Krishnan, S., 2019. GC-MS analysis of metabolites in leaf and stem of medicinally important *Strobilanthes* species from North Western Ghats of India. *Phytomorphology*. 69(3&4), 69–79.
- Geetha, T., Varalakshmi, P., 2001. Anti-inflammatory activity of lupeol and lupeollinoleate in rats. *Journal of Ethnopharmacology*. 76(1), 175–176. [https://doi.org/10.1016/S0378-8741\(01\)00175-1](https://doi.org/10.1016/S0378-8741(01)00175-1)
- Gurning, K., Simanjuntak, H.A., Purba, H., Situmorang, R.F.R., Barus, L., Silaban, S., 2021. Determination of total tannins and antibacterial activities ethanol extraction seri (*Muntingia calabura* L) leaves. *Journal of Physics: Conference Series*. 1811(1), 1–6. <https://doi.org/10.1088/1742-6596/1811/1/012121>
- Iwu, M.W., Duncan, A.R., Okunji, C.O., 1999. *New antimicrobials of plant origin Perspectives on new crops and new uses*, ASHS Press, Alexandria, VA, pp. 457–462.
- Jose, B.E., Panneerselvam, P., 2019. Identification of Phytochemical Constituents within the Leaf Extracts of *Azima tetracantha* Lam using Gas Chromatography-Mass Spectrometry. *Research Journal of Pharmacology and Pharmacodynamics*. 11, 17–22. <https://doi.org/10.5958/2321-5836.2019.00004.1>
- Karker, P.R., Kumar, R., S, Y., 2019. Exploring Flavonoids for the Inhibition of Hepatitis C Virus by Targeting NS3 Protease. *Journal of Pharmaceutical Sciences and Research*. 11(9), 3195–3200.
- Konappa, N., Udayashankar, A.C., Krishnamurthy, S., Pradeep, C.K., Chowdappa, S., Jogaiah, S., 2020. GC-MS analysis of phytoconstituents from *Amomum nilgiriicum* and molecular docking interactions of bioactive serverogenin acetate with target proteins. *Scientific Reports*. 10(1), 1–23. <https://doi.org/10.1038/s41598-020-73442-0>
- Kopoytkoba, Ø.B., Caxho, Ø.B., 2019. Nanoparticles with the aggregation-induced emission effect: properties and applications. *Biopolymers and Cell*. 35(4), 249–267. <https://doi.org/10.7124/bc.000A08>
- Krishnaveni, M., Raginabanu, C., Kalaivani, M., Krishnakumari, G., 2015. GC-MS/MS analysis of phytochemicals in *Muntingia calabura* L, Antimicrobial assay. *World journal of pharmaceutical research*. 4(4), 1855–1859.
- Lakshmanan, V., Thangaraj, M., Ponnusamy, B., Santhirakasan, S., Kannan, R., Regunathan, U., Selvaraj, S., 2019. Antibacterial Activity of *Rhizophora apiculata* Leaf Extract for the Management of Rice Bacterial Blight Disease. *Plant Pathology Journal*. 18, 39–46. <https://doi.org/10.3923/ppj.2019.39.46>
- Landim, E.M.D.S., Ruiz, A.L., Carvalho, J.E.D., Pomini, A.M., Pastorini, L.H., Santin, S.M.O., 2019. Antiproliferative activity and chemical constituents of *Lonchocarpus cultratus* (Fabaceae). *Natural product research*. 10, 1–4. <https://doi.org/10.1080/14786419.2019.1647427>
- Lanuzza, F., Occhiuto, F., Monforte, M.T., Tripodo, M.M., Angelo, D.V., Galati, E.M., 2017. Antioxidant phytochemicals of *Opuntia ficus indica* (L) Mill Cladodes with potential anti-spasmodic activity. *Pharmacognosy magazine*. 3(13), 424–429. https://doi.org/10.4103/pm.pm_495_16
- Leema, H.P., Prakash, O., 2019. Chemical composition and in vitro antioxidant potential of pili and safedshatavar (*Asparagus racemosus*). *The Pharma Innovation Journal*. 8, 153–158.
- Marques, J., Silva, A.M., Marques, M.P., Braga, S.S., 2019. Ruthenium (III) trithiacyclonane complexes of 7, 3', 4'-trihydroxyflavone, chrysin and tectochrysin: Synthesis, characterisation, and cytotoxic evaluation. *Inorganica Chimica Acta*. 488, 71–79. <https://doi.org/10.1016/j.ica.2019.01.003>
- Mishra, D., Patnaik, S., 2020. GC-MS Analyses phyto-chemicals and antibacterial activity of *Withania somnifera* L Dunal extract in the context of treatment to liver cirrhosis. *Biomed Pharmacol J*. 13, 71–78. <https://doi.org/10.13005/bpj/1862>
- Momin, M.A., Bellah, S.F., Rahman, S.M.R., Rahman, A.A., Murshid, G.M.M., Emran, T.B., 2014. Phytopharmacological evaluation of ethanol extract of *Sida cordifolia* L. roots. *Asian Pacific Journal of Tropical Biomedicine*. 4(1), 18–24. [https://doi.org/10.1016/s2221-1691\(14\)60202-1](https://doi.org/10.1016/s2221-1691(14)60202-1)
- Mu, Y.L., Xie, Y.Y., Zhou, L., Zhong, Y., Liu, L., Bai, H., Wang, Y.S., Zhang, X.M., 2009. Cardioprotective Effect of 'Methylamine Irisolidone. Hypoxia/Reoxygenation Injury in Cultured Rat Cardiac Myocytes. 26, 1170–1177. <https://doi.org/10.1002/cbdv.200800314>
- Naik, S., Gurushanthaiah, M., Kavimani, M., Mahesh, G.M., 2019. Extraction of Phytochemical constituents from the root of *Eclipta Alba* through gas chromatography-mass spectrometry procedure. *International Journal of Research in Pharmaceutical Sciences*. 10(2), 1238–1242. <https://doi.org/10.26452/ijrps.v10i2.414>
- Olabisi, A.E., Olubunmi, O.O., 2019. Essential oil compositions and antioxidant properties of *Cycas revoluta* (Thunb). *GSC Biological and Pharmaceutical Sciences*. 9, 102–109. <https://doi.org/10.30574/gscbps.2019.9.2.0203>
- Pok, P.S., Londoño, V.A.G., Vicente, S., Romero, S.M., Pacin, A., Tolaba, M., Alzamora, S.M., Resnik, S.L., 2020. Evaluation of citrus flavonoids parasiticus in maize: Aflatoxins reduction and ultrastructure alterations. *Food chemistry*, 126414. <https://doi.org/10.1016/j.foodchem.2020.126414>
- Preethi, R., Devanathan, V.V., Loganathan, M., 2010. Antimicrobial and antioxidant against *Aspergillus* efficacy of some medicinal plants against food borne pathogens. *Advances in Biological Research*. 4(2), 122–125.
- Priyanka, C., Kumar, P., Bankar, S.P., Karthik, L., 2015. In vitro antibacterial activity and gas chromatography-mass spectroscopy analysis of *Acacia karoo* and *Ziziphus mauritiana* extracts. *Journal of Taibah University for Science*. 9(1), 13–19. <https://doi.org/10.1016/j.jtusc.2014.06.007>
- Revathi, N., Dhanaraj, T.S., 2019. Evaluation of bioactive phytochemicals in leaves extract of *Dodonaea angustifolia* using gas chromatography and mass spectroscopic. *J Pharmacognosy and Phytochemistry*. 8(3), 4406–4409.
- Rubaye, A.F., Hameed, I.H., Kamal, S.A., 2018. Screening of metabolites products of *Fusarium oxysporum* and determination of its antibacterial and antifungal activity using medicinal plants extract. *Indian Journal Public Health Research Development*. 9(3), 399–404. <https://doi.org/10.5958/0976-5506.2018.00243.7>
- Sadek, M.M.E., Nagwa, S., El-Dayem, Seham, Y., Hassan, M.A., Momin, M.A., Bellah, S.F., Rahman, S.M.R., Rahman, A.A., Murshid, G.M.M., Emran, T.B., 2014. Phytopharmacological evaluation of ethanol extract of *Sida cordifolia* L roots. *Asian Pacific Journal of tropical biomedicine*. 4(1), 60202–60203. [https://doi.org/10.1016/S2221-1691\(14\)60202-1](https://doi.org/10.1016/S2221-1691(14)60202-1)
- Safwat, G.M., Hamed, M.M., Moatamed, S.A., 2018. Studies of the biological activity of *Cassia fistula*. *Pharmacologyonline*. 1, 75–85.
- Sahu, N., Mishra, G., Chandra, H.K., Nirala, S.K., Bhadauria, M., 2020. Naringenin mitigates anti tuberculosis drugs induced hepatic and renal injury in rats. *Journal of Traditional and Complementary Medicine*. 10(1), 26–35. <https://doi.org/10.1016/j.jtcm.2019.01.001>
- Shahaby, O.A., Zayat, M., Fattah, A., Hefny, M.M., 2019. Evaluation of the biological activity of *Capparis spinosa* var. *aegyptiaca* essential oils and fatty constituents as Anticipated Antioxidant and Antimicrobial

- Agents. Progress in Chemical and Biochemical Research. 2(4). <https://doi.org/10.33945/SAMI/PCBR.2019.4.6>
- Shaheed, K.A., Algaraawi, N.I., Alsultany, A.K., Abbas, Z.H., Khshayish, I.K., Alkhazali, M.T., 2019. Analysis of bioactive phytochemical compound of *Cyperus iria* L. By using gas chromatography-mass spectrometry. *Earth and Environmental Science*. 388(1), 1–11. <https://doi.org/10.1088/1755-1315/388/1/012064>
- Shelke, V., Bhot, M., 2019. GC-MS Analysis of Bio-active Compounds in Ethanolic extract of leaf and stem of *Asclepias curassavica* L. *International Journal of Pharmaceutical Investigation*. 9(2), 67–70. <https://doi.org/10.5530/ijpi.2019.2.13>
- Sufian, A.S., Ramasamy, K., Ahmat, N., Zakaria, Z.A., Yusof, M.I., 2013. Isolation and identification of antibacterial and cytotoxic compounds from the leaves of *Muntingia calabura* L. *Journal of Ethnopharmacology*. 146(1), 198–204. <https://doi.org/10.1016/j.jep.2012.12.032>
- Sukprasert, S., Pansuksan, K., Sriyakul, K., 2020. Lysiphyllum strychnifolium (Craib) A. Schmitz extract, a novel neuraminidase inhibitor of avian influenza virus subtype H5N1. *Journal of Herbal Medicine*(20), 100330–100330. <https://doi.org/10.1016/j.hermed.2020.100330>
- Suttiarporn, P., Sookwong, P., Mahatheerant, S., 2016. Fractionation and identification of antioxidant compounds from bran of Thai black rice cv. Riceberry. *International Journal of Chemical Engineering and Applications*. 7(2), 109–114. <https://doi.org/10.7763/IJCEA.2016.V7.552>
- Swantara, M.D., Rita, W.S., Suartha, N., Agustina, K.K., 2019. Anticancer activities of toxic isolate of *Xestospongia testudinaria* sponge. *Veterinary world*. 12(9), 1434–1434. <https://doi.org/10.14202/vetworld.2019.1434-1440>
- Tatipamula, Vb, Killari, K.N., Prasad, K., Rao, G.S., Talluri, M.R., Vantaku, S., Bilakanti, D., Srilakshmi, N., 2019. Cytotoxicity studies of the chemical constituents from marine algae *Charabaltica*. *Indian Journal of Pharmaceutical Sciences*. 81(5). <https://doi.org/10.36468/pharmaceutical-sciences.575>
- Uma, B., Prabhakar, K., Rajendran, S., Lakshmi, S.Y., 2009. Studies on GC/MS Spectroscopic Analysis of some Bioactive Antimicrobial Compounds from *Cinnamomum zeylanicum*. *Journal of Medicinal Plants*. 8, 125–131.
- Vital, P., Rivera, W.L., 2009. Antimicrobial activity and cytotoxicity of *Chromolaena odorata* (L.f) King and Robinson and *Uncaria perrottetii* (A. Rich) Merr. extracts. *Journal of Medicinal Plants Research*. 3(7), 511–518.
- Wang, Y., Ke, R.J., Jiang, P.R., Ying, J.H., Lou, E.Z., Chen, J.Y., 2019. The effects of tectochrysin on prostate cancer cells apoptosis and its mechanism. *Chinese journal of applied physiology*. 35(3), 283–291.
- Wei, Z., Sun, Z., Wang, Y., Dong, S., Lv, C., Wang, T., Yan, S., Wang, C., 2020. Fourier transform infrared and gas chromatography-mass extracts from *Euscaphis japonica* bark. *Thermal Science*. 24(3A), 1673–1680. <https://doi.org/10.2298/TSCI190611038W>
- Weng, X., Guo, M., Luo, F., Chen, Z., 2017. One-step green synthesis of bimetallic Fe/Ni nanoparticles by eucalyptus leaf extract: Biomolecules identification, characterization and catalytic activity. *Chemical Engineering Journal*. 308, 904–911. <https://doi.org/10.1016/j.cej.2016.09.134>
- Yamuna, P., Abirami, P., Vijayashalini, P., Sharmila, M., 2017. GC-MS analysis of bioactive compounds in the entire plant parts of ethanolic extract of *Gomphrena decumbens* Jacq. *Journal of Medicinal Plants Studies*. 5(3), 31–37.
- Yang, H.Q., Chen, G.H., Dong, S., Sun, Z., Wang, Y., Luo, X., Chen, B., Yao, G., Gao, Y., Lv, C., Zheng, D., 2020. Chemical constituents and medical function of leaves of *Diospyros lotus* L. *Thermal Science*. 24(3), 1633–1639. <https://doi.org/10.2298/TSCI190525033Y>
- Yausheva, E.V., Duskaev, G.K., Levakhin, G.I., Nurzhanov, B.S., Yuldashbaev, Y.A., Rysaev, A.F., Rakhmatullin, S.G., Inchagova, K.S., 2019. Evaluation of the effects of plant extracts on cattle rumen micro-biome. *InIOP Conference Series: Earth and Environmental Science*. <https://doi.org/10.1088/1755-1315/341/1/012165>
- Yue, X., Yang, H., Wang, T., Dong, S., Sun, Z., Wang, Y., Luo, X., Chen, B., Yao, G., Gao, Y., Lv, C., 2020. Molecules and medical function of *Diospyros lotus* L. *Thermal Science*. 24(3), 1705–1712. <https://doi.org/10.2298/TSCI190529042Y>
- Zakaria, Z.A., Hazalin, N.M., Zaid, S.M., Ghani, M.A., Hassan, M.H., Gopalan, H.K., Sulaiman, M., 2007. Antinociceptive, anti-inflammatory and antipyretic effects of *Muntingia calabura* aqueous extract in animal models. *Journal of Natural Medicines*. 61(4), 443–448. <https://doi.org/10.1007/s11418-007-0167-2>
- Zakaria, Z.A., Mohamed, A.M., Jamil, N.M., Rofee, M.S., Husain, M.K., Sulaiman, M.R., The, L.K., Salleh, M.Z., 2011. In vitro antiproliferative and antioxidant activities of the extracts of *Muntingia calabura* leaves. *The American Journal of Chinese Medicine*. 39(01), 183–200. <https://doi.org/10.1142/S0192415X11008749>
- Zakaria, Z.A., Sani, M.H., Cheema, M.S., Kader, A.A., Kek, T.L., Salleh, M.Z., 2014. Antinociceptive activity of methanolic extract of *Muntingia calabura* leaves: further elucidation of the possible mechanisms. *BMC Complementary and Alternative Medicine*(1), 1–12. <https://doi.org/10.1186/1472-6882-14-63>
- Zakaria, Z.A., Sufian, A.S., Ramasamy, K., Ahmat, N., Sulaiman, M.R., Arifah, A.K., Zuraini, A., Somchit, M.N., 2010. In vitro antimicrobial activity of *Muntingia calabura* extracts and fractions. *African Journal of Microbiology Research*. 4(4), 304–308.