

Original Research

View Article Online



Received 18 October 2021  
Revised 20 January 2022  
Accepted 21 January 2022  
Available online 07 February 2022

Edited by Barbara Sawicka

## KEYWORDS:

*Cleome rutidosperma*  
biological activity  
photochemical  
FTIR

Natr Resour Human Health 2022; (0): 1-6  
<https://doi.org/10.53365/nrhh/146009>  
eISSN: 2583-1194  
Copyright © 2022 Visagaa Publishing House

## Comparative antioxidant, antibacterial and phytochemical analysis of roots, stems, leaves and seeds from *Cleome rutidosperma* DC

Radjasagarin Arumugam<sup>1,\*</sup>, Balachandravinayagam Elanchezhian<sup>2</sup>, Jayakumar Samidurai<sup>3</sup>, Kaliyamoorthy Amirthaganesan<sup>2</sup>

<sup>1</sup>Department of Botany, A.V.C. College (Autonomous), Mannampandal, Tamil Nadu, India

<sup>2</sup>Department of Chemistry, A.V.C. College (Autonomous), Mannampandal, Tamil Nadu, India

<sup>3</sup>P.G. and Research Department of Zoology and Wildlife Biology, A.V.C. College (Autonomous), Mannampandal, Tamil Nadu, India

**ABSTRACT:** The emerging microbial infections and their resistance to the existing antibiotics lead to discovering novel compounds, primarily from medicinal plants with secondary metabolites having several bioactive potentials, including antioxidants. The current investigation aims to measure the antioxidant and antibacterial efficacy of ethanolic extracts from roots, stems, leaves and seeds of *Cleome rutidosperma*. The extracts were subjected to quantitative (total phenolic and flavonoid), qualitative phytochemical studies, and functional groups identification by FT-IR analysis. The extract of leaves showed the highest total antioxidant ( $54.21 \pm 1.56$  mg ABAE/g), DPPH ( $62.92 \pm 1.94$  mg GAEs/g), and FRAP ( $71.64 \pm 2.02$  mg GAEs/g) activity among the all-tested parts. The antibacterial efficacy of extracts was determined by the microdilution bioassay method, which demonstrated that G(+ve) bacteria appear to be more susceptible to the crude extracts than G(-ve) bacteria. The qualitative phytochemical screening-detected alkaloids, flavonoids, phenols, sugars, proteins, saponins, sterols, tannins, and terpenoids. The leaves have the highest levels of phenolics ( $70.451.23$  mg GAE/g DW) and flavonoids ( $32.261.12$  mg RE/g DW) among the all-tested parts. The extracts' functional group was validated using the FT-IR spectra. Polyphenols, flavonoids, and tannins were identified in the crude extracts. These findings imply that *C. rutidosperma* could be a promising candidate for further research into infectious illness treatment and as a resource of novel antioxidants in nutraceutical and biopharmaceutical industries as a functional additive.

## 1. INTRODUCTION

For the past few decades, researchers have been examining the biological properties of medicinal and food plants in the hopes of discovering a cure for a variety of modern diseases and a way to delay the onset of ageing symptoms. Free radicals and other reactive oxygens are highly reactive atoms or groups of atoms that contain an unpaired electron. The term "reactive oxygen species" (ROS) refers to oxygen derivatives that have the potential to be reactive (e.g.,  $O_2^{\cdot-}$ ,  $H_2O_2$ ,  $\cdot OH$ , and  $\cdot NO$ ). ROS are involved in oxidative damage to essential cellular components like proteins, lipids, lipoproteins, and DNA, which frequently occurs in biological systems and is associated with the etiology of diabetes, atherosclerosis, arthritis, cancer, cardiac problems, and neurological diseases, as well as the aging process (Farber, 1994; Halliwell & Gutteridge, 1985; Venkatesh & Dorai, 2011). Antioxidants can assist in preventing the

oxidation of essential cellular components and guard against free radical damage. The scientific community is looking for new antioxidant agents with less deleterious effects than existing synthetic antioxidants (butylated hydroxyanisole and butylated hydroxytoluene) (Bajpai et al., 2014; Jun et al., 2014). Another primary public health concern is antibiotic resistance, particularly in developing nations where transmittable ailments are the primary reason for death (N.D. Silva et al., 2016).

Numerous plants have been utilized as alternative medicines for many infections, ailments, and food preservatives since ancient times, implying the existence of antimicrobial and antioxidant elements (Panovska et al., 2005). Plant-derived antioxidants guard against oxidative stress by discarding ROS and RNS (reactive nitrogen species). They also suppress the polymerization chain reaction that free radicals initiate (Ahmed et al., 2015; Bianco & Uccella, 2000; Céspedes et al., 2008; Lafka et al., 2007; O.A. Silva et al., 2012). Nearly 200 species

\* Corresponding author.

E-mail address: [rmugam@gmail.com](mailto:rmugam@gmail.com) (Radjasagarin Arumugam)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

are found in the genus *Cleome*, the Capparidaceae family. Since ancient times *Cleome* species has been used in folk medicine for its ethnomedicinal properties, including antimicrobial, antidiabetic, anthelmintic, anti-inflammatory, anticonvulsant, antipyretic, antidiarrheal, and carminative properties, as well as in the therapy of wounds, epilepsy, convulsions, spasm, pain, and skin ailments (Archi et al., 2016; U. Bose et al., 2011; Devi et al., 2002; Motaal et al., 2011; Parimaladevi et al., 2003; Thomas et al., 2014). So far, there is no literature on the comparative antioxidant and antimicrobial potential of different parts of *C. rutidosperma*. Therefore this study aimed to determine the antioxidant and antibacterial potential of ethanolic extracts from roots, stems, leaves and seeds of *C. rutidosperma* and to reveal the functional groups present in the test plant using FTIR, which possibly open a new perspective for rising novel products such as medicine.

## 2. MATERIALS AND METHODS

### 2.1. Chemicals and reagents

Sigma-Aldrich provided the DPPH. All other compounds were of analytical quality and were obtained from Qualigens – Thermo Fisher Scientific, Mumbai, India.

### 2.2. Plant material

*Cleome rutidosperma* DC was collected in January 2020 on the campus of A.V.C. College in Mannampandal, Mayiladuthurai, Tamil Nadu, India. The root, stem, leaves, and seeds of the plant were removed. After thorough rinsing, the samples were dried in the shade until they attained a consistent weight. Using a mixer grinder, the dry samples were crushed to a fine powder and stored for later use. The extract preparation and analytical methods followed in this study were given in the supplementary file.

### 2.3. Statistical analysis

All tests were performed in triplicate, and the findings are expressed as mean  $\pm$  SD. The data were evaluated statistically using one-way ANOVA with Tukey's ( $p < 0.01$ ) test using the SPSS v22.

## 3. RESULTS AND DISCUSSION

### 3.1. Total antioxidant activity

Table 1 shows the findings of various assays (total antioxidant, DPPH, and FRAP) used to investigate the antioxidant potential of ethanolic extracts from *C. rutidosperma*. The total antioxidant activity of different parts of *C. rutidosperma* extracts ranged from  $30.63 \pm 1.34$  to  $54.21 \pm 1.56$  mg ABAE/g, with higher antioxidant activity in leaves and lower antioxidant activity in roots. In terms of total antioxidant activity, there was a significant ( $p < 0.01$ ) difference across the different parts of *C. rutidosperma*. Molybdenum VI is reduced in the phosphomolybdenum assay to generate a green phosphate/Mo<sup>5+</sup> combination.

### 3.2. DPPH radical scavenging activity

The radical scavenging potential of the plant components evaluated here showed significant ( $p < 0.01$ ) differences. DPPH radical scavenging activity ranged from  $36.62 \pm 1.62$  to  $62.92 \pm 1.94$  mg GAEs/g. The leaves had a significantly higher radical scavenging potential than the roots, significantly lower. The capacity of antioxidants to donate hydrogen is thought to be the mechanism by which they scavenge DPPH radicals. DPPH utilises an electron or hydrogen radical to create a stable diamagnetic molecule. When DPPH is combined with a hydrogen atom donor, a stable non-radical form of DPPH is generated, and a colour shifts from violet to light yellow. (Molyneux, 2004). As a result, DPPH is commonly used as a free radical in evaluating reducing chemicals (Cotelle et al., 1996) and is a useful reagent for evaluating compounds' free radical scavenging characteristics (Duan et al., 2006). According to Ghimire et al. (2011), the existence of bioactive chemicals such as phenolics, flavonoids, and tannins in different plant sections could reflect the varying scavenging activities. The perusal of available literature indicates a clear link between the radical scavenging ability of plant extracts and phytochemical contents (Arumugam et al., 2019, 2020; Namvar et al., 2017). The current findings also support the view of Yingming et al. (2004), who found that extracts with a high phenolic content were more effective in scavenging free radicals. Nonetheless, other extracts with lower phenolic levels demonstrated considerable activity, suggesting that other secondary metabolites may contribute to the scavenging activity.

### 3.3. Ferrous reducing ability (FRAP)

The ferrous reducing ability of different parts of *C. rutidosperma*, measured in Gallic acid equivalent (mg/g), was shown to be significantly different ( $p < 0.01$ ) and increased in the order of roots, stems, seeds, and leaves in the current investigation. Like other antioxidant activities, the leaves ( $71.64 \pm 2.02$  mg GAEs/g) demonstrated a stronger ability to reduce Fe<sup>3+</sup> than roots ( $42.51 \pm 1.08$  mg GAEs/g). In the FRAP assay, antioxidants in the samples used a redox-linked colorimetric mechanism involving single electron transfer to convert ferric (III) to ferrous (II) (Li et al., 2006). Flavonoids have a perfect structure for scavenging free radicals because they include several hydroxyls that act as hydrogen donors, making them potent antioxidant agents (Marya et al., 2011). According to Barros et al. (2007), the existence of reductones, which act as antioxidants by donating a hydrogen atom to the free radical chain, is associated with the presence of reducing capabilities.

### 3.4. Antibacterial activity

Table 2 summarises the antibacterial efficacy of ethanol extracts from *C. rutidosperma* roots, stems, leaves, and seeds against six human pathogens. The MIC was estimated over the test pathogens. The lowest MIC values among the four extracts indicated the highest antibacterial activity. All *C. rutidosperma* extracts exhibited a broad antibacterial activity, as they inhibited all microorganisms tested. The range of MIC

**Table 1**  
Antioxidant activity of ethanolic extracts from *C. rutidosperma*.

| Assays  | Roots                   | Stems                   | Leaves                   | Seeds                    |
|---|-------------------------|-------------------------|--------------------------|--------------------------|
| Total antioxidant activity (mg ABAE/g) <sup>x</sup> | 30.63±1.34 <sup>d</sup> | 43.22±1.85 <sup>c</sup> | 54.21±1.56 <sup>a</sup>  | 47.22± 1.24 <sup>b</sup> |
| DPPH radical (mg GAEs/g) <sup>y</sup>               | 36.62±1.62 <sup>c</sup> | 49.12±1.33 <sup>b</sup> | 62.92±1.94 <sup>a</sup>  | 51.51±1.55 <sup>b</sup>  |
| FRAP reducing power (mg GAEs/g) <sup>y</sup>        | 42.51±1.08 <sup>c</sup> | 54.85±2.45 <sup>b</sup> | 71.64±2.02 <sup>a</sup>  | 58.12± 1.43 <sup>b</sup> |
| Total phenolic (mg GAEs/g) <sup>y</sup>             | 37.23±0.62 <sup>c</sup> | 49.31±1.44 <sup>b</sup> | 70.45 ±1.23 <sup>a</sup> | 51.22±1.31 <sup>b</sup>  |
| Total flavonoid (mg REs/g) <sup>z</sup>             | 20.11±0.89 <sup>c</sup> | 25.33±1.62 <sup>b</sup> | 32.26±1.12 <sup>a</sup>  | 27.41±1.52 <sup>b</sup>  |

Different subscripts in the same column indicate significant difference (p < 0.01).

to inhibit *Salmonella typhii* (1.56 - 6.25 µg/mL), *Escherichia coli* (1.56 - 6.25 µg/mL), *Pseudomonas aeruginosa* (3.125 - 6.25 µg/mL), *Staphylococcus aureus* (6.25 - 3.125 µg/mL), *Streptococcus pyogenes* (1.56 - 3.125 µg/mL) and *Bacillus subtilis* (0.78 - 3.125 µg/mL). Significant inhibition was defined as MIC less than or equal to 1.56 mg/mL. The extracts were more sensitive to G(+) bacteria than G(-) bacteria. *B. subtilis* and *S. pyogenes* were most susceptible to the extracts, and the rest of the pathogens showed almost similar kinds of MIC values. Similarly, N.D. Silva et al. (2016) found that *Cleome spinosa* extracts were more active against G(+) bacteria. McNeil et al. (2018) reported that essential oil from above-ground portions of *C. rutidosperma* displayed antibacterial and antifungal action, with the most substantial inhibitory effect against *Bacillus cereus*. The ethanolic extract of *C. rutidosperma* leaves has higher antimicrobial activity than the aqueous extract (Ghosh et al., 2019).

**Table 2**

Minimum inhibitory concentrations (MIC - µg/mL) of ethanolic extracts from *C. rutidosperma* against selected bacterial pathogens.

| Pathogens                     | Roots | Stem | Leaves | Seeds | Gentamycin |
|-------------------------------|-------|------|--------|-------|------------|
| <i>Salmonella typhii</i>      | 6.25  | 12.5 | 3.12   | 6.25  | 0.195      |
| <i>Escherichia coli</i>       | 12.5  | 12.5 | 6.25   | 6.25  | 0.195      |
| <i>Pseudomonas aeruginosa</i> | 6.25  | 6.25 | 3.12   | 3.12  | 0.195      |
| <i>Bacillus subtilis</i>      | 1.56  | 1.56 | 0.39   | 0.78  | 0.049      |
| <i>Staphylococcus aureus</i>  | 1.56  | 1.56 | 0.39   | 0.78  | 0.024      |
| <i>Streptococcus pyogenes</i> | 3.12  | 3.12 | 1.56   | 3.12  | 0.097      |

In the present study, *Bacillus* sp. was most susceptible to the extracts. Against all the tested bacterial pathogens, leaves extract of *C. rutidosperma* had the highest inhibition effects. When compared to the antibacterial activity of standard antibiotics (gentamicin), the antimicrobial potential of the extracts was in the following order: roots > stems > seeds > leaves. Numerous secondary metabolites such as alkaloids, flavonoids, phenols, saponins, steroids, and others may be responsible for medicinal plants' therapeutic powers. Many fungi, bacteria, and viruses are inhibited by polyphenolic chemicals found in medicinal plants (Sodipo et al., 2000). A. Bose et al. (2007) reported that an ethanolic extract of *C. rutidosperma* revealed the presence of lipids, steroids, terpenoids, flavonoids, tannins, saponins and sugars. According to Donkor et al. (2014), alkaloids and tannins

presented in *C. viscosa* are associated with antibacterial activity. The presence of tannins, triterpenoids, and flavonoids in ethanol extract may explain its antibacterial properties. Tannins have been demonstrated to form irreversible compounds with prolene-rich proteins, resulting in the inhibition of cell wall synthesis (Chandrika & Chellaram, 2015; Mamtha et al., 2004).


### 3.5. Qualitative phytochemical analysis

Analysis of the extract yield showed that the *C. rutidosperma* leaves had the highest percentage compared to roots, stems and seeds (Table S1, Appendix A). A total of 13 qualitative phytochemical tests were carried out to detect their presence in plant extracts. The extracts contained alkaloids, flavonoids, phenols, sugars, proteins, saponins, sterols, tannins, and terpenoids, which conferred to the qualitative phytochemical screening study. Furthermore, glycosides were found in the stem, leaves, and roots, whereas coumarin was only found in the roots. On the other hand, quinones were found in none of the extracts (Table S1, Appendix A). Secondary metabolites found in *C. rutidosperma* samples include saponins, flavonoids, tannins, coumarins, and terpenoids, all of which have antibacterial and antifungal properties (Hayek et al., 2013).

### 3.6. Total phenolic and flavonoid contents

The therapeutic qualities of medicinal plants are due to secondary metabolites produced by the plants. Phenolic compounds (flavonoids, phenolic acids, quinones, phenylpropanoids, tannins, lignins, and hydroxyl compounds) are among the most diverse families of secondary metabolites found in natural products (Alu'datt et al., 2017; Nile et al., 2017). Table 1 shows the total phenolic and flavonoid content of different parts of ethanolic extracts from *C. rutidosperma*. The total phenolic content ranged between 37.230.62mg GAE/g DW and 70.451.23 mg GAE/g DW. Total phenolic content was significantly higher (p < 0.01) in leaves than in other parts tested. Phenolic chemicals are a diverse category of secondary metabolites that vary in distribution and concentration across and within plant species (Robards et al., 1999). Like phenol, the leaves had a significantly (p < 0.01) greater total flavonoid concentration. The total flavonoid content ranged between 20.110.89mg RE/g DW and 32.261.12mg RE/g DW. Ghosh et al. (2019) reported that the total phenolic (121.671.82 mg GAE/g DW) and flavonoids (65.080.98 mg QE/g DW) content

of ethanolic extracts of the same species was more significant than the concentration found in this study. According to *D. Silva et al. (2006)*, a higher concentration of phenolics and flavonoids in plant leaves might result from photosynthesis. Although no significant ( $p < 0.01$ ) variation was observed in total phenolic and flavonoid contents between the stem and seed extracts of *C. rutidosperma*. Polyphenolic compounds, such as total phenolic and total flavonoids, have been linked to many biological functions, including antioxidants (*Céspedes et al., 2010; Djeridane et al., 2006; Lamien-Meda et al., 2008; Meda et al., 2013*), antimicrobial activities (*Shan et al., 2007*). The capacity of phenolic compounds and flavonoids to scavenge hazardous free radicals and reactive species is well documented (*Hall & Cuppett, 1997; Jørgensen et al., 1999; Pietta, 2000; Rice-Evans et al., 1997*). Flavonoids play a significant role in both human and animal nutrition. They are a well-known class of natural chemicals that have been shown in various studies to have antioxidant and hepatoprotective properties (*Bratkov et al., 2016; Krasteva et al., 2016; Pistelli, 2002*). Plant secondary metabolites involve a range of biological actions, and therefore their regular consumption can have beneficial and harmful health repercussions (*Stobiecki & Kachlicki, 2006*).



**Figure 1.** Fourier Transform Infrared Spectroscopy (FTIR) spectrum of ethanolic extracts from *C. rutidosperma*

### 3.7. Fourier Transform Infrared Spectroscopy analysis

**Figure 1** and **Table S2, Appendix A** exhibit sample IR spectra analyses in the mid-infrared region ( $4000\text{--}400\text{ cm}^{-1}$ ) for aqueous ethanolic extracts from *C. rutidosperma*. For all plant parts examined, similar peaks were found at 2921, 1509, 1383, 1248, and  $1044\text{ cm}^{-1}$ . Absorbance bands at  $3420\text{--}3447\text{ cm}^{-1}$  were identified as polyphenols in the plant parts used in this study. The O-H stretching vibrations between  $3447\text{ and }3159\text{ cm}^{-1}$  confirmed the presence of water, alcohol, and phenols in the sample (*Larkin, 2011*). The C=O stretching vibration in

carbonyl compounds, defined by the presence of flavonoids in the samples, is ascribed to the absorption peak at  $1644\text{--}1636\text{ cm}^{-1}$  (*Socrates, 2001*). The absorption bands at  $2921\text{ cm}^{-1}$  and  $2925\text{ cm}^{-1}$  correspond to  $\text{CH}_2$  group C-H stretching, indicating the presence of several amino acids. The presence of aliphatic and aromatic CH groups in these compounds may also be indicated by these bands (*Ramamurthy & Kannan, 2007; Sivakesava & Irudayaraj, 2000*). The bands at 2925 and  $2921\text{ cm}^{-1}$  are caused by asymmetric stretching of the aromatic rings due to methyl ( $-\text{CH}_3$ ) and methylene ( $-\text{CH}_2-$ ) substitutions, respectively (*Socrates, 2001*). Absorption bands at  $1509\text{ cm}^{-1}$  suggest the existence of the benzene ring in aromatic chemicals in all plant parts of *C. rutidosperma*. The absorption band at  $1044\text{ cm}^{-1}$  in this study could be the stretching vibration of the C-O-C group of esters *Vlachos et al. (2006)*.

## 4. CONCLUSION

The plants under investigation can provide adequate protection against oxidative stress through complementary systems such as free radical scavenging metal ion reduction and antibacterial activity. More research into phytochemical compound isolation through chromatographic techniques demonstrating antioxidant, antimicrobial, and other pharmacological activity is required before being used as valuable additives in the nutraceutical and biopharmaceutical sectors.

## CONFLICTS OF INTEREST

The authors declare no competing interests.

## ACKNOWLEDGMENTS

The authors are grateful to the authorities of A.V.C. College (Autonomous), Tamil Nadu, India, for providing the facilities to conduct this study.

## ORCID

|                                   |                     |
|-----------------------------------|---------------------|
| Radjasagarin Arumugam             | 0000-0003-1980-1820 |
| Balachandravinayagam Elanchezhian | 0000-0002-2253-9205 |
| Jayakumar Samidurai               | 0000-0002-9509-5625 |
| Kaliyamoorthy Amirthagesan        | 0000-0003-3356-0234 |

## A. APPENDIX A. SUPPLEMENTARY INFORMATION

Supplementary information to this article can be found online at <https://doi.org/10.53365/nrfhh/146009>.

## REFERENCES

- Ahmed, I.A., Mikail, M.A., Ibrahim, M.B., Hazali, N.B., Rasad, M.S.B.A., Ghani, R.A., Wahab, R.A., Arief, S.J., Yahya, M.N.A., 2015. Antioxidant activity and phenolic profile of various morphological parts of underutilised *Baccaureaangulata* fruit. *Food Chemistry*. 172(1), 778–787. <https://doi.org/10.1016/j.foodchem.2014.09.122>
- Alu'datt, M.H., Rababah, T., Alhamad, M.N., Al-Mahasneh, M.A., Almajwal, A., Gammoh, S., Ereifej, K., Johargy, A., Alli, I., 2017. A review of phenolic compounds in oil-bearing plants: Distribution,

- identification and occurrence of phenolic compounds. *Food Chemistry*. 218(1), 99–106. <https://doi.org/10.1016/j.foodchem.2016.09.057> <https://doi.org/10.1016/j.foodchem.2016.09.057>
- Archi, F.F., Islam, S., Babu, M.A.H.K., Ullah, A., Azam, S., Chowdhury, A., Rahman, M., Karim, M.S., Goswami, S., 2016. Potential evaluation of central nervous system anti-depressant activity of *Cleome rutidosperma* in mice. *Biomedical Research and Therapy*. 3(10), 889–901.
- Arumugam, R., Kirkan, B., Sarikurkcu, C., 2019. Phenolic profile, antioxidant and enzyme inhibitory potential of methanolic extracts from different parts of *Astragalus ponticus* Pall. *South African Journal of Botany*. 120, 268–273. <https://doi.org/10.1016/j.sajb.2018.07.002>
- Arumugam, R., Sarikurkcu, C., Mutlu, M., Tepe, B., 2020. *Sophora alopecuroides* var. *alopecuroides*: Phytochemical composition, antioxidant and enzyme inhibitory activity of the methanolic extract of aerial parts, flowers, leaves, roots, and stems. *South African Journal of Botany*. 143, 282–290. <https://doi.org/10.1016/j.sajb.2020.10.009>
- Bajpai, V.K., Sharma, A., Kang, S.C., Baek, K.H., 2014. Antioxidant, lipid peroxidation inhibition and free radical scavenging efficacy of a diterpenoid compound sugiol isolated from *Metasequoia glyptostroboides*. *Asian Pacific Journal of Tropical Medicine*. 7(1), 9–15. [https://doi.org/10.1016/S1995-7645\(13\)60183-2](https://doi.org/10.1016/S1995-7645(13)60183-2)
- Barros, L., Ferreira, M.J., Queirós, B., Ferreira, I.C.F.R., Baptista, P., 2007. Total phenols, ascorbic acid,  $\beta$ -carotene and lycopene in Portuguese wild edible mushrooms and their antioxidant activities. *Food Chemistry*. 103(2), 413–419. <https://doi.org/10.1016/j.foodchem.2006.07.038>
- Bianco, A., Uccella, N., 2000. Biophenolic components of olives. *Food Res. Int.* 33(6), 72–79. [https://doi.org/10.1016/S0963-9969\(00\)00072-7](https://doi.org/10.1016/S0963-9969(00)00072-7)
- Bose, A., Mondal, S., Gupta, J.K., Ghosh, T., Si, S., Debbhuti, D., 2007. A study on antimicrobial activity of *Cleome rutidosperma* DC. *Journal of Natural Remedies*. 7(1), 132–134. <https://doi.org/10.18311/jnr/2007/205>
- Bose, U., Bala, V., Ghosh, T.N., Gunasekaran, K., Rahman, A.A., 2011. Antinociceptive, cytotoxic and antibacterial activities of *Cleome viscosa* leaves. *Rev. Bras. Farmacogn.* 21(1), 165–169. <https://doi.org/10.1590/S0102-695X2011005000023>
- Bratkov, V.M., Shkondrov, A.M., Zdraveva, P.K., Krasteva, I.N., 2016. Flavonoids from the genus *Astragalus*: Phytochemistry and biological activity. *Pharmacognosy Reviews*. 10(19), 11–32. <https://doi.org/10.4103/0973-7847.176550>
- Céspedes, C.L., El-Hafidi, M., Pavon, N., Alarcon, J., 2008. Antioxidant and cardioprotective activities of phenolic extracts from fruits of Chilean blackberry *Aristotelia chilensis* (Elaeocarpaceae), Maqui. *Food Chemistry*. 107(2), 820–829. <https://doi.org/10.1016/j.foodchem.2007.08.092>
- Céspedes, C.L., Valdez-Morales, M., Avila, J.G., El-Hafidi, M., Alarcón, J., Paredes-López, O., 2010. Phytochemical profile and the antioxidant activity of Chilean wild black-berry fruits, *Aristotelia chilensis* (Mol) Stuntz (Elaeocarpaceae). *Food Chemistry*. 119(3), 886–895. <https://doi.org/10.1016/j.foodchem.2009.07.045>
- Chandrika, M., Chellaram, C., 2015. Antibacterial properties of leaf extracts of *Borreria hispida* from Tamilandu regions, India. *Der Pharma Chemica*. 7, 389–394.
- Cotelle, N., Bernier, J.L., Catteau, J.P., Pommery, J., Wallet, J.C., Gaydou, E.M., 1996. Antioxidant properties of hydroxy-flavones. *Free Radical Biology and Medicine*. 20(1), 35–43. [https://doi.org/10.1016/0891-5849\(95\)02014-4](https://doi.org/10.1016/0891-5849(95)02014-4)
- Devi, B.P., Boominathan, R., Mandal, S.C., 2002. Evaluation of anti-diarrheal activity of *Cleome viscosa* L. extract in rats. *Phytomedicine*. 9(8), 739–742. <https://doi.org/10.1078/094471102321621368>
- Djeridane, A., Yousfi, M., Nadjemi, B., Maamri, S., Djireb, F., Stocker, P., 2006. Phenolic extracts from various Algerian plants as strong inhibitors of porcine liver carboxylesterase. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 21(6), 716–726. <https://doi.org/10.1080/14756360600810399>
- Donkor, A.M., Atindaana, E., Bugri, K.G., Atindaana, E.A., 2014. Evaluation of antibacterial potentiation of crude extracts of *Phyllanthus amarus*, *Tamarindus indica* and *Cleome viscosa* and their formulation. *International Journal of Plant Research*. 4, 23–28. [10.5923/j.plant.20140401.05](https://doi.org/10.5923/j.plant.20140401.05)
- Duan, X.J., Zhang, W.W., Li, X.M., Wang, B.G., 2006. Evaluation of antioxidant property of extract and fractions obtained from a red alga, *Polysiphonia urceolata*. *Food Chemistry*. 95(1), 37–43. <https://doi.org/10.1016/j.foodchem.2004.12.015>
- Farber, J.L., 1994. Mechanisms of cell injury by activated oxygen species. *Environmental Health Perspectives*. 102(10), 17–24. <https://doi.org/10.1289/ehp.94102s1017>
- Ghimire, B., Kumar, Seong, E.S., Kim, E.H., Ghimera, A.K., Yu, C.Y., Ghimire, Krishna, Chung, I.M., 2011. A comparative evaluation of the antioxidant activity of some medicinal plants popularly used in Nepal. *Journal of Medicinal Plant Research*. 5(10), 1884–1891.
- Ghosh, P., Biswas, M., Biswas, S., Dutta, A., 2019. Phytochemical Screening, Antioxidant and Antimicrobial Activity of leaves of *Cleome rutidosperma* DC. (Cleomaceae). *Journal of Pharmaceutical Sciences and Research*. 11(5), 1790–179.
- Hall, C.A., Cuppett, S.L., 1997. Antioxidant methodology: In vivo and in vitro concepts, Aruoma, O.I., Cuppett, S. L., (Eds.). AOCs Press, Illinois, pp. 141–172.
- Halliwell, B., Gutteridge, J.M.C., 1985. Free radicals in biology and medicine. *Journal of Free Radicals in Biology & Medicine*. 1(4), 331–332. [https://doi.org/10.1016/0748-5514\(85\)90140-0](https://doi.org/10.1016/0748-5514(85)90140-0)
- Hayek, S.A., Gyawali, R., Ibrahim, S.A., 2013. *Formatex*.Info.
- Jørgensen, L.V., Madsen, H.L., Thomsen, M.K., Dragsted, L.O., Skibsted, L.H., 1999. Regeneration of phenolic antioxidants from phenoxyl radicals: An ESR and electrochemical study of antioxidant hierarchy. *Free Radical Research*. 30(3), 207–220. <https://doi.org/10.1080/10715769900300231>
- Jun, H., Il, Kim, B.T., Song, G.S., Kim, Y.S., 2014. Structural characterization of phenolic antioxidants from purple perilla (*Perilla frutescens* var. *acuta*) leaves. *Food Chemistry*. 148(1), 367–372. <https://doi.org/10.1016/j.foodchem.2013.10.028>
- Krasteva, I., Shkondrov, A., Ionkova, I., Zdraveva, P., 2016. Advances in phytochemistry, pharmacology and biotechnology of Bulgarian *Astragalus* species. *Phytochemistry Reviews*. 15, 567–590. <https://doi.org/10.1007/s11101-016-9462-4>
- Lafka, T.I., Sinanoglou, V., Lazos, E.S., 2007. On the extraction and antioxidant activity of phenolic compounds from winery wastes. *Food Chemistry*. 104(3), 1206–1214. <https://doi.org/10.1016/j.foodchem.2007.01.068>
- Lamien-Meda, A., Lamien, C.E., Compaoré, M.M.Y., Meda, R.N.T., Kiendrebeogo, M., Zeba, B., Millogo, J.F., Nacoulma, O.G., 2008. Polyphenol content and antioxidant activity of fourteen wild edible fruits from Burkina Faso. *Molecules*. 13(3), 581–594. <https://doi.org/10.3390/molecules13030581>
- Larkin, P., 2011. Infrared and Raman Spectroscopy Principles and Spectral Interpretation, and others, (Eds.). Elsevier, Waltham, MA, USA, p. 228. <https://doi.org/10.1016/C2010-0-68479-3>
- Li, Y., Guo, C., Yang, J., Wei, J., Xu, J., Cheng, S., 2006. Evaluation of antioxidant properties of pomegranate peel extract in comparison

- with pomegranate pulp extract. *Food Chemistry*. 96(2), 254–260. <https://doi.org/10.1016/j.foodchem.2005.02.033>
- Mamtha, B., Kavitha, K., Srinivasan, K.K., Shivananda, P.G., 2004. An in vitro study of the effect of *Centella asiatica* [Indian pennywort] on enteric pathogens. *Indian Journal of Pharmacology*. 36(1), 41.
- Marya, B., Dattani, K.H., Patel, D.D., Patel, P.D., Patel, D., Suthar, M.P., Patel, V.P., Bothara, S.B., 2011. In-vitro cytotoxicity evaluation of aqueous fruit and leaf extracts of *Grewia asiatica* using MTT assay. *Der Pharma Chemica*. 3.
- McNeil, M.J., Porter, R.B.R., Rainford, L., Dunbar, O., Francis, S., Laurieri, N., Delgoda, R., 2018. Chemical composition and biological activities of the essential oil from *Cleome rutidosperma* DC. *Fitoterapia*. 129, 191–197. <https://doi.org/10.1016/j.fitote.2018.07.006>
- Meda, N.T.R., Bangou, M.J., Bakasso, S., Millogo-Rasolodimby, J., Nacoulma, O.G., 2013. Antioxidant activity of phenolic and flavonoid fractions of *Cleome gynandra* and *Maerua angolensis* of Burkina faso. *Journal of Applied Pharmaceutical Science*. 3(2), 36–42.
- Molyneux, P., 2004. The Use of the Stable Free Radical Diphenylpicrylhydrazyl (DPPH) for Estimating Antioxidant Activity. *Songklanakarin Journal of Science and Technology*. 26(2), 211–219.
- Motaal, A.A., Ezzat, S.M., Haddad, P.S., 2011. Determination of bioactive markers in *Cleome droserifolia* using cell-based bioassays for antidiabetic activity and isolation of two novel active compounds. *Phytomedicine*. 19(1), 38–41. <https://doi.org/10.1016/j.phymed.2011.07.003>
- Namvar, K., Mohammadi, A., Salehi, E.A., Feyzi, P., 2017. Evaluation of Solvent Effect (Methanol: Water Mixture) on the Phenolic Content and Antioxidant Activities of *Stachys turcomanica* Trautv. *Pharmaceutical Sciences*. 23(3), 244–248. <https://doi.org/10.15171/PS.2017.36>
- Nile, S.H., Nile, A.S., Keum, Y.S., 2017. Total phenolics, antioxidant, antitumor, and enzyme inhibitory activity of Indian medicinal and aromatic plants extracted with different extraction methods. 3 *Biotech*. 7, 76. <https://doi.org/10.1007/s13205-017-0706-9>
- Panovska, T.K., Kulevanova, S., Stefova, M., 2005. In vitro antioxidant activity of some *Teucrium* species (Lamiaceae). *Acta Pharmaceutica*. 55(2), 207–214.
- Parimaladevi, B., Boominathan, R., Mandal, S.C., 2003. Studies on analgesic activity of *Cleome viscosa* in mice. *Fitoterapia*. 74(3), 262–266. [https://doi.org/10.1016/s0367-326x\(03\)00020-0](https://doi.org/10.1016/s0367-326x(03)00020-0)
- Pietta, P.G., 2000. Flavonoids as antioxidants. *Journal of Natural Products*. 63(7), 1035–1042. <https://doi.org/10.1021/np9904509>
- Pistelli, L.F., 2002. Secondary metabolites of genus *Astragalus*: Structure and biological activity. *Studies in Natural Products Chemistry*. 27, 443–545. [https://doi.org/10.1016/S1572-5995\(02\)80043-6](https://doi.org/10.1016/S1572-5995(02)80043-6)
- Ramamurthy, N., Kannan, S., 2007. Fourier Transform Infrared Spectroscopic analysis of a plant (*Calotropis gigantea* linn) from an industrial village. *J. Biophys*.
- Rice-Evans, C.A., Sampson, J., Bramley, P.M., Holloway, D.E., 1997. Why do we expect carotenoids to be antioxidants in vivo? *Free Radical Research*. 26(4), 381–398. <https://doi.org/10.3109/10715769709097818>
- Robards, K., Prenzler, P.D., Tucker, G., Swatsitang, P., Glover, W., 1999. Phenolic compounds and their role in oxidative processes in fruits. *Food Chemistry*. 66(4), 401–436. [https://doi.org/10.1016/S0308-8146\(99\)00093-X](https://doi.org/10.1016/S0308-8146(99)00093-X)
- Shan, B., Cai, Y.Z., Brooks, J.D., Corke, H., 2007. The in vitro antibacterial activity of dietary spice and medicinal herb extracts. *International Journal of Food Microbiology*. 117(1), 112–119. <https://doi.org/10.1016/j.ijfoodmicro.2007.03.003>
- Silva, D., Souza, J.F.M.D., Matta, M.C., Andrade, S.R.D., Vidal, M.R., N, F.V., 2006. Correlation analysis between phenolic levels of Brazilian propolis extracts and their antimicrobial and antioxidant activities. *Food Chemistry*. 99(3), 431–435. <https://doi.org/10.1016/j.foodchem.2005.07.055>
- Silva, N.D., Fonseca, L.C.M.D., Araújo, C.S.D., Correia, J.M., Do, M.T.S., Cavalcanti, M., de Menezes Lima, V.L., 2016. Antimicrobial Activity and Phytochemical Analysis of Organic Extracts from *Cleome spinosa* Jacq. *Frontiers in Microbiology*. 7, 963. <https://doi.org/10.3389/fmicb.2016.00963>
- Silva, O.A., Oliveira, F.R.D., De, A.M., Lima, T.C., Sousa, D.P.D., Souza, A.A.D., Freitas, R.M.D., 2012. Evaluation of the antioxidant effects in vitro of the isopulegone. *Free Radicals and Antioxidants*. 2(4), 50–55. <https://doi.org/10.5530/ax.2012.4.9>
- Sivakesava, S., Irudayaraj, J., 2000. Analysis of potato chips using FTIR photoacoustic spectroscopy. *Journal of the Science of Food and Agriculture*. 80, 1805–1810. [https://doi.org/10.1002/1097-0010\(20000915\)80:12%3C1805::AID-JSFA710%3E3.0.CO;2-L](https://doi.org/10.1002/1097-0010(20000915)80:12%3C1805::AID-JSFA710%3E3.0.CO;2-L)
- Socrates, G., 2001. Infrared and Raman Characteristic Group Frequencies: Tables and Charts, In: 3rd (Eds.); and others, (Eds.). John Wiley & Sons, LTD, p. 364.
- Sodipo, O.A., Akinniyi, J.A., Ogunbameru, J.V., 2000. Studies on certain characteristics of extracts of bark of *Pausinystalia johimbe* and *Pausinystalia macroceras* (K Schum) Pierre ex Beille. *Global Journal of Pure and Applied Sciences*. 6(1), 83–87. <https://doi.org/10.4314/gjpas.v6i1.16082>
- Stobiecki, M., Kachlicki, P., 2006. Isolation and identification of flavonoids, G. E. et al., (Eds.), *The Science of Flavonoids*. Springer, New York, pp. 47–69. [https://doi.org/10.1007/978-0-387-28822-2\\_2](https://doi.org/10.1007/978-0-387-28822-2_2)
- Thomas, B., Arumugam, R., Veerasamy, A., Ramamoorthy, S., 2014. Ethnomedicinal plants used for the treatment of cuts and wounds by Kuruma tribes, Wayanadu districts of Kerala, India. *Asian Pacific Journal of Tropical Biomedicine*. 4(1), S488–S491. <https://doi.org/10.12980/apjtb.4.2014b571>
- Venkatesh, B., Dorai, A., 2011. Antibacterial and Antioxidant potential of White and Pink *Nelumbo Nucifera* Gaertn Flowers. *International Conference on Bioscience, Biochemistry and Bioinformatics*. 5, 213–217.
- Vlachos, N., Skopelitis, Y., Psaroudaki, M., Konstantinidou, V., Chatzilarou, A., Tegou, E., 2006. Applications of Fourier transform-infrared spectroscopy to edible oils. *Analytica Chimica Acta*. 573-574, 459–465. <https://doi.org/10.1016/j.aca.2006.05.034>
- Yingming, P., Ying, L., Hengshan, W., Min, L., 2004. Antioxidant activities of several Chinese medicine herbs. *Food Chemistry*. 88(3), 347–350. <https://doi.org/10.1016/j.foodchem.2004.02.002>