

Original Research

View Article Online



Received 10 January 2022

Revised 18 January 2022

Accepted 28 January 2022

Available online 07 February 2022

Edited by Carlos L. Cespedes Acuña

KEYWORDS:

ABTS

DPPH

flavonoid content

FRAP

Plasmodium falciparum

Natr Resour Human Health 2022; (0): 1-7

<https://doi.org/10.53365/nrhh/146223>

eISSN: 2583-1194

Copyright © 2022 Visagaa Publishing House

Antimalarial, cytotoxic and antioxidant activities of 14 medicinal plants from Sudan

Ibrahim B. E. El Bashir¹, Maha Y. Kordofani¹, Sakina Yagi¹, Abdulrahman A. Al-Atar², Ahmed A. Qahtan², Eslam M. Abdel-Salam², Abdullah M. Alkahtani², Majed A. Al-Mansoub³, Ammar Mohammed Ahmed Ali⁴, Ali M. Batah⁵, Gokhan Zengin^{6,*}

¹Department of Botany, Faculty of Science, University of Khartoum P.O. Box 321, Khartoum, Sudan

²Department of Botany and Microbiology, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Saudi Arabia

³Department of Chemistry, College of Science, Ibb University, P.O.70270, Ibb, Yemen

⁴Department of Biology, Faculty of Education, Hajjah University, Hajjah, Yemen

⁵Tropical Disease Research Center, University of Science and Technology, Sana'a, Yemen

⁶Physiology and Biochemistry Laboratory, Department of Biology, Science Faculty, Selcuk University, Campus, 42103 Konya, Turkey

ABSTRACT: Fourteen plants endogenous to Sudan were selected to evaluate their total phenolic content, antioxidant, antimalarial and cytotoxicity potential in the present work. Extracts were prepared by maceration of each plant material in chloroform: methanol (C: M; 1:1 v/v) mixture. The antioxidant activity was determined by measuring the radical scavenging effects against 2,2-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS), and ferric-reducing (FRAP) capacity. The antiplasmodial activity was determined using the NF54 strain of *Plasmodium falciparum*. Cytotoxicity was evaluated towards human colorectal carcinoma cell line (HCT-116), human hormone-sensitive and invasive breast cancer cell line (MCF-7), and human hormone-resistant breast cancer cell line (MDA-MB-231), in addition to endothelial normal EAhy-296 cell line. Results showed that the plants' total phenolic and flavonoid contents were variable. Of the 14 plant species, only *Burnatia enneandra* showed high *in vitro* antiplasmodial activity (IC₅₀ 5758 ng/mL). Some plants possessed considerable free radical scavenging ability and reducing power. *Coccinia grandis* fruit extract (IC₅₀ 13.23 ± 0.51 µg/mL) and *Geigeria alata* root extract (IC₅₀ 35.54 ± 0.27 µg/mL) displayed the highest DPPH and ABTS scavenging activity, respectively. *Striga hermonthica* whole plant extract exhibited the highest FRAP reducing power (107.15 ± 0.11 nmol Fe⁺²eq./mg). At a 100 µg/mL concentration, *Dioscorea hirtiflora* bulb extract displayed the highest cytotoxicity (74.23 ± 03.72%), followed by *Mitragyna inermis* fruit extract (65.28 ± 04.60%) against HCT-116 cell line. *Aerva javanica* leaf extract showed toxicity to the MDA-MB-231 cell line (50.82 ± 07.46%) at 100 µg/mL. The current study results showed that endogenous medicinal plants might represent a rich source of natural antioxidant, antimalarial and antitumor agents.

1. INTRODUCTION

Chronic diseases represent a significant cause of increased mortality and morbidity cases worldwide. The number of cancer patients is expected to reach 24 million in 2035, as estimated by the International Agency for Research on Cancer (IARC) (Bray et al., 2013). In Sudan, although there is no national, systematic screening programme for cancer registry,

the number of diagnosed cancer cases is increasing (Elamin et al., 2015; Elebead et al., 2012). Furthermore, the number of malaria patients was estimated by the WHO as 229 million cases worldwide in 2019, with more than 1.8 million cases from Sudan (OCHA, 2019; WHO, 2020) (WHO, 2020; OCHA, 2019). Several oxidative stress-induced pathological disorders and disease conditions, including diabetes, cancers,

* Corresponding author.

E-mail address: gokhanzengin@selcuk.edu.tr (Gokhan Zengin)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1

Profile of the investigated plants

Name of Plant	Family	Voucher No	Ailment treated in traditional medicine	References
<i>Aerva javanica</i> (Burm.f.) Juss. ex Schult.	Amaranthaceae	94/19	Wounds, cough, diarrhoea, ulcer, hyperglycaemia	(Ghazali et al., 1997)
<i>Burnatia enneandra</i> Micheli	Alismataceae	2011/BE	Wounds, intestinal parasites	(I.G. Doka, 2012)
<i>Cissus quadrangularis</i> L.	Vitaceae	2014/CQ	Syphilis, dandruff, back pain, wound, rheumatism	(Ghazali et al., 1997; Issa et al., 2018)
<i>Coccinia grandis</i> (L.) Voigt	Cucurbitaceae	1109KD7	Diabetes, cataracts, skin eruptions	(I.G. Doka, 2012)
<i>Dioscorea dumetorum</i> (Kunth) Pax	Dioscoreaceae	1109KD3	Rheumatoid, arthritis	(I.G. Doka, 2012)
<i>Dioscorea hirtiflora</i> Benth.	Dioscoreaceae	1109KD4	Malaria, stomach troubles	(I.G. Doka, 2012)
<i>Fagonia arabica</i> L.	Zygophyllaceae	95/13	Skin allergy	(Issa et al., 2018)
<i>Geigeria alata</i> Benth. & Hook.f. ex Oliv. & Hiern	Compositae	41935/HNC	Diabetes	(Issa et al., 2018)
<i>Leptadenia arborea</i> (Forssk.) Schweinf.	Apocynaceae	99/8	Snake bits	(Ghazali et al., 1997)
<i>Maerua pseudopetalosa</i> (Gilg & Gilg-Ben.) DeWolf	Capparaceae	83/107	Diabetes, sexual debility, hypertension, kidney disorders	(Issa et al., 2018)
<i>Mitragyna inermis</i> (Willd.) Kuntze	Rubiaceae	2013-12/HBD	Diabetes	(Alamin et al., 2015)
<i>Raphionacme splendens</i> Schltr.	Apocynaceae	1109KR6	Famine food	(I. Doka & Yagi, 2009)
<i>Striga hermonthica</i> (Delile) Benth.	Orobanchaceae	2013-16/HBD	Menstrual cramps, diabetes	(Issa et al., 2018)
<i>Tephrosia apollinea</i> (Delile) DC.	Leguminosae	94/16	Swelling, bone fracture, cough, earache, wounds	(Hassan et al., 2014)

aging, neurological, and cardiovascular disorders, generate free radicals (Ramana et al., 2018). Antioxidants play key roles in counteracting oxidative stress by protecting cells against the effects of free radicals. Natural antioxidants are generally categorized into dietary sources, including vegetables, fruits, cereals, and beans and those from medicinal plants and wild herbs (Wilson et al., 2017). Studies have indicated that antioxidants have a high potential to inhibit the development of many diseases, including cancer (Ramana et al., 2018). Nowadays, the pharmaceutical industries are searching for potential natural and safe raw materials to overcome the side effects of synthetic drugs and problems related to drug-resistant (Abdin et al., 2020). Many investigated plant extracts and isolated phyto-components have a wide range of pharmacological properties and could be used as the safest, most effective and natural source instead of chemical drugs. Karunakaran et al. (2020) isolated two anticancer compounds (ananixanthone and trapezifolixanthone) from *Calophyllum macrocarpum* and proved their cytotoxicity against HeLa Chang liver and HEK-293 cell lines. Also, some Omani *Haplophyllum tuberculatum* species recorded potent antioxidant and intense cytotoxic activities (Al-Muniri & Hossain, 2017). Furthermore, several examined plant extracts exhibited considerable effect against Plasmodium parasites, as displayed by *Syzygium guineense* and *Parinari congensis* extracts (Laryea and Borquaye, 2019) and evaluated by leaf extract of *Fadogia cienkowskii* (Orabueze et al., 2019). Therefore, plants are considered a potential natural source for new lead

antimalarial and antitumor compounds and antioxidant agents. The diversity of climates in Sudan results in a wide variety of flora species which comprises about 3137 plant species belonging to 170 families and 1280 genera (H. Khalid et al., 2012). Considering all the issues mentioned above and in an attempt to explore plant-based alternative solutions in promoting health, the objective of this study was to examine 14 medicinal plants, traditionally used in Sudan to cure different diseases, for their antioxidant, antimalarial, and cytotoxic potential and to determine their total phenolic and flavonoid contents. These plants are; *Aerva javanica* (Burm.f.) Juss. ex Schult., *Burnatia enneandra* Micheli, *Cissus quadrangularis* L., *Coccinia grandis* (L.) Voigt, *Dioscorea dumetorum* (Kunth) Pax, *Dioscorea hirtiflora* Benth., *Fagonia arabica* L., *Geigeria alata* Benth. & Hook.f. ex Oliv. & Hiern, *Leptadenia arborea* (Forssk.) Schweinf., *Maerua pseudopetalosa* (Gilg & Gilg-Ben.) DeWolf, *Mitragyna inermis* (Willd.) Kuntze, *Raphionacme splendens* Schltr., *Striga hermonthica* (Delile) Benth., and *Tephrosia apollinea* (Delile) DC. The identities and traditional medicinal uses of the investigated plants are compiled in Table 1.

2. MATERIALS AND METHODS

2.1. Plant materials

Samples of the selected plants were collected from different regions in Sudan. The plants were identified and authenticated at the Department of Botany, Faculty of Science, University of Khartoum, by Maha Y. Kordofani. Voucher numbers are given

in Table 1.

2.2. Preparation of extracts

Samples were dried in the shade at ambient temperature and then grounded into fine powder. Fifty grams of each plant material was macerated in chloroform: methanol (1:1 v/v) solvent for 24 hours, then filtrated and concentrated using a rotary evaporator under reduced pressure and weighed.

2.3. Determination of total phenolic (TPC) and (total flavonoids (TFC) contents

The TPC and TFC were determined by adopting the method described by Wolfe et al. (2003) and Ordoñez et al. (2006). The detailed procedures were given in the Appendix A.

2.4. Pharmacological evaluation

Antioxidant activity of the extracts was estimated using DPPH in vitro method (K.J. Wang et al., 2007), ABTS radical-scavenging activity (Re et al. (1999) Re et al. (1999) and Ferric reducing antioxidant power (FRAP) (Benzie and Strain (1996) Benzie and Strain (1996). The antiplasmodial activity was determined by the in vitro microplate assay against the NF54 strain of *Plasmodium falciparum* (sensitive to all known drugs). A modification of the [3H] hypoxanthine incorporation assay was used (Witschel et al., 2012). Cytotoxicity of the extracts, at concentrations of 50 and 100 µg/mL, was performed as described by Mosmann (1983) using the thiazolyl blue tetrazolium bromide (MTT) procedure. The detailed pharmacological procedures were given in the Appendix A.

2.5. Statistical Analysis

All experiments were performed in triplicate, and the results were expressed as mean ± standard deviation (SD) values. Significant differences between samples were analyzed using analysis of variance (ANOVA) and Duncan's multiple range test ($P < 0.05$).

3. RESULTS AND DISCUSSION

Plant bioactivity and medicinal aspects are usually linked with various classes of secondary phytochemicals such as phenols, flavonoids, terpenes, etc. Their presence and concentration vary from one plant to another and even from part to another in the same plant, mainly used in Ethnobotany (Kumar et al., 2019).

3.1. Phenolic content

Phenolics are well recognized to exert several biological properties, including antioxidant, antiproliferation, and anti-carcinogenic activities (Wilson et al., 2017). Accordingly, the total polyphenolics and flavonoids contents of the studied plant's extracts were determined (Table 2). The values of total polyphenolic content were ranged between 11.78 ± 0.07 to 94.43 ± 0.63 µg gallic acid eq./mg extract. *G. alata* exhibited the highest phenolic content, followed by *D. hirtiflora* (66.45 ± 0.30 µg gallic acid eq./mg), *L. arborea* (stem) (57.25 ± 1.37 µg

gallic acid eq./mg), *C. quadrangularis* (56.01 ± 0.64 µg gallic acid eq./mg), *M. inermis* (51.25 ± 0.51 µg gallic acid eq./mg) and *S. hermonthica* (51.05 ± 0.07 µg gallic acid eq./mg). Other species extracts had total polyphenolic content of less than 40 µg gallic acid eq./mg.

The total flavonoid content values in the extracts ranged between 2.63 ± 0.03 to 31.81 ± 0.74 µg quercetin eq./mg extract. The extract of *S. hermonthica* showed the highest flavonoid content followed by that of *G. alata* (30.20 ± 0.79 µg quercetin eq./mg extract), *L. arborea* (stem) (25.32 ± 1.66 µg quercetin eq./mg extract), *L. arborea* (leaf) (24.88 ± 0.41 µg quercetin eq./mg extract), *C. quadrangularis* (24.52 ± 0.65 µg quercetin eq./mg extract), *Aerva javanica* (22.86 ± 0.95 µg quercetin eq./mg extract) and *D. dumetorum* (19.92 ± 0.26 µg quercetin eq./mg extract) respectively. Other species extracts showed total flavonoid content < 11 µg quercetin eq./mg extract.

3.2. Antioxidant activity

The antioxidant capacity for the selected 14 plant extracts was evaluated using three complementary assays; DPPH, ABTS and FRAP (Table 2). Different extracts exhibited various free radical scavenging activities, as indicated by their IC₅₀ values. In the DPPH assay, the IC₅₀ values of plants extracts ranged from 13.23 to 1233.24 µg/mL. *C. grandis* displayed the highest DPPH scavenging activity, followed by *S. hermonthica* (IC₅₀ 28.14 ± 0.25 µg/mL), *D. hirtiflora* (IC₅₀ 47.28 ± 3.24 µg/mL), *M. inermis* (IC₅₀ 66.69 ± 1.12 µg/mL), *G. alata* (IC₅₀ 68.54 ± 0.71 µg/mL) respectively. The IC₅₀ values of different plant extracts ranged from 35.54 to 712.72 µg/mL from the ABTS assay. The highest ABTS scavenging activity was obtained from *G. alata*, followed by *D. hirtiflora* (IC₅₀ 59.9 µg/mL), *S. hermonthica* (IC₅₀ 71.21 µg/mL) and *M. inermis* (IC₅₀ 73.19 µg/mL) respectively. From the FRAP assay values (were ranged from 43.24 ± 0.62 to 107.15 ± 0.11 nmol Fe⁺² eq./mg where *S. hermonthica* exhibited the highest FRAP reducing power (107.15 ± 0.11 nmol Fe⁺² eq./mg), followed by *B. enneandra* (106.31 ± 0.92 nmol Fe⁺² eq./mg), *A. javanica* (100.71 ± 0.4992 nmol Fe⁺² eq./mg), *L. arborea* (stem) (99.76 ± 3.42 nmol Fe⁺² eq./mg), *C. quadrangularis* (96.38 ± 2.69 nmol Fe⁺² eq./mg), *G. alata* (83.13 ± 0.86 nmol Fe⁺² eq./mg) and *M. inermis* (82.69 ± 1.18 nmol Fe⁺² eq./mg) respectively.

It was clear that extracts displayed variable antioxidant capacity in different assays. Although both ABTS and DPPH assays are associated with electron and radical scavenging, the studied extracts revealed variable capacity in both assays. This observation has been reported previously, where it was suggested that factors like stereo-selectivity of the radicals or the solubility of the extracts in different testing systems affect the capacity of extracts to react and quench other radicals (Jimoh et al., 2010). The observed high antioxidant activity of some plant extracts may be attributed to high concentrations of phenolic compounds with potential protective effects (Spanou et al., 2010). Furthermore, these findings supported previous results on the antioxidant activity of *S. hermonthica* (Kiendrebeogo

Table 2

Phenolic contents and antioxidant activity of extracts of the investigated plants.

Name of Plant	Part used	TPC ^A	TFC ^B	DPPH ^C	ABTS ^C	FRAP ^D
<i>Aerva javanica</i>	leaf	39.96 ± 0.36 ^d	22.86 ± 0.95 ^b	154.09 ± 4.49 ^d	133.67 ± 7.50 ^e	100.71 ± 0.49 ^b
<i>Burnatia enneandra</i>	Fruit	32.01 ± 0.30 ^d	10.88 ± 6.60 ^d	183.05 ± 4.32 ^d	162.26 ± 14.94 ^e	106.31 ± 0.92 ^a
<i>Cissus quadrangularis</i>	Whole plant	56.01 ± 0.64 ^b	24.52 ± 0.65 ^b	89.02 ± 2.80 ^c	82.72 ± 4.12 ^c	96.38 ± 2.69 ^b
<i>Coccinia grandis</i>	Fruit	31.35 ± 0.47 ^d	04.95 ± 0.10 ^e	13.23 ± 0.51 ^a	181.81 ± 12.80 ^f	60.44 ± 1.49 ^e
<i>Dioscorea dumetorum</i>	Tuber	22.94 ± 0.24 ^e	19.92 ± 0.26 ^c	215.15 ± 16.03 ^e	299.84 ± 24.87 ^g	69.15 ± 0.16 ^d
<i>Dioscorea hirtiflora</i>	Tuber	66.45 ± 0.30 ^b	08.32 ± 0.57 ^d	47.28 ± 3.24 ^b	59.90 ± 7.50 ^b	78.61 ± 1.32 ^c
<i>Fagonia arabica</i>	Whole plant	39.18 ± 1.14 ^d	09.43 ± 0.13 ^d	272.67 ± 6.82 ^e	98.95 ± 4.32 ^d	55.96 ± 0.33 ^e
<i>Geigeria alata</i>	Root	94.43 ± 0.63 ^a	30.20 ± 0.79 ^a	68.54 ± 0.71 ^b	35.54 ± 0.27 ^a	83.13 ± 0.86 ^c
<i>Leptadenia arborea</i>	Stem	57.25 ± 1.37 ^b	25.32 ± 1.66 ^b	88.28 ± 1.70 ^c	94.85 ± 3.41 ^d	99.76 ± 3.42 ^b
	leaf	30.57 ± 0.07 ^d	24.88 ± 0.41 ^b	127.78 ± 3.89 ^d	137.07 ± 1.67 ^e	71.50 ± 1.40 ^d
<i>Maerua pseudopetalosa</i>	Seed	11.78 ± 0.07 ^f	8.84 ± 0.20 ^d	1233.24 ± 28.3 ^g	482.24 ± 136.9 ^h	43.24 ± 0.62 ^f
<i>Mitragyna inermis</i>	Leaf	51.25 ± 0.51 ^c	10.94 ± 0.97 ^d	66.69 ± 1.12 ^b	73.19 ± 1.90 ^c	82.69 ± 1.18 ^c
<i>Raphionacme splendens</i>	Root	27.31 ± 0.30 ^e	02.63 ± 0.03 ^e	197.35 ± 26.02 ^d	182.41 ± 4.72 ^f	61.98 ± 0.23 ^e
<i>Striga hermonthica</i>	Whole plant	51.05 ± 0.07 ^c	31.81 ± 0.74 ^a	28.14 ± 0.25 ^a	71.21 ± 1.28 ^c	107.15 ± 0.11 ^a
<i>Tephrosia apollinea</i>	Seed	12.89 ± 0.26 ^f	01.54 ± 0.01 ^f	423.73 ± 9.11 ^f	712.72 ± 185.1 ⁱ	51.01 ± 0.57 ^e

^A, Total polyphenolic content (μg gallic acid eq/mg extract); ^B, Total flavonoid content (μg quercetin eq./mg extract); ^C, IC₅₀ ($\mu\text{g}/\text{mL}$); ^D, nmol Fe²⁺ eq./mg extract. Extract: chloroform:methanol (1:1). Different superscript letters in the same column indicate significant difference ($p < 0.05$).

et al., 2005), *C. grandis* (Deshpande et al., 2011), *C. quadrangularis* (Prabhavathi et al., 2016; Tiwari et al., 2021), *D. hirtiflora* (Adeniran et al., 2020; Sonibare & Abegunde, 2012), *G. alata* (Zheleva-Dimitrova et al., 2017) and *M. inermis* (Ouedraogo et al., 2020) using different extraction solvents.

3.3. Antimalarial activity

In vitro antimalarial activity was measured using the chloroquine-sensitive NF54 strains of *Plasmodium falciparum* (Table 3). Of the 14 plant species, only *B. enneandra* showed considerable in vitro antiplasmodial activity with an IC₅₀ value of 5758 ng/mL. However, except for determining total phenolic content, this plant has never been investigated for its secondary metabolite content. Other plants extract showed IC₅₀ value >10 000 ng/mL. However, some of these plant species are reported to be used to treat malaria in other countries and were found to exert antiplasmodial activity in other plasmodial strains (Philip et al., 2017; Tefera & Kim, 2019).

For example, leaves of *M. inermis* is used to treat malaria in Guinea, and an antiplasmodial activity test on strain W2 showed that it has potent activity with an IC₅₀ value of 4.32 $\mu\text{g}/\text{mL}$ (Traoré et al., 2015). Leaf of *C. grandis* possessed antiplasmodial activity against *P. falciparum* with an inhibition value of 69 % (Ravikumar et al., 2012). *S. hermonthica* exhibited a high intrinsic antimalarial activity (68.5% suppression of parasitaemia) against *P. berghei* in mice (Okpako & Ajaiyeoba, 2004). Dichloromethane extract of *C. quadrangularis* whole plant had moderate (IC₅₀ 23.9 $\mu\text{g}/\text{mL}$) antiplasmodial activity against *P. falciparum* chloroquine-sensitive strain 3D7 (Bah et al., 2007). Leaves and roots of *D. dumetorum* are commonly used to treat malarias in Southwest Negeria (Dike et al., 2012), but no scientific evidence was found. Moreover, pure compounds, tephroglabrin, (-)-semiglabin, semiglabinone, (+)-glabratephrin isolated from *T. apollinea* were found to be

Table 3In vitro inhibition of plant species extracts under the study against *Plasmodium falciparum* strain NF54.

Name of Plant	Part used	IC ₅₀ NF54 (ng/mL)
<i>Aerva javanica</i>	leaf	> 10 000
<i>Burnatia enneandra</i>	Fruit	5758
<i>Cissus quadrangularis</i>	Whole plant	> 10 000
<i>Coccinia grandis</i>	Fruit	> 10 000
<i>Dioscorea dumetorum</i>	Tuber	> 10 000
<i>Dioscorea hirtiflora</i>	Tuber	> 10 000
<i>Fagonia arabica</i>	Whole plant	> 10 000
<i>Geigeria alata</i>	Root	> 10 000
<i>Leptadenia arborea</i>	Stem	> 10 000
	leaf	> 10 000
<i>Maerua pseudopetalosa</i>	Seed	> 10 000
<i>Mitragyna inermis</i>	Leaf	> 10 000
<i>Raphionacme splendens</i>	Root	> 10 000
<i>Striga hermonthica</i>	Whole plant	> 10 000
<i>Tephrosia apollinea</i>	Seed	> 10 000
Chloroquine	Standard	2500
Artesunate	Standard	1400

*Extract: chloroform:methanol (1:1).

inactive against *P. falciparum* (S.A. Khalid et al., 1986). Thus, it could be concluded that the antimalarial activity of plant extracts depends on many factors, among them the type of extract and plasmodial strains.

3.4. Cytotoxicity

The cytotoxicity of the 15 plant extracts against three cancerous cell lines, HCT 116, MCF 7 and MDA-MB-231, in addition, to one human normal cell line, EAhy 926, were determined after treatment with 50 and 100 $\mu\text{g}/\text{mL}$ of extracts for 48 hours (Figure 1). Cytotoxicity of extracts against HCT 116 showed that at 50 $\mu\text{g}/\text{mL}$, *D. hirtiflora* displayed




Figure 1. Cytotoxicity of 15 crude extracts of investigated plants towards HCT-116, MCF-7, MDA-MB-231 and EAhy-296 cells lines at concentrations 50 and 100 µg/mL for 48 hours. A.j, *Aerva javanica*; B.e, *Burnatia enneandra*; C.q, *Cissus quadrangularis*; C.g, *Coccinia grandes*; D.d, *Dioscorea dumetorum*; D.h, *Dioscorea hirtiflora*; F.a, *Fagonia arabica*; G.a, *Geigeria alata*; L.s & L.l, *Leptadenia arborea* (s:stem, l:leaf); M.p, *Maerua pseudopetalosa*; M.i, *Mitragyna inermis*; R.s, *Raphionacme splendens*; S.h, *Striga hermonthica*; T.a, *Tephrosia apollinea*. Different superscript letters indicate significant difference ($p < 0.05$).

cytotoxicity with inhibition values of $59.59 \pm 13.03\%$. At 100 µg/mL, the same extract displayed the highest cytotoxicity ($74.23 \pm 03.72\%$), followed by those of *M. inermis* ($65.28 \pm 04.60\%$) and *B. enneandra* ($64.79 \pm 02.05\%$) respectively (Figure 1-a). All the 15 studied extracts did not exert cytotoxicity against the MCF-7 cells line at both concentrations used (Figure 1-b). Extract of *A. javanica* was the only extract with toxicity against the MDA-MB-231 cell line with an inhibition value equal to $50.82 \pm 07.46\%$ at 100 µg/mL (Figure 1-c). Furthermore, all extracts were not toxic to human normal cell line EAhy-926 at 50 µg/mL. Nevertheless, at 100 µg/mL, *B. enneandra* and *T. apollinea* extracts were relatively toxic ($60.04 \pm 01.21\%$ and $57.07 \pm 03.11\%$, respectively), while all other extracts were not toxic (Figure 1-d).

The results of cytotoxicity screening of the present study revealed that *D. hirtiflora* and *M. inermis* displayed interesting cytotoxic effects towards HCT-116. Both plant species were not toxic to the normal EAhy-926 cell line. No published data on the antitumor activity of *D. hirtiflora*, *A. javanica* and *B. enneandra* were found. However, the antitumor effect of other *Dioscorea* spp like *D. bulbifera* (J.M. Wang et al., 2012), *D. esculenta* (Murugan & Mohan, 2012), and *D. opposita* (Liu et al., 2016; Zhang et al., 2016) was reported. A previous study on *M. inermis* showed that the alkaloid rich extract did not exert mutagenic or genotoxic activity (Traore et al., 2000)

4. CONCLUSION

In conclusion, the results obtained in this study showed that total polyphenolic and flavonoid contents and antioxidant activity differ among extracts of the studied medicinal plants. Some of these medicinal plants are promising sources of natural antioxidants. Results of antiplasmodial activity demonstrated that *B. enneandra* could be a potential source of antimalarial agent(s). Cytotoxicity screening suggested that *D. hirtiflora* and *M. inermis* are promising candidates with antitumor activity. It is well known that plant extracts comprise mixtures of complex metabolites, which exert their action on different levels and through several mechanisms. Thus, isolation and determination of bioactive molecule(s) responsible for the antitumor activity of these two species and their mode of action are warranted to evaluate their potential for cancer therapy. Moreover, bioassay-guided fractionation of crude extracts of *B. enneandra* would be interesting to determine and verify whether the same constituents or different ones exert their toxicity towards cancer and normal cells. To the best of our knowledge, this is the first report on the cytotoxicity property of these two plants and the antiplasmodial activity of *B. enneandra*.

CONFLICTS OF INTEREST

Given his role as Associate Editor, Gokhan Zengin has not been involved and has no access to information regarding the

peer review of this article. Full responsibility for the editorial process for this article was delegated to Editor-in-Chief Carlos L. Cespedes Acuña. All authors reported that there is no conflict of interest associated to this work.

ACKNOWLEDGMENTS

The authors are thankful to the Researchers Supporting Project number (RSP-2021/86), King Saud University, Riyadh, Saudi Arabia.

ORCID

Ibrahim B. E. El Bashir	0000-0002-9104-5743
Maha Y. Kordofani	0000-0003-1334-215X
Sakina Yagi	0000-0002-9600-3526
Abdulrahman A. Al-Atar	0000-0002-8884-9820
Ahmed A. Qahtan	0000-0002-0064-1516
Eslam M. Abdel-Salam	0000-0003-3297-9551
Abdullah M. Alkahtani	0000-0002-4780-786X
Majed A. Al-Mansoub	0000-0003-0162-5830
Ammar Mohammed Ahmed Ali	0000-0002-3270-6396
Ali M. Batah	0000-0003-1117-376X
Gokhan Zengin	0000-0001-6548-7823

A. APPENDIX, SUPPLEMENTARY INFORMATION

Supplementary information to this article can be found online at <https://doi.org/10.53365/nrfhh/146223>.

AUTHOR CONTRIBUTIONS

IBEEB, SY, AAAA, AMA - Research concept and design; IBEEB, MYK, AAQ, EMAS - Collection and/or assembly of data; MYK, AAAA, AAQ, EMAS - Data analysis and interpretation, SY, AAQ, AMA, MAAM, AMAA - Writing the article; SY, AMA, MAAM, AMAA, GZ - Critical revision of the article, GZ - Final approval of the article.

REFERENCES

- Adeniran, A., Sonibare, M., Ajayi, O., Rajacharya, G., Kumar, S., 2020. Antioxidant Activity of column fractions and Caryatin isolated from the ethyl acetate extract of *Dioscorea hirtiflora* tuber. *Tropical Journal of Natural Product Research*. 4, 276–281. <https://doi.org/10.26538/tjnpr/v4i7.4>
- Alamin, M.A., Yagi, A.I., Yagi, S.M., 2015. Evaluation of antidiabetic activity of plants used in Western Sudan. *Asian Pacific Journal of Tropical Biomedicine*. 5, 395–402. [https://doi.org/10.1016/S2221-1691\(15\)30375-0](https://doi.org/10.1016/S2221-1691(15)30375-0)
- Al-Muniri, R.M.S., Hossain, M.A., 2017. Evaluation of antioxidant and cytotoxic activities of different extracts of folk medicinal plant *Haplophyllum tuberculatum*. *Egyptian Journal of Basic and Applied Sciences*. 4, 101–106. <https://doi.org/10.1016/j.ejbas.2017.04.003>
- Bah, S., Jäger, A.K., Adersen, A., Diallo, D., Paulsen, B.S., 2007. Antiplasmodial and GABAA-benzodiazepine receptor binding activities of five plants used in traditional medicine in Mali, West Africa. *Journal of Ethnopharmacology*. 110, 451–457. <https://doi.org/10.1016/j.jep.2006.10.019>
- Benzie, I.F.F., Strain, J.J., 1996. The Ferric Reducing Ability of Plasma (FRAP) as a Measure of “Antioxidant Power”: The FRAP Assay. *Analytical Biochemistry*. 239, 70–76. <https://doi.org/10.1006/abio.1996.0292>
- Bray, F., Ren, J.S., Masuyer, E., Ferlay, J., 2013. Global estimates of cancer prevalence for 27 sites in the adult population in 2008. *International Journal of Cancer*. 132, 1133–1145. <https://doi.org/10.1002/ijc.27711>
- Deshpande, S., Patil, M., Parmar, K., Daswadkar, S., Khodade, R., 2011. A study of antioxidant activity of fruit extract of *Coccinia grandis* (L., Voigt). *International Journal of Drug Research and Technology*. 1, 69–72.
- Doka, I., Yagi, S., 2009. Ethnobotanical survey of medicinal plants in West Kordofan (Western Sudan).
- Doka, I.G., 2012. Evaluation of some wild Plant from south-west Kordofan As food sources, and others, (Eds.). LAP LAMBERT Academic Publishing, pp. 1–132.
- Elamin, A., Ibrahim, M.E., Abuidris, D., Mohamed, K.E.H., Mohammed, S.I., 2015. Part I: cancer in Sudan-burden, distribution, and trends breast, gynecological, and prostate cancers. *Cancer medicine*. 4, 447–456. <https://doi.org/10.1002/cam4.378>
- Elebead, F.M., Hamid, A., Hilmi, H.S.M., Galal, H., 2012. Mapping Cancer Disease Using Geographical Information System (GIS) in Gezira State-Sudan. *Journal of Community Health*. 37, 830–839. <https://doi.org/10.1007/s10900-011-9517-9>
- Ghazali, G.E., Tohami, M.E., Egami, A.E., Abdalla, W., Mohammed, M., 1997. Medicinal plants of the Sudan. Part IV. Medicinal plants of northern Kordofan. .
- Hassan, L.E., A., Ahamed, K., Majid, M.B., Iqbal, A.S., Suede, M.A., Haque, F.S.R., Ismail, R.A., Ein, Z., Majid, O.C., A, A.M.S., 2014. Crystal structure elucidation and anticancer studies of (-)-pseudosemiglabin: a flavanone isolated from the aerial parts of *Tephrosia apollinea*. *PLoS One*. 9, 90806. <https://doi.org/10.1371/journal.pone.0090806>
- Issa, T.O., Mohamed, Y.S., Yagi, S., Ahmed, R.H., Najeeb, T.M., Makhawi, A.M., Khider, T.O., 2018. Ethnobotanical investigation on medicinal plants in Algoz area (South Kordofan), Sudan. *Journal of Ethnobiology and Ethnomedicine*. 14, 31. <https://doi.org/10.1186/s13002-018-0230-y>
- Jimoh, F.O., Adedapo, A.A., Afolayan, A.J., 2010. Comparison of the nutritional value and biological activities of the acetone, methanol and water extracts of the leaves of *Solanum nigrum* and *Leonotis leonorus*. *Food and Chemical Toxicology*. 48, 964–971. <https://doi.org/10.1016/j.fct.2010.01.007>
- Karunakaran, T., Firouz, N.S., Santhanam, R., Jong, V.Y.M., 2020. Phytochemicals from *Calophyllum macrocarpum* Hook.f. and its cytotoxic activities. *Natural Product Research*. 36(2), 654–659. <https://doi.org/10.1080/14786419.2020.1795658>
- Khalid, H., Abdalla, W.E., Abdelgadir, H., Opatz, T., Efferth, T., 2012. Gems from traditional north-African medicine: medicinal and aromatic plants from Sudan. *Natural Products and Bioprospecting*. 2, 92–103. <https://doi.org/10.1007/s13659-012-0015-2>
- Khalid, S.A., Farouk, A., Geary, T.G., Jensen, J.B., 1986. Potential antimalarial candidates from African plants: An in vitro approach using *Plasmodium falciparum*. *Journal of Ethnopharmacology*. 15, 201–209. [https://doi.org/10.1016/0378-8741\(86\)90156-X](https://doi.org/10.1016/0378-8741(86)90156-X)
- Kiendrebeogo, M., Dijoux-Franca, M.G., Lamien, C.E., Meda, A., Wouessidjewe, D., 2005. Acute toxicity and antioxidant property of *Striga hermonthica* (Del.) Benth (Scrophulariaceae). *African Journal*

- of Biotechnology. 4(9), 919–922.
- Kumar, J.U., Chaitanya, S., J, M., Semotiuk, K., J, A., V, K., 2019. Indigenous knowledge on medicinal plants used by ethnic communities of South India. *Ethnobotany Research and Applications*. 18, 1–112. <https://doi.org/10.32859/era.18.4.1-112>
- Liu, Y., Li, H., Fan, Y., Man, S., Liu, Z., Gao, W., Wang, T., 2016. Antioxidant and antitumor activities of the extracts from Chinese yam (*Dioscorea opposita* Thunb.) flesh and peel and the effective compounds. *Journal of Food Science*. 81, 1553–1564. <https://doi.org/10.1111/1750-3841.13322>
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*. 65, 55–63. [https://doi.org/10.1016/0022-1759\(83\)90303-4](https://doi.org/10.1016/0022-1759(83)90303-4)
- Murugan, M., Mohan, V.R., 2012. In vitro antioxidant studies of *Dioscorea esculenta* (Lour). Burkill. *Asian Pacific Journal of Tropical Biomedicine*. 2, 1620–1624. [https://doi.org/10.1016/S2221-1691\(12\)60464-X](https://doi.org/10.1016/S2221-1691(12)60464-X)
- OCHA., 2019. SUDAN Situation Report. <https://reports.unocha.org/en/country/sudan/> Accessed on 15th February 2021.
- Okpako, L.C., Ajaiyeoba, E.O., 2004. In vitro and in vivo antimalarial studies of *Striga hermonthica* and *Tapinanthus sessilifolius* extracts. *African Journal of Medicine and Medical Sciences*. 33, 73–75.
- Orabueze, C.I., Adesegun, S.A., Ejeatuluchukwu, O., Ota, D.A., Coker, H.A., 2019. In vivo antimalarial and in vitro antioxidant activities of hydro-methanol leaf extract of *Fadogia cienkowskii* Schweinf.(Rubiaceae). *African Journal of Pharmacology and Therapeutics*. 8(1), 6–13.
- Ordoñez, A.A.L., Gomez, J.D., Vattuone, M.A., Lsla, M.I., 2006. Antioxidant activities of *Sechium edule* (Jacq.) Swartz extracts. *Food Chemistry*. 97, 452–458. <https://doi.org/10.1016/j.foodchem.2005.05.024>
- Ouédraogo, R.J., Somda, M.B., Ouattara, L., Kagambega, W., Ouoba, P., Ouédraogo, G.A., 2020. Evaluation of the Antioxidant and α -amylase Inhibitory Activities of *Mitragyna inermis* (Willd) O. Kuntze. *Journal of Experimental Biology and Agricultural Sciences*. 8(5), 676–682. [https://doi.org/10.18006/2020.8\(5\).676.682](https://doi.org/10.18006/2020.8(5).676.682)
- Philip, K., Elizabeth, M.M., Cheplogoi, P.K., Samuel, K.T., 2017. Ethnobotanical survey of antimalarial medicinal plants used in Butebo County, Eastern Uganda. *European Journal of Medicinal Plants*. 21, 1–22.
- Prabhavathi, R., Prasad, M., Jayaramu, M., 2016. In-vitro antioxidant studies of *Cissus quadrangularis* (L) extracts. *European Journal of Experimental Biology*. 6, 1–6.
- Ramana, K.V., Reddy, A.B.M., Majeti, N.V.R.K., Singhal, S.S., 2018. Therapeutic Potential of Natural Antioxidants. *Oxidative Medicine and Cellular Longevity*. 2018, 9471051. <https://doi.org/10.1155/2018/9471051>
- Ravikumar, S., Inbaneson, S.J., Suganthi, P., 2012. In vitro antiplasmodial activity of ethanolic extracts of South Indian medicinal plants against *Plasmodium falciparum*. *Asian Pacific Journal of Tropical Disease*. 2, 180–183. [https://doi.org/10.1016/S2222-1808\(12\)60043-7](https://doi.org/10.1016/S2222-1808(12)60043-7)
- Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice-Evans, C., 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*. 26, 1231–1237. [https://doi.org/10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3)
- Sonibare, M.A., Abegunde, R.B., 2012. In vitro antimicrobial and antioxidant analysis of *Dioscorea dumetorum* (Kunth) Pax and *Dioscorea hirtiflora* (Linn.) and their bioactive metabolites from Nigeria. *Journal of Applied Biosciences*. 51, 3583–3590.
- Spanou, C., Stagos, D., Aligiannis, N., Kouretas, D., 2010. Influence of Potent Antioxidant Leguminosae Family Plant Extracts on Growth and Antioxidant Defense System of Hep2 Cancer Cell Line. *Journal of Medicinal Food*. 13, 149–155. <https://doi.org/10.1089/jmf.2009.0058>
- Tefera, B.N., Kim, Y.D., 2019. Ethnobotanical study of medicinal plants used as antimalarial and repellent by Sidama people of Hawassa Zuria district, Southern Ethiopia. *Complementary Medicine Research*. 10, 13–26. <https://doi.org/10.5455/jcmr.20181102063241>
- Tiwari, M., Gupta, P.S., Sharma, N., 2021. A Preliminary Study on In Vitro Antioxidant and In Vivo Anti-Inflammatory Activity of *Cissus quadrangularis* Linn. *Research Journal of Pharmacy and Technology*. 14, 2619–2624. <https://doi.org/10.52711/0974-360X.2021.00461>
- Traore, F., Gasquet, M., Laget, M., Guiraud, H., Giorgio, C.D., Azas, N., Doumbo, O., Timon-David, P., 2000. Toxicity and genotoxicity of antimalarial alkaloid rich extracts derived from *Mitragyna inermis* O. Kuntze and *Nauclea latifolia*. *Phytotherapy Research*. 14, 608–611. [https://doi.org/10.1002/1099-1573\(200012\)14:8<608::AID-PTR667>3.0.CO;2-D](https://doi.org/10.1002/1099-1573(200012)14:8<608::AID-PTR667>3.0.CO;2-D)
- Traoré, M.S., Baldé, M., Camara, A., Baldé, E.S., Diané, S., Diallo, M.S.T., Keita, A., Cos, P., Maes, L., Pieters, L., Baldé, A.M., 2015. The malaria co-infection challenge: An investigation into the antimicrobial activity of selected Guinean medicinal plants. *Journal of Ethnopharmacology*. 174, 576–581. <https://doi.org/10.1016/j.jep.2015.03.008>
- Wang, J.M., Ji, L.L., Branford-White, C.J., Wang, Z.Y., Shen, K.K., Liu, H., Wang, Z.T., 2012. Antitumor activity of *Dioscorea bulbifera* L. rhizome in vivo. *Fitoterapia*. 83, 388–394. <https://doi.org/10.1016/j.fitote.2011.12.001>
- Wang, K.J., Yang, C.R., Zhang, Y.J., 2007. Phenolic antioxidants from Chinese toon (fresh young leaves and shoots of *Toona sinensis*). *Food Chemistry*. 101, 365–371. <https://doi.org/10.1016/j.foodchem.2006.01.044>
- WHO., 2020. Malaria Fact Sheets. <https://www.who.int/news-room/fact-sheets/detail/malaria>. Date accessed: 2021-02-15
- Wilson, D.W., Nash, P., Buttar, H.S., Griffiths, K., Singh, R., Meester, F.D., Horiuchi, R., Takahashi, T., 2017. The Role of Food Antioxidants, Benefits of Functional Foods, and Influence of Feeding Habits on the Health of the Older Person: An Overview. *Antioxidants*. 6, 81. <https://doi.org/10.3390/antiox6040081>
- Witschel, M., Rottmann, M., Kaiser, M., Brun, R., 2012. Agrochemicals against malaria, sleeping sickness, leishmaniasis and Chagas disease. *PLoS Neglected Tropical Diseases*. 6, e1805. <https://doi.org/10.1371/journal.pntd.0001805>
- Wolfe, K., Wu, X., Liu, R.H., 2003. Antioxidant activity of apple peels. *Journal of Agricultural and Food Chemistry*. 51, 609–614. <https://doi.org/10.1021/jf020782a>
- Zhang, Z., Wang, X., Liu, C., Li, J., 2016. The degradation, antioxidant and antimutagenic activity of the mucilage polysaccharide from *Dioscorea opposita*. *Carbohydrate Polymers*. 150, 227–231. <https://doi.org/10.1016/j.carbpol.2016.05.034>
- Zheleva-Dimitrova, D., Gevrenova, R., Zaharieva, M.M., Najdenski, H., Ruseva, S., Lozanov, V., Balabanova, V., Yagi, S., Momekov, G., Mitev, V., 2017. HPLC-UV and LC-MS Analyses of Acylquinic Acids in *Geigeria alata* (DC) Oliv. & Hiern. and their Contribution to Antioxidant and Antimicrobial Capacity. *Phytochemical Analysis*. 28, 176–184. <https://doi.org/10.1002/pca.2658>