

Original Research

View Article Online



Received 02 September 2022
Revised 11 September 2022
Accepted 15 September 2022
Available online 21 September 2022

Edited by Shaif Ullah Khan

KEYWORDS:

Benzo[c]phenanthridine alkaloids
Chemotaxonomy
Cytotoxicity
Flavonoids
Rutaceae
Zanthoxylum gillettii

Natr Resour Human Health 2022; 1-5
<https://doi.org/10.53365/nrfhh/154598>
eISSN: 2583-1194
Copyright © 2022 Visagaa Publishing House

Benzo[c]phenanthridine alkaloids and glycosylated flavonoids from the roots of *Zanthoxylum gillettii* (Rutaceae)

Liliane Clotilde Dzouemo¹, Emmanuel Ngeufa Happi^{1,*}, Mouthe Gervais Happi², Willifred Dongmo Tekapi Tsopgni¹, Jean Duplex Wansi¹

¹Department of Chemistry, Faculty of Sciences, University of Douala, P.O. Box 24157, Douala, Cameroon

²Department of Chemistry, Higher Teacher Training College Bambili, The University of Bamenda, P.O. Box 39, Bambili, Cameroon

ABSTRACT: The roots of *Zanthoxylum gillettii* have been chemically investigated and led to the isolation and characterization of nine compounds including three benzo[c]phenanthridine alkaloids oxychelerythrine (1), angoline (2) and arnottianamide (3); two glycosylated flavonoids hesperidin (4) and its analogue neohesperidin (5) along with four common specialized metabolites including sesamin (6), lupeol (7), β -sitosterol (8) and its derivative daucosterol (9). Their structures were established based on their recorded 1D and 2D-NMR data. Some compounds (1, 2, 3, 6 and 7) did not display a cytotoxic activity until the highest concentration of 200 μ M against the human cell lines MCF-7 and A549. The chemotaxonomic relevance of this study has been discussed. To the best of our knowledge, except for the five compounds 3, 6-9, all the four others have been for the first time isolated from *Z. gillettii* (Rutaceae).

1. INTRODUCTION

Rutaceae represents a large plant family which along with Meliaceae and Simaroubaceae families are called Sapindales which are distinguishable by their ability to produce limonoids (Happi et al., 2018; Kemayou et al., 2021). Rutaceae consists to a total of 148 plant genera and 1,487 accepted species including *Zanthoxylum gillettii* (De Wild.) P.G. Waterman (Rutaceae) also known as *Fagara tessmanii* or *Fagara macrophylla* (Gaya et al., 2013; List, 2022). Recent review on the genus *Zanthoxylum* revealed that its plant species display a large variety of bioactive specialized metabolites belonging to various classes of compounds including lignans, alkaloids, flavonoids, coumarins and triterpenoids with a large scale of activities related to cancer, infectious diseases and sickle cell anaemia (Okagu et al., 2021). The plant *Z. gillettii* is widely used in Cameroon as a spice (due to its relevant fragrance and taste) and in African traditional medicine for the cure of numerous diseases viz cancer, malaria, inflammation or microbial infections (Zirihi et al., 2009). As part of our search for bioactive natural products with good cytotoxicity or less toxicity (Happi et al., 2021; Makong et al., 2019; Mbougna et al., 2021), we have investigated the roots of *Z. gillettii* collected in the West region of Cameroon. The chemical investigations of the methanolic extract led to the isolation of nine compounds

which were screened for their cytotoxic potency against the two cell lines MCF-7 (human breast adenocarcinoma cell line) and A549 (human lung cancer cell line). The chemotaxonomy relevance of the isolated compounds in this study is also discussed in the paper.

2. MATERIALS AND METHODS

2.1. General instrumentation

Details can be retrieved from the supplementary information associated with this paper (Appendix A).

2.2. Plant material

The roots of *Zanthoxylum gillettii* (De Wild.) P.G. Waterman were harvested at Bana-Tentcheu (GPS coordinates: Latitude 5°07'57"N, Longitude 10°17'41"E, Elevation: 1412 m), West region, Cameroon, in December 2021. Some leaves and bark samples have been used for the identification of the plant has been done by Mr Victor Nana in comparison to some collected leaves with the plant specimen registered under the voucher number 38960 HNC in the National Herbarium of Cameroon database.

* Corresponding author.

E-mail address: ngeufa@yahoo.fr (Emmanuel Ngeufa Happi)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2.3. Extraction and isolation

The air-dried and powdered roots of *Zanthoxylum gillettii* (~3 kg) were macerated twice in methanol at room temperature for 48 h, each, to give a dark brown gum as crude extract (63.2 g) after removing of solvent. Small amount (~3.8 g) of the crude extract was kept for further investigations (biological tests and chemical analyses) and the remaining extract was suspended in water (100 ml) and partitioned with *n*-hexane, dichloromethane and ethyl acetate (500 ml, each), as well as *n*-butanol (200 ml), to afford four main fractions A (12.7 g), B (22.3 g), C (10.4 g) and D (4.7 g), respectively.

Fraction A was mainly a yellowish oil and fats. Its purification by silica gel column chromatography (230–400 mesh, 5.0 x 75.0 cm) using an increasing polarity of ethyl acetate (EtOAc) (0 to 40%) in *n*-hexane (*n*-Hex). The process led to the isolation of two common compounds namely lupeol (7) (5.8 mg) and β -sitosterol (8) (13.6 mg) at the polarities *n*-Hex/AcOEt 19:1 and *n*-Hex/AcOEt 9:1, respectively.

However, dichloromethane-soluble fraction B was further purified using silica gel column chromatography eluting with increasing polarity of ethyl acetate (EtOAc) (10 to 70%) in *n*-hexane (*n*-Hex). A total of 221 sub-fractions (ca. 100 ml each) were collected and combined into 7 series B1–B7 based on their TLC profiles. Arnottianamide (3, 4.6 mg), sesamin (6, 12.7 mg) and angoline (2, 3.2 mg) were obtained as powders from series B3 (*n*-Hex/AcOEt 3:1), B4 (*n*-Hex/AcOEt 7:3) and B5 (*n*-Hex/AcOEt 3:2), respectively.

Furthermore, the series B7 (2.4 g, *n*-Hex/AcOEt 1:1), the ethyl acetate soluble fraction C and the *n*-butanol soluble fraction D displayed similar TLC profiles and were combined, then further chromatographed over silica gel column eluting with increasing polarity of methanol (MeOH) (0 to 30%) in dichloromethane (DCM). Following the collection of a total of 237 sub-fractions (ca. 100 ml each), combined into six series D1 – D6, oxychelerythrine (1, 6.2 mg) was recrystallised from series D3 (DCM/MeOH 37:3), hesperidin (4) and its analogue neohesperidin (5) (8.2 mg) were obtained as an unseparated mixture of yellow powders from series D4 (DCM/MeOH 4:1) with the relative proportion 1:1 in the mixture based on the heights of ^{13}C NMR peaks of corresponding carbon atoms in the two compounds as displayed on Figures 1S and 2S (see Appendix A), while the common steroid daucosterol (9, 12.2 mg) was precipitated as a white powder from D2 (DCM/MeOH 9:1).

2.4. Physical aspects and spectral data of isolated compounds

Further details on the spectral data of isolated compounds are available in the supplementary information (Appendix A).

2.5. Cytotoxicity activity

The detailed protocol for the biological test is given in the supplementary information associated with this paper (Appendix A).

3. RESULTS AND DISCUSSION

3.1. Phytochemical study

The chemical study of the roots of *Zanthoxylum gillettii* led to the isolation of nine compounds (Figure 1 for compounds 1–5 and Figure 2 for compounds 6–9). The structures of the isolated compounds have been established using their spectroscopic data, as well as the comparison of these data with those reported in the literature. The nine isolated compounds have been identified as oxychelerythrine (1) (Fuchino et al., 2010), angoline (2) (Lee et al., 1998), arnottianamide (3) (Yang et al., 2009), hesperidin (4) (Tine et al., 2020), neohesperidin (5) (D.I. Hamdan et al., 2014), sesamin (6) (Yang et al., 2009), lupeol (7) (Jouwa et al., 2020; Tsopgni et al., 2019; Wouamba, Happi, Lenta, et al., 2020; Wouamba, Happi, Poufofo, et al., 2020), β -sitosterol (8) (Keugwa et al., 2021) and its derivative daucosterol (9) (Happi et al., 2020).




Figure 1. Benzo[*c*]phenanthridine alkaloids and flavonoids from the roots of *Zanthoxylum gillettii*




Figure 2. Structures of sesamin (6), lupeol (7), β -sitosterol (8) and daucosterol (9) isolated from the roots of *Zanthoxylum gillettii*.

3.2. Cytotoxicity of some isolated compounds

Five compounds including the three benzo[*c*]phenanthridine alkaloids (1–3) and the two major compounds sesamin (6) and lupeol (7) were evaluated for their cytotoxicity against the human breast adenocarcinoma cell line (MCF-7) and the human lung cancer cell line (A549) using doxorubicin as reference. The recorded results showed that all the tested compounds did not display an activity until the highest concentration of 200 μM ($\text{IC}_{50} > 200 \mu\text{M}$).

3.3. Chemotaxonomic significance of the study

The chemical investigations of the methanol extract of *Z. gilletii* roots resulted in the identification of nine compounds namely oxchelerythrine (1), angoline (2), arnottianamide (3), the mixture hesperidin (4) and neohesperidin (5), as well as the four common compounds sesamin (6), lupeol (7), β -sitosterol (8) and daucosterol (9).

The literature survey on the previous chemical studies of the genus *Zanthoxylum* indicated that, to the best of our knowledge, it is the first report of compounds 1, 2, 4 and 5 from the plant species *Z. gilletii* but have been already isolated from other species in the genus and the family Rutaceae.

Benzo[c]phenanthridine alkaloids are well-known to be found in the family Rutaceae (Laines-Hidalgo et al., 2022). Therefore, the isolation of the three benzo[c]phenanthridine alkaloids 1-3 further supported the classification of the plant as belonging to the family Rutaceae.

Moreover, the literature indicated that oxchelerythrine (1) has been reported from *Z. nitidum*, *Z. schinifolium* and *Z. capense* (Cabral et al., 2015); angoline (2) from *Z. nitidum* or *Z. thomense* (J. Liu et al., 2014; Simeray et al., 1985), while arnottianamide (3) was obtained from roots of *Z. gilletii*, *Z. nitidum* and *Z. leprieurii* (Adesina & Reisch, 1988; Zeng et al., 2022). This may suggest that benzo[c]phenanthridine alkaloids might be considered as markers compounds for the genus *Zanthoxylum*.

Further chemotaxonomic insights of the study have been deduced from the presence of the other compounds isolated during the current investigation of the roots of *Z. gilletii*. Unsurprisingly, the lignan sesamin has been isolated during our investigation as one of the major compounds and it has been mainly reported from several species of the genus *Zanthoxylum* including *Z. chalybeum*, *Z. paracanthum*, *Z. leprieurii*, *Z. rigidum*, *Z. paracanthum*, *Z. zanthoxyloides*, *Z. nitidum* and *Z. heitzii* (Andima et al., 2020; Chakthong et al., 2019; Kaigongi et al., 2020; Muema et al., 2021; Oloya et al., 2021; Omosa et al., 2022; Santos et al., 2020; Wangenstein et al., 2016).

Additionally, the compounds identified as lupeol, β -sitosterol, daucosterol are mainly distributed in the plant kingdom and within the family Rutaceae, they have been isolated from *Z. acanthopodium*, *Z. schinifolium*, *Z. tessmannii* or *Citrus aurantium* (Bissim et al., 2019; Cheng et al., 2002; Kenmoe et al., 2019). Finally, flavonoids are usually obtained from several plants and more particularly from the genus *Zanthoxylum*. Hesperidin (4) was previously isolated from *Z. scandens*, *Z. leprieurii*, *Z. zanthoxyloides* and *Z. myriacanthum* var. *pubescens* (Ngoumfo et al., 2010; Nguyen et al., 2002; Queiroz et al., 2006; Zhang et al., 2017), while neohesperidin (5) was reported from *Z. dissitum* (S. Liu et al., 2009) but more from some Rutaceae plants including *Murraya tetramera* and *Citrus pyriformis* (D. Hamdan et al., 2011; Zhou et al., 2014).

Taken together, all the isolated compounds from the roots of *Z. gilletii* provide significant insights into the taxonomy of the plant and extend the knowledge of its chemistry.

4. CONCLUDING REMARKS

The present paper deals with the chemical study of the methanol soluble extract of *Z. gilletii* roots and the cytotoxic evaluation of the isolated compounds against the tumour cell lines MCF-7 and A549. Nine compounds have been identified using their spectral data as three alkaloids 1-3, two glycosylated flavonoids 4 and 5, as well as four other common compounds including sesamin (6), lupeol (7), β -sitosterol (8) and daucosterol (9). The results from biological tests showed that compounds 1-3, 6 and 7 did not display any cytotoxicity against the two selected cell lines until the concentration of 200 μ M. Furthermore, the chemotaxonomic study of the isolated compounds further supports the classification of the plants by the presence of benzo[c]phenanthridine alkaloids majorly reported from Rutaceae, Fumariaceae and Papaveraceae; sesamin (6) abundantly distributed in the genus *Zanthoxylum* as well as lupeol (7) which is largely found in the family Rutaceae. The plant deserves further study for its minor compounds and another batch of biological activities with different cell lines in order to identify the most potent metabolites that can be classified as lead compounds for the development of new drugs for the local population. However, this study furnishes four additional compounds which enrich the chemistry of the plant *Z. gilletii*.

CONFLICTS OF INTEREST

The authors disclose that they have no competing interests.

ORCID

Liliane Clotilde Dzuemo	0000-0003-4967-4316
Emmanuel Ngeufa Happi	0000-0002-2276-3709
Mouthe Gervais Happi	0000-0001-9659-6125
Willifred Dongmo Tekapi Tsopgni	0000-0002-1644-8171
Jean Duplex Wansi	0000-0002-5111-4361

A. APPENDIX. SUPPLEMENTARY INFORMATION

The supplementary information associated with this article available at <https://doi.org/10.53365/nrfhh/154598>.

AUTHOR CONTRIBUTIONS

Research concept and design : LCD, GMH; Collection of data : WDTT; Data analysis and interpretation: LCD; Writing the article : GMH, LCD; Supervision, critical revision and final approval of the article : ENH, JDW.

REFERENCES

- Adesina, S.K., Reisch, J., 1988. Natural product chemistry. Part 119. Arnottianamide and other constituents of *Zanthoxylum gilletii* root. Journal of Natural Products. 51, 601-602. <https://doi.org/10.1021/np50057a031>
- Andima, M., Coghi, P., Yang, L.J., Wong, V.K.W., Ngule, C.M., Heydenreich, M., Ndakala, A.J., Yenesew, A., Derese, S., 2020.

- Antiproliferative activity of secondary metabolites from *Zanthoxylum zanthoxyloides* Lam: *in-vitro* and *in-silico* studies. *Pharmacognosy Communications*. 10, 44–51. <http://dx.doi.org/10.5530/pc.2020.1.8>
- Bissim, S., Kenmogne, S.B., Tcho, A.T., Lateef, M., Ahmed, A., Nguefa, H.E., Wansi, J.D., Ali, M.S., Kamdem, W.A.F., 2019. Bioactive acridone alkaloids and their derivatives from *Citrus aurantium* (Rutaceae). *Phytochemistry Letters*. 29, 148–153. <https://doi.org/10.1016/j.phytol.2018.12.010>
- Cabral, V., Luo, X., Junqueira, E., Costa, S.S., Mulhovo, S., Duarte, A., Couto, I., Viveiros, M., Ferreira, M.-J.U., 2015. Enhancing activity of antibiotics against *Staphylococcus aureus*: *Zanthoxylum capense* constituents and derivatives. *Phytomedicine*. 22, 469–476. <https://doi.org/10.1016/j.phymed.2015.02.003>
- Chakthong, S., Ampaprom, R., Inparn, S., Phetkul, U., Chusri, S., Limsuwan, S., Voravuthikunchai, S.P., 2019. New alkylamide from the stems of *Zanthoxylum nitidum*. *Natural Product Research*. 33, 153–161. <https://doi.org/10.1080/14786419.2018.1440218>
- Cheng, M.J., Yang, C.H., Lin, W.Y., Lin, W.Y., Tsai, I.L., Chen, I.S., 2002. Chemical constituents from the leaves of *Zanthoxylum schinifolium*. *Journal of the Chinese Chemical Society*. 49, 125–128. <https://doi.org/10.1002/jccs.200200021>
- Fuchino, H., Kawano, M., Yasumoto, K.M., Sekita, S., Satake, M., Ishikawa, T., Kiuchi, F., Kawahara, N., 2010. In vitro leishmanicidal activity of benzophenanthridine alkaloids from *Bocconia pearcei* and related compounds. *Chemical and Pharmaceutical Bulletin*. 58, 1047–1050. <https://doi.org/10.1248/cpb.58.1047>
- Gaya, C.H., Kawaka, J.F., Muchugi, A., Ngeranwa, J.J., 2013. Variation of alkaloids in the Kenyan *Zanthoxylum gillettii* (De wild Waterman). *African Journal of Plant Science*. 7, 438–444. <https://doi.org/10.5897/AJPS2013.1008>
- Hamdan, D., El-Readi, M.Z., Tahrani, A., Herrmann, F., Kaufmann, D., Farrag, N., El-Shazly, A., Wink, M., 2011. Secondary metabolites of ponderosa lemon (*Citrus pyriformis*) and their antioxidant, anti-inflammatory, and cytotoxic activities. *Zeitschrift für Naturforschung C*. 66, 385–393. <https://doi.org/10.1515/znc-2011-7-810>
- Hamdan, D.I., Mahmoud, M.F., Wink, M., El-Shazly, A.M., 2014. Effect of hesperidin and neohesperidin from bitter orange (*Citrus aurantium* var. *bigaradia*) peel on indomethacin-induced peptic ulcers in rats. *Environmental Toxicology and Pharmacology*. 37, 907–915. <https://doi.org/10.1016/j.etap.2014.03.006>
- Happi, G.M., Kemayou, G.P.M., Stammer, H.G., Neumann, B., Ismail, M., Kouam, S.F., Wansi, J.D., Tchouankeu, J.C., Frese, M., Lenta, B.N., Sewald, N., 2021. Three phragmalin-type limonoids orthoesters and the structure of odorone isolated from the bark of *Entandrophragma candollei* (Meliaceae). *Phytochemistry*. 181, 112537. <https://doi.org/10.1016/j.phytochem.2020.112537>
- Happi, G.M., Ngadjui, B.T., Green, I.R., Kouam, S.F., 2018. Phytochemistry and pharmacology of the genus *Entandrophragma* over the 50 years from 1967 to 2018: a ‘golden’ overview. *Journal of Pharmacy and Pharmacology*. 70, 1431–1460. <https://doi.org/10.1111/jphp.13005>
- Happi, G.M., Wouamba, S.C.N., Ismail, M., Kouam, S.F., Frese, M., Lenta, B.N., Sewald, N., 2020. Ergostane-type steroids from the Cameroonian ‘white tiama’ *Entandrophragma angolense*. *Steroids*. 156, 108584. <https://doi.org/10.1016/j.steroids.2020.108584>
- Jouwa, N.S.T., Mvot-Akak, C., Happi, G.M., Frese, M., Stammer, H.G., Neumann, B., Lenta, B.N., Sewald, N., Nkengfack, A.E., 2020. Antioxidant norbergenin derivatives from the leaves of *Diospyros gillettii* De Wild (Ebenaceae). *Phytochemistry Letters*. 36, 63–67. <https://doi.org/10.1016/j.phytol.2020.01.012>
- Kaigongi, M.M., Lukhoba, C.W., Yaouba, S., Makunga, N.P., Githiomi, J., Yenesew, A., 2020. In vitro antimicrobial and antiproliferative activities of the root bark extract and isolated chemical constituents of *Zanthoxylum paracanthum* Kokwaro (Rutaceae). *Plants*. 9, 920. <https://doi.org/10.3390/plants9070920>
- Kemayou, G.P.M., Kache, S.F., Dzuemo, L.C., Happi, G.M., Kouam, S.F., Tchouankeu, J.C., 2021. Phytochemistry, traditional uses, and pharmacology of the genus *Ekebergia* (Meliaceae): A review. *Trends in Phytochemical Research*. 5(3), 110–125. <https://dx.doi.org/10.30495/tpr.2021.1928329.1204>
- Kenmoe, D.C., Kouam, K.A., Azebaze, G.B.A., Belle, E.K.P., Vardamides, J.C., Sewald, N., Wansi, J.D., 2019. A new aromatic amide from the roots of *Zanthoxylum tessmannii* (Rutaceae). *Chemistry & Biodiversity*. 16, 1800590. <https://doi.org/10.1002/cbdv.201800590>
- Keugwa, C.V.T., Ehawa, F.R.E., Happi, G.M., Tsopgni, W.D.T., Waffo, A.F.K., Ndom, J.C., Wansi, J.D., 2021. Three new 30-norfriedelane and a new friedelane triterpenes from the trunk bark of *Caloncoba welwitschii* (Oliv.) Gilg (Achariaceae). *Natural Product Research*, 1–10. <https://doi.org/10.1080/14786419.2021.1981314>
- Laines-Hidalgo, J.I., Muñoz-Sánchez, J.A., Loza-Müller, L., Vázquez-Flota, F., 2022. An update of the sanguinarine and benzophenanthridine alkaloids’ biosynthesis and their applications. *Molecules*. 27, 1378. <https://doi.org/10.3390/molecules27041378>
- Lee, S.K., Qing, W.G., Mar, W., Luyengi, L., Mehta, R.G., Kawanishii, K., Fong, H.H.S., Beecher, C.W.W., Kinghorn, A.D., Pezzuto, J.M., 1998. Angoline and chelerythrine, benzophenanthridine alkaloids that do not inhibit protein kinase C. *Journal of Biological Chemistry*. 273, 19829–19833. <https://doi.org/10.1074/jbc.273.31.19829>
- List, T.P., 2022. Rutaceae. <http://www.theplantlist.org/browse/A/Rutaceae/>. Date accessed: 2022-09-01
- Liu, J., Zhang, Q., Ye, Y., Li, W., Qiu, J., Liu, J., Zhan, R., Chen, W., Yu, Q., 2014. Angoline: A selective IL-6/STAT3 signaling pathway inhibitor isolated from *Zanthoxylum nitidum*. *Phytomedicine*. 21, 1088–1091. <https://doi.org/10.1016/j.phymed.2014.04.001>
- Liu, S., Zhang, W., He, G., Lei, P., Li, X., Liang, Y., 2009. Studies on chemical constituents in stems of *Zanthoxylum dissitum*. *Zhongguo Zhongyao Zazhi*. 34, 571–573.
- Makong, Y.S., Happi, G.M., Djouaka, J.L.B., Wansi, J.D., Nahar, L., Waffo, A.F.K., Martin, C., Sewald, N., Sarker, S.D., 2019. Cytotoxic stilbenes and canthinone alkaloids from *Brucea antidysenterica* (Simaroubaceae). *Molecules*. 24(23), 4412. <https://doi.org/10.3390/molecules24234412>
- Mbougna, J.F.T., Happi, G.M., Bitchagno, G.T.M., Awouafack, M.D., Lenta, B.N., Kouam, S.F., Tane, P., Sewald, N., Tene, M., 2021. Chemical constituents from *Ficus natalensis* Hochst (Moraceae) and their chemophenetic significance. *Biochemical Systematics and Ecology*. 95, 104227. <https://doi.org/10.1016/j.bse.2021.104227>
- Muema, J.M., Bargul, J.L., Mutunga, J.M., Obonyo, M.A., Asudi, G.O., Njeru, S.N., 2021. Neurotoxic *Zanthoxylum chalybeum* root constituents invoke mosquito larval growth retardation through ecdysteroidogenic CYP450s transcriptional perturbations. *Pesticide Biochemistry and Physiology*. 178, 104912. <https://doi.org/10.1016/j.pestbp.2021.104912>
- Ngoumfo, R.M., Jouda, J.B., Mouafo, F.T., Komguem, J., Mbazoa, C.D., Shiao, T.C., Choudhary, M.I., Laatsch, H., Legault, J., Pichette, A., 2010. In vitro cytotoxic activity of isolated acridones alkaloids from *Zanthoxylum leprieurii* Guill. et Perr. *Bioorganic & Medicinal Chemistry*. 18, 3601–3605. <https://doi.org/10.1016/j.bmc.2010.03.040>
- Nguyen, Q.A., Van-Dufat, H.T., Michel, S., Tillequin, F., Dumontet, V., Sevenet, T., 2002. A new phenylpropanoid ester from the bark of *Zanthoxylum scandens* (Rutaceae). *Zeitschrift für Naturforschung C*.

- 57, 986–989. <https://doi.org/10.1515/znc-2002-11-1205>
- Okagu, I.U., Ndefo, J.C., Aham, E.C., Udenigwe, C.C., 2021. *Zanthoxylum* species: a review of traditional uses, phytochemistry and pharmacology in relation to cancer, infectious diseases and sickle cell anemia. *Frontiers in Pharmacology*. 12, 713090. <https://doi.org/10.3389/fphar.2021.713090>
- Oloya, B., Namukobe, J., Heydenreich, M., Ssengooba, W., Schmidt, B., Byamukama, R., 2021. Antimycobacterial activity of the extract and isolated compounds from the stem bark of *Zanthoxylum leprieurii* Guill. and Perr. *Natural Product Communication*. 16, 1–8. <https://doi.org/10.1177/1934578X2111035>
- Omosa, L.K., Nchiozem-Ngitedem, V.A., Mukavi, J., Atieno, O.B., Ombui, N.H., Hashim, I., Obegi, M.J., Efferth, T., Spitteller, M., 2022. Cytotoxic alkaloids from the root of *Zanthoxylum paracanthum* (mildbr) Kokwaro. *Natural Product Research*. 36, 2528–2525. <https://doi.org/10.1080/14786419.2021.1913586>
- Queiroz, E.F., Hay, A.E., Chaaib, F., Van Diemen, D., Diallo, D., Hostettmann, K., 2006. New and bioactive aromatic compounds from *Zanthoxylum zanthoxyloides*. *Planta Medica*. 72, 746–750. <https://doi.org/10.1055/s-2006-941504>
- Santos, C.H.C., Souza, M.C., De Sousa, P.T., Ribeiro, T.A.N., Jacinto, M.J., Santos, R.A.N., Silva, V.C.P., 2020. Lignans from stem bark of *Zanthoxylum rigidum* (Rutaceae) and their leishmanicidal activities. *International Journal of Advanced Biotechnology and Research*. 11, 26–33.
- Simeray, J., Chaumont, J.P., Bevalot, F., Vaquette, J., 1985. Zanthomamide: an aromatic amide from *Zanthoxylum thomense*. *Phytochemistry*. 24, 2720–2721. [https://doi.org/10.1016/S0031-9422\(00\)80704-9](https://doi.org/10.1016/S0031-9422(00)80704-9)
- Tine, Y., Diallo, A., Ndoye, I., Yang, Y., Renucci, F., Guèye, R.S., Baldé, M., Diédhiou, A., Fall, D., Seck, M., Desjobert, J.M., Costa, J., Wélé, A., Paolini, J., 2020. Flavonoid compounds from *Zanthoxylum leprieurii* Guill. et Perr (Rutaceae) extracts and their antioxidant activity against ABTS•+. *Journal of Drug Delivery and Therapeutics*. 10, 120–124. <https://doi.org/10.22270/jddt.v10i2-s.4010>
- Tsopgni, W.D.T., Happi, G.M., Stammler, H.G., Neumann, B., Mbobda, A.S.W., Kouam, S.F., Frese, M., Azebaze, A.G.B., Lenta, B.N., Sewald, N., 2019. Chemical constituents from the bark of the Cameroonian mahogany *Trichilia emetica* Vahl (Meliaceae). *Phytochemistry Letters*. 33, 49–54. <https://doi.org/10.1016/j.phytol.2019.07.009>
- Wangenstein, H., Ho, G.T.T., Tadesse, M., Miles, C.O., Moussavi, N., Mikolo, B., Malterud, K.E., 2016. A new benzophenanthridine alkaloid and other bioactive constituents from the stem bark of *Zanthoxylum heitzii*. *Fitoterapia*. 109, 196–200. <https://doi.org/10.1016/j.fitote.2016.01.012>
- Wouamba, S.C.N., Happi, G.M., Lenta, B.N., Sewald, N., Kouam, S.F., 2020. Vernoguinamide: A new ceramide and other compounds from the root of *Vernonia guineensis* Benth. and their chemophenetic significance. *Biochemical Systematics and Ecology*. 88, 103988. <https://doi.org/10.1016/j.bse.2019.103988>
- Wouamba, S.C.N., Happi, G.M., Poufo, M.N., Tchamgoue, J., Jouda, J.B., Longo, F., Lenta, B.N., Sewald, N., Kouam, S.F., 2020. Antibacterial flavonoids and other compounds from the aerial parts of *Vernonia guineensis* Benth. (Asteraceae). *Chemistry & Biodiversity*. 17, e2000296. <https://doi.org/10.1002/cbdv.202000296>
- Yang, C.H., Cheng, M.J., Lee, S.J., Yang, C.W., Chang, H.S., Chen, I.S., 2009. Secondary metabolites and cytotoxic activities from the stem bark of *Zanthoxylum nitidum*. *Chemistry & Biodiversity*. 6, 846–857. <https://doi.org/10.1002/cbdv.200800107>
- Zeng, Q., Wang, Z.J., Chen, S., Wang, H., Xie, T.Z., Xu, X.J., Xiang, M.L., Chen, Y.C., Luo, X.D., 2022. Phytochemical and anti-MRSA constituents of *Zanthoxylum nitidum*. *Biomedicine & Pharmacotherapy*. 148, 112758. <https://doi.org/10.1016/j.biopha.2022.112758>
- Zhang, H.L., Gan, X.Q., Fan, Q.F., Yang, J.J., Zhang, P., Hu, H.B., Song, Q.S., 2017. Chemical constituents and anti-inflammatory activities of Maqian (*Zanthoxylum myriacanthum* var. *pubescens*) bark extracts. *Scientific Reports*. 7, 45805. <https://doi.org/10.1038/srep45805>
- Zhou, Y., Lv, H., Wang, W., Tu, P., Jiang, Y., 2014. Flavonoids and anthraquinones from *Murraya tetramera* C. C. Huang (Rutaceae). *Biochemical Systematics and Ecology*. 57, 78–80. <https://doi.org/10.1016/j.bse.2014.07.016>
- Zirihi, G., N'guessan, K., Etien, D., Serikouassi, B., 2009. Evaluation *in vitro* of antiplasmodial activity of ethanolic extracts of *Funtumia elastica*, *Rauwolfia vomitoria* and *Zanthoxylum gillettii* on *Plasmodium falciparum* isolates from Côte d'Ivoire. *Journal of Animal and Plant Sciences*. 5, 406–413.