


Original Research

View Article online



Received 15 September 2025  
Revised 04 December 2025  
Accepted 17 December 2025  
Available Online 27 February 2026

Edited by Kannan RR Rengasamy

## KEYWORDS:

*Eleutherine bulbosa*  
Gibberellic acid  
Bulblet formation  
Bulb growth  
Flavonoid  
Polyphenol

<https://doi.org/10.53365/nrfhh/215711>  
eISSN: 2583-1194  
Copyright © 2026 Visagaa Publishing House

## Gibberellic acid enhances growth, bulblet formation, and secondary metabolite accumulation in *Eleutherine bulbosa* (Mill.) Urb

Thi Thanh Hien Tran<sup>1</sup>, Thi My Tu Vang<sup>1</sup>, Kieu Uyen Vi Nguyen<sup>1</sup>, Bach Mai Thi Vo<sup>1</sup>

<sup>1</sup>School of Biology – Biotechnology, University of Science – VNU HCM, Vietnam National University, Ho Chi Minh City, Viet Nam

**ABSTRACT:** *Eleutherine bulbosa* is a medicinally important plant, yet strategies to enhance both yield and bioactive compound content remain limited. This study investigated the role of gibberellic acid (GA<sub>3</sub>) in bulblet induction from mother bulbs and subsequent growth responses in 6-week-old plants. Different GA<sub>3</sub> concentrations were tested independently at each stage, and the most effective treatments were then applied sequentially. GA<sub>3</sub> at 30 ppm applied to mother bulbs significantly promoted bulblet initiation, increasing fresh and dry mass, bulb length, and respiration activity. At the 6-week growth stage, GA<sub>3</sub> at 40 ppm stimulated biomass accumulation and secondary metabolite production, whereas 50 ppm exerted an inhibitory effect. Flavonoid and polyphenol levels increased under GA<sub>3</sub> application, with maximum accumulation observed at 40–50 ppm depending on the developmental phase. Sequential treatments combining 30 ppm at planting and 40 ppm during later growth achieved the greatest improvements in bulb productivity as well as flavonoid and polyphenol enrichment. The findings also highlight potential physiological links among respiration intensity, carbohydrate reserves, and secondary metabolite biosynthesis.

## 1. INTRODUCTION

*Eleutherine bulbosa* (Mill.) Urb is a bulbous herb belonging to the Iridaceae family, rich in secondary compounds such as phenolic and flavonoid derivatives, naphthalene, anthraquinone, and naphthoquinone (Sun et al., 2024). As a traditional remedy, the bulb has been used to treat a number of conditions, such as diabetes, breast cancer, high blood pressure, and sexual issues (Kamarudin et al., 2020; Naspiah et al., 2014). Numerous investigations have confirmed the pharmacological applications of *E. bulbosa*, particularly its antibacterial, antihypertensive, antidiabetic, anti-inflammatory, and antiviral properties (Da Silva et al., 2024; Harlita et al., 2018; Herman et al., 2024; Jiang et al., 2020; Panyachariwat et al., 2024). Given its medicinal and other values, studies aimed

at enhancing bulb production efficiency and increasing the accumulation of pharmaceutically important secondary metabolites are essential.

*Eleutherine bulbosa* bulbs are composed of a vertically compressed stem with apical and axillary buds encircled by multiple layers of overlapping modified leaves (enlarged leaf bases) that resemble onions and serve as the main organs for storage (Borges et al., 2020; Tribble et al., 2021). Bulbs can store carbohydrates and other nutrients due to thickening development, which requires cell expansion following proliferation at different lateral meristems (Plunkert et al., 2024). Bulb initiation and growth are regulated by exogenous and environmental factors such as phytohormones. Phytohormones regulate plant growth and are thought to have a crucial role in bulb formation and bulb enlargement (Atif et al., 2020).

\* Corresponding author.

E-mail address: [vtbmai@hcmus.edu.vn](mailto:vtbmai@hcmus.edu.vn) (Bach Mai Thi Vo)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Gibberellins are among the most important (Shu et al., 2024). A phytohormone that promotes growth, gibberellins increase the quantity of cloves per bulb and boost their production efficiency by promoting cell elongation, leaf expansion, axillary meristem initiation, and lateral bud creation (Liu et al., 2020; Khokhar, 2023; Ritonga et al., 2023). Furthermore, GA<sub>3</sub> can promote the growth of lateral buds and enhance tuber count (Liu et al., 2019). Therefore, applying GA<sub>3</sub> at appropriate concentrations to affect bulblet formation and number, as well as the storage compounds in the bulb, is the main objective of this study.

## 2. MATERIALS AND METHODS

### 2.1. Materials

- *E. bulbosa* mother bulbs from the National Institute of Medical Materials (Figure 2A and B)
- The 6-, 12-, and 18-week-old *E. bulbosa* were grown in the experimental garden of the Department of Plant Physiology, University of Sciences, Vietnam National University, Ho Chi Minh City (Figures 1, 2C, D, and E)



Figure 1. *E. bulbosa* plant in the experimental garden.




Figure 2. (A) *E. bulbosa* mother bulb (A) and after removing the dry leaf layer showing buds (B) (arrows) and (C) the 6-week-old, (D) the 12-week-old, (E) the 18-week-old from the mother bulb was grown in the experimental garden.

## 3. METHODS

### 3.1. Study on the effect of GA<sub>3</sub> on *E. bulbosa* bulblet formation from mother bulbs

*E. bulbosa* mother bulbs were treated with GA<sub>3</sub> by soaking in GA<sub>3</sub> solution at different concentrations (from 20 to 50 ppm) (0.5 cm from the base) for a week, then placed in pots with a volume of (20 cm × 21 cm × 20 cm) containing organic soil and coconut fiber mixed in a 1:1 ratio. After 12 weeks, the following parameters were determined: number of bulblets, diameter, fresh weight, dry weight, bulb length, respiration intensity, total sugar content, flavonoid, and polyphenol content (PC).

### 3.2. Study on the effect of GA<sub>3</sub> after 6 weeks of treatments at the 6-week-old plant stage of *E. bulbosa*

*E. bulbosa* mother bulbs had the outer dry leaf scales removed, the leaf top and old roots cut off, and were soaked in water (0.5 cm from the base) for a week. After a week, the mother bulbs developed about 2 leaves and 7–10 roots, which were placed in plastic cups with a volume of (95 mm × 58 mm × 125 mm), containing organic soil and coconut fiber mixed in a 1:1 ratio. After 6 weeks of planting, the seedlings from the original mother bulbs were transferred to pots (20 cm × 21 cm × 20 cm) to adapt for 3 days and treated with GA<sub>3</sub> at different concentrations (from 20 to 50 ppm). The parameters, including number of bulblets, bulb diameter, fresh weight, dry weight, and length, total sugar content, respiration intensity, flavonoid, and PC in bulb on week 12th, were determined.

### 3.3. Study on the effect of GA<sub>3</sub> on the growth stages of *E. bulbosa*

*E. bulbosa* mother bulbs were treated with GA<sub>3</sub> by soaking in GA<sub>3</sub> 30 ppm for a week, then placed in pots with a volume of (20 cm × 21 cm × 20 cm) containing organic soil and coconut fiber mixed in a 1:1 ratio. A 40 ppm GA<sub>3</sub> treatment was continued at weeks 6, 12, and 18. The experiment consisted of six treatments (Table 1). The parameters for fresh weight, dry weight, number of bulbs, bulb yield per pot, flavonoid, and PC were determined at the 24th week of treatment.

### 3.4. Analysis of physiological and biochemical parameters

- Determination of the diameter and length of the bulb: The diameter of the bulb is measured with a ruler, clamped

**Table 1**

Processing method effect of GA<sub>3</sub> on the growth stages of *E. bulbosa*.

Treatments	Processing method			
	Week 0	Week 6	Week 12	Week 18
Control 1	Water	Water	Water	Water
Control 2	GA <sub>3</sub> 30 ppm	Water	Water	Water
GA <sub>3</sub> (w6)	GA <sub>3</sub> 30 ppm	GA <sub>3</sub> 40 ppm	Water	Water
GA <sub>3</sub> (w12)	GA <sub>3</sub> 30 ppm	Water	GA <sub>3</sub> 40 ppm	Water
GA <sub>3</sub> (w18)	GA <sub>3</sub> 30 ppm	Water	Water	GA <sub>3</sub> 40 ppm
GA <sub>3</sub> (w6+w12+w18)	GA <sub>3</sub> 30 ppm	GA <sub>3</sub> 40 ppm	GA <sub>3</sub> 40 ppm	GA <sub>3</sub> 40 ppm

at the largest area of edema, and the length of the bulb is measured with a ruler in millimeters, from the position of the root to the position where the leaf sheath separates.

- Determination of fresh and dry weight of bulbs: *E. bulbosa* bulbs were weighed using an analytical balance with an error of 0.01 mg of OHAUS (USA). Then, dried at 120 °C for 1 hour, then at 80 °C for about 72 hours until the weight remained unchanged to determine the dry weight (Sade et al., 2015).
- Determination of total sugar content of bulbs: Total sugar content in *E. bulbosa* bulb were determined using the phenol-sulfuric acid method according to Combs et al. (1987) and González-Vázquez (2022). Absorbance was measured at a wavelength of 490 nm using a spectrophotometer, compared to the sucrose standard curve to obtain the total sugar content.
- Measurement of respiratory intensity: *E. bulbosa* bulb during treatment was determined using a Leaf Lab 2 oxygen electrode gas exchanger (Hansatech, UK) with an improved measuring chamber with a height of 6 cm. Oxygen exchange rate was calculated based on the amount of oxygen absorbed per gram of fresh weight per minute  $\mu\text{mol O}_2 \text{ g}^{-1} \text{ min}^{-1}$ .
- Determination of total phenolic and flavonoid content Polyphenol and flavonoid of bulb dried powder were extracted in 60 % ethanol at a ratio of 1:35 (w/v) (at pH 3.0) for 1 h at 50 °C, then the extract was filtered and allowed to dry. PC was determined spectrophotometrically using the Folin–Ciocalteu method described by Shi et al. (2019), which used a color reaction with Folin–Ciocalteu reagent and 20 % Na<sub>2</sub>CO<sub>3</sub> with gallic acid as the standard, then measured OD at 765 nm. Aluminum chloride spectrophotometry was used to assess total flavonoid content (TFC), as described by Atanassova et al. (2011), which involved a color reaction with 5 % NaNO<sub>2</sub> and 10 % AlCl<sub>3</sub>, with OD measured at 510 nm. Rutin was used as

a standard. Polyphenol and flavonoid content in bulb were expressed as mg/g dry weight.

#### ■ Experimental design and statistical analysis

The plants are placed in the experimental garden under light intensity of 10,000–25,000 lux, temperature of 33 ± 5 °C, and humidity of 60 – 65 %. Ten replicates of three plants each were used in the fully randomized design of the studies. Plants were fertilized with NPK 16:16:16 (starting at week 4, once every 4 weeks). Watering was performed once daily at 4 p.m., with a uniform spray until the soil and leaves were wet.

SPSS (Statistical Package for the Social Sciences) 20.0 was used to analyze the data. Statistical differences were significant at the 95 % level, and results were expressed as means ± standard deviations, with accompanying text samples.

## 4. RESULTS


### 4.1. Effect of GA<sub>3</sub> on growth and bulblet formation of *E. bulbosa*

After 12 weeks of treatment, in the 30 ppm GA<sub>3</sub> treatment, the number of tubers increased rapidly to reach 6–7 tubers, much higher than the control (2–3 bulbs) and other treatments (2–3 or 5–6 bulbs) (Figure 3A–E). The number of bulbs, bulb fresh weight, and dry weight increased sharply at 30 ppm GA<sub>3</sub> treatment, then declined at 40 ppm and dropped markedly at 50 ppm. Bulb diameter was reduced in all treatments compared to the control, and the sharpest drop was observed in the 50 ppm GA<sub>3</sub> treatment (Table 2 and Figure 3 A1E1). Bulb length also increased sharply at 30 ppm and 40 ppm GA<sub>3</sub> treatment but decreased at 50 ppm GA<sub>3</sub> treatment (Table 2).

The sugar and starch content decreased from 20 ppm to 30 ppm after GA<sub>3</sub> treatment and then remained constant at 50 ppm treatment. The polyphenol and flavonoid levels began to increase at 30 ppm GA<sub>3</sub> treatment and then remained constant at 50 ppm GA<sub>3</sub> treatment. The respiration intensity of bulbs also increased sharply at 30 ppm GA<sub>3</sub> treatment and then decreased, especially at 50 ppm GA<sub>3</sub> treatment (Table 3).

### 4.2. Effect of GA<sub>3</sub> after 6 weeks of treatments at the 6-week-old plant stage of *E. bulbosa*

After 6 weeks of treatment at the 6-week-old plant stage, there was no significant difference in bulb count between the control and the treatments (Figure 4A–E), except for the 50 ppm GA<sub>3</sub> treatment, which had fewer bulbs. The bulb diameter, fresh weight, and dry weight gradually increased with the GA<sub>3</sub> treatment concentration and peaked at the GA<sub>3</sub> 40 ppm,



**Figure 3.** *Eleutherine bulbosa* after 12 weeks of treatments, with (A): Control; (B): GA<sub>3</sub> 20 ppm; (C): GA<sub>3</sub> 30 ppm; (D): GA<sub>3</sub> 40 ppm; (E): GA<sub>3</sub> 50 ppm and horizontal cutting bulb detail with (A1) Control (A1); (B1): GA<sub>3</sub> 20 ppm; (C1): GA<sub>3</sub> 30 ppm; (D1): GA<sub>3</sub> 40 ppm; (E1): GA<sub>3</sub> 50 ppm.

**Table 2**

Changes in bulb diameter, fresh weight (FW), dry weight (DW), bulb length, and the number of leaf sheaths of *E. bulbosa* from mother bulbs after 12 weeks of treatments

Treatments	Number of bulblets	Bulb diameter (mm)	Bulb length (cm)	Fresh weight (g)	(g)DW. g <sup>-1</sup> FW
mother bulbs					
Control	2.6 ± 0.2 <sup>c</sup>	13.14 ± 0.15 <sup>a</sup>	4.12 ± 0.08 <sup>c</sup>	1.153 ± 0.009 <sup>e</sup>	0.346 ± 0.008 <sup>d</sup>
GA <sub>3</sub> 20 ppm	2.8 ± 0.2 <sup>c</sup>	12.14 ± 0.14 <sup>b</sup>	4.24 ± 0.08 <sup>b</sup>	1.379 ± 0.008 <sup>d</sup>	0.418 ± 0.006 <sup>c</sup>
GA <sub>3</sub> 30 ppm	6.8 ± 0.4 <sup>a</sup>	12.14 ± 0.16 <sup>b</sup>	4.52 ± 0.06 <sup>a</sup>	1.930 ± 0.010 <sup>a</sup>	0.692 ± 0.008 <sup>a</sup>
GA <sub>3</sub> 40 ppm	5.8 ± 0.4 <sup>b</sup>	12.12 ± 0.11 <sup>b</sup>	4.54 ± 0.02 <sup>a</sup>	1.643 ± 0.009 <sup>b</sup>	0.563 ± 0.009 <sup>b</sup>
GA <sub>3</sub> 50 ppm	3.2 ± 0.2 <sup>c</sup>	11.06 ± 0.15 <sup>c</sup>	4.10 ± 0.07 <sup>c</sup>	1.450 ± 0.012 <sup>c</sup>	0.439 ± 0.009 <sup>c</sup>

The average numbers in the column with different letters are meaningful at  $p \leq 0.05$ .

**Table 3**


Total sugar content (TSC), starch content (StC), flavonoids, polyphenols content, and respiration intensity (RI) of *E. bulbosa* bulbs after 12 weeks of treatment with GA<sub>3</sub>.

Treatments	TSC (mg.g <sup>-1</sup> DW)	StC (mg.g <sup>-1</sup> DW)	Flavonoid (mg.g <sup>-1</sup> DW)	Polyphenol (mg.g <sup>-1</sup> DW)	RI (μLO <sub>2</sub> .g <sup>-1</sup> FW.h <sup>-1</sup> )
Control	32.23 ± 0.07 <sup>a</sup>	27.24 ± 1.5 <sup>a</sup>	2.89 ± 0.10 <sup>c</sup>	5.07 ± 0.05 <sup>b</sup>	53.50 ± 1.09 <sup>c</sup>
GA <sub>3</sub> 20 ppm	29.75 ± 1.15 <sup>b</sup>	22.68 ± 1.34 <sup>b</sup>	3.01 ± 0.02 <sup>c</sup>	5.09 ± 0.08 <sup>b</sup>	50.16 ± 1.46 <sup>c</sup>
GA <sub>3</sub> 30 ppm	26.02 ± 0.50 <sup>c</sup>	19.26 ± 1.21 <sup>c</sup>	3.54 ± 0.02 <sup>b</sup>	5.15 ± 0.11 <sup>ab</sup>	73.22 ± 0.02 <sup>a</sup>
GA <sub>3</sub> 40ppm	26.48 ± 0.50 <sup>c</sup>	19.41 ± 1.18 <sup>c</sup>	3.56 ± 0.01 <sup>b</sup>	5.23 ± 0.11 <sup>a</sup>	59.23 ± 0.72 <sup>b</sup>
GA <sub>3</sub> 50 ppm	27.73 ± 0.19 <sup>c</sup>	19.27 ± 1.51 <sup>c</sup>	3.29 ± 0.04 <sup>b</sup>	5.35 ± 0.12 <sup>a</sup>	48.19 ± 0.02 <sup>d</sup>

The average numbers in the column with different letters are meaningful at  $p \leq 0.05$ .

after which they declined at 50 ppm (Table 3 and Figure 3A1–E1). Similarly, bulb length increased most at 30 and 40 ppm GA<sub>3</sub> and decreased sharply at 50 ppm GA<sub>3</sub> (Table 3). Sugar and starch contents were reduced in all treatments relative to the control. The starch content plummeted at the 40 ppm GA<sub>3</sub>

treatment and then remained unchanged at the 50 ppm GA<sub>3</sub> treatment, whereas the sugar content plummeted at the 50 ppm GA<sub>3</sub> treatment. The polyphenol and flavonoid contents increased sharply at 40 ppm GA<sub>3</sub> treatment and then remained constant at 50 ppm GA<sub>3</sub> treatment (Table 5).



**Figure 4.** *E. bulbosa* after 6 weeks of treatments, with (A1): Control; (B1): GA<sub>3</sub> 20 ppm; (C1): GA<sub>3</sub> 30 ppm; (D1): GA<sub>3</sub> 40 ppm; (E1): GA<sub>3</sub> 50 ppm and horizontal cutting bulb detail with (A1) Control; (B1): GA<sub>3</sub> 20 ppm; (C1): GA<sub>3</sub> 30 ppm; (D1): GA<sub>3</sub> 40 ppm; (E1): GA<sub>3</sub> 50 ppm.

**Table 4**

Changes in bulb diameter, fresh weight (FW), dry weight (DW), bulb length, and the number of leaf sheaths of *E. bulbosa* from after 6 weeks of treatments at the 6-week-old plant stage.

Treatments	Number of bulblets	Bulb diameter (mm)	Bulb length (cm)	Fresh weight (g)	DW. g <sup>-1</sup> FW (g)
Control	2.7 ± 0.2 <sup>a</sup>	10.06 ± 0.06 <sup>d</sup>	4.13 ± 0.07 <sup>c</sup>	1.14 ± 0.06 <sup>d</sup>	0.35 ± 0.03 <sup>d</sup>
GA <sub>3</sub> 20 ppm	2.7 ± 0.2 <sup>a</sup>	11.01 ± 0.10 <sup>c</sup>	4.41 ± 0.12 <sup>b</sup>	1.42 ± 0.09 <sup>c</sup>	0.43 ± 0.03 <sup>c</sup>
GA <sub>3</sub> 30 ppm	3.3 ± 0.4 <sup>a</sup>	11.50 ± 0.06 <sup>b</sup>	4.70 ± 0.12 <sup>a</sup>	1.86 ± 0.02 <sup>b</sup>	0.67 ± 0.03 <sup>b</sup>
GA <sub>3</sub> 40 ppm	3.0 ± 0.1 <sup>a</sup>	12.93 ± 0.03 <sup>a</sup>	4.73 ± 0.09 <sup>a</sup>	2.01 ± 0.03 <sup>a</sup>	0.88 ± 0.02 <sup>a</sup>
GA <sub>3</sub> 50 ppm	1.9 ± 0.2 <sup>b</sup>	8.27 ± 0.06 <sup>d</sup>	4.13 ± 0.06 <sup>c</sup>	1.08 ± 0.02 <sup>d</sup>	0.46 ± 0.03 <sup>c</sup>

The average numbers in the column with different letters are meaningful at  $p \leq 0.05$ .

**Table 5**

Total sugar content (TSC), starch content (StC), flavonoids, polyphenols content, and respiration intensity (RI) of *E. bulbosa* bulbs after 6 weeks of treatment with GA<sub>3</sub> at the 6-week-old plant stage.

Treatments	TSC (mg.g <sup>-1</sup> DW)	StC (mg.g <sup>-1</sup> DW)	Flavonoid (mg.g <sup>-1</sup> DW)	Polyphenol (mg.g <sup>-1</sup> DW)	RI (μLO <sub>2</sub> g <sup>-1</sup> FW.h <sup>-1</sup> )
Control	32.23 ± 0.07 <sup>a</sup>	17.24 ± 0.5 <sup>a</sup>	2.93 ± 0.10 <sup>d</sup>	5.01 ± 0.05 <sup>c</sup>	51.32 ± 1.09 <sup>c</sup>
GA <sub>3</sub> 20 ppm	30.73 ± 0.19 <sup>b</sup>	12.68 ± 0.34 <sup>b</sup>	3.41 ± 0.01 <sup>c</sup>	5.29 ± 0.08 <sup>b</sup>	55.76 ± 1.46 <sup>c</sup>
GA <sub>3</sub> 30 ppm	29.82 ± 0.50 <sup>b</sup>	11.26 ± 0.21 <sup>c</sup>	5.53 ± 0.02 <sup>b</sup>	5.39 ± 0.11 <sup>b</sup>	60.28 ± 0.72 <sup>b</sup>
GA <sub>3</sub> 40 ppm	27.75 ± 0.15 <sup>c</sup>	10.41 ± 0.98 <sup>d</sup>	6.55 ± 0.01 <sup>a</sup>	6.60 ± 0.11 <sup>a</sup>	78.22 ± 0.02 <sup>a</sup>
GA <sub>3</sub> 50 ppm	23.48 ± 0.50 <sup>d</sup>	10.27 ± 0.57 <sup>d</sup>	6.49 ± 0.04 <sup>a</sup>	6.35 ± 0.12 <sup>a</sup>	59.11 ± 0.02 <sup>d</sup>

The average numbers in the column with different letters are meaningful at  $p \leq 0.05$ .

### 4.3. Effect of GA<sub>3</sub> combined treatments on the growth stages of *E. bulbosa*

The fresh and dried weight of the bulbs increased in all treatments compared to the control. Between the 2 control experiments, Control 2, which was treated with 30 ppm GA<sub>3</sub> at week 0 (mother bulb), had a significantly higher fresh and dry weight. There was no difference in fresh and dry weight between GA<sub>3</sub> treatments at weeks 6, 12, and 18. When combined treatments were applied at three stages, these indicators reached their highest values. Likewise, flavonoid and PCs and tuber yield per pot also showed the greatest increases under the combined-stage treatment. The polyphenol content (PC) in Control 2 increased compared to Control 1, but flavonoid content did not differ. Comparing single-stage applications showed that GA<sub>3</sub> at week 12 and week 18 did not differ, and both increased markedly versus week 6, while PC at week 18 increased sharply compared with the other 2 weeks. However, bulb yield per pot did not differ among single-stage treatments (Table 6).

## 5. DISCUSSION

### 5.1. Effect of GA<sub>3</sub> on growth and bulblet formation

*Eleutherine bulbosa* bud derived from mother bulbs after 12 weeks of GA<sub>3</sub> treatment showed that the number of bulblets, bulb length, and both fresh and dry weight gradually increased at GA<sub>3</sub> concentrations of 20–30 ppm, with a significant improvement recorded at 30 ppm compared with the control. In contrast, a higher level (50 ppm) caused a decrease in these parameters (Table 2). This suggests that GA<sub>3</sub>, like many other plant growth regulators, has a concentration-dependent dual effect. At moderate levels, it promotes axillary meristem initiation and stimulates cell elongation (Liu et al., 2020), thereby enhancing bulblet formation and elongation

in *E. bulbosa*, which contributes to increased fresh weight. However, when the hormone is applied at excessive concentrations, growth parameters are suppressed. These results are similar to reports on garlic and lily that GA<sub>3</sub> at appropriate levels enhanced bulblet production, but excessive application reduced yield (Khokhar, 2023; Anisah et al., 2023).

Furthermore, during the early growth stage of *E. bulbosa*, exogenous GA<sub>3</sub> application may impact both bulblet initiation and longitudinal cell expansion within the bulb. This results in an increase in bulblet number and bulb length but is often accompanied by a reduction in bulb diameter, as studied in *Tulipa saxatilis* (Sari, 2024). This may be explained by the fact that gibberellins regulate growth mainly by stimulating structural proteins and enzymes to promote cell elongation (Kou et al., 2021), rather than enhancing radial expansion.

The effect of GA<sub>3</sub> on bulb information and growth in *E. bulbosa* also depends on the bulbs' growth stage. At the early stage of growth 0 week (mother bulb stage), the optimal concentration for improving growth goals is GA<sub>3</sub> 30 ppm, especially increasing the number of bulbs while at the 6 week stage, GA<sub>3</sub> at 40 ppm significantly increases fresh weight, dry weight, and length, but does not increase the number of bulbs, and at 50 ppm all parameters decrease (Figures 3 and 4; Table 4). This indicates that in later stages of growth, GA<sub>3</sub> favors biomass accumulation rather than only bulblet initiation. These results correspond to the role of GA in promoting cell expansion in preparation for dry matter build-up in the storage organs (Liao et al., 2025).

### 5.2. Effect of GA<sub>3</sub> on secondary compounds

GA<sub>3</sub> treatments increase respiration and reduce sugar and starch levels both early from mother bulbs and at 6 weeks (Table 5). This may be because exogenous gibberellin inhibits the activity of sucrosesynthase (SS) and soluble starch

**Table 6**

Changes in bulb fresh weight (FW), dry weight (DW), flavonoid content, polyphenol content, and bulb yield per plant of *E. bulbosa* after 12 weeks in treatments GA<sub>3</sub> at week 12 (the 24-week-old plant).

Treatments	FW (g) bulb/plant	DW/g FW	Flavonoid (mg/g DW)	Polyphenol (mg/g DW)	Bulb yield (g)/pot
Control 1	78.55 ± 0.02 <sup>d</sup>	4.92 ± 0.15 <sup>d</sup>	4.51 ± 0.11 <sup>d</sup>	5.25 ± 0.07 <sup>e</sup>	221.55 ± 3.69 <sup>d</sup>
Control 2	103.79 ± 0.02 <sup>c</sup>	5.73 ± 0.07 <sup>c</sup>	4.72 ± 0.12 <sup>d</sup>	5.81 ± 0.07 <sup>d</sup>	380.59 ± 3.06 <sup>e</sup>
GA <sub>3</sub> (w6)	180.15 ± 0.05 <sup>b</sup>	6.13 ± 0.18 <sup>b</sup>	7.05 ± 0.08 <sup>c</sup>	7.35 ± 0.08 <sup>c</sup>	543.56 ± 6.36 <sup>b</sup>
GA <sub>3</sub> (w12)	183.13 ± 0.05 <sup>b</sup>	6.23 ± 0.18 <sup>b</sup>	7.35 ± 0.08 <sup>b</sup>	7.45 ± 0.04 <sup>c</sup>	550.06 ± 6.34 <sup>b</sup>
GA <sub>3</sub> (w18)	184.12 ± 0.05 <sup>b</sup>	6.23 ± 0.18 <sup>b</sup>	7.33 ± 0.08 <sup>b</sup>	7.85 ± 0.05 <sup>b</sup>	560.06 ± 7.24 <sup>b</sup>
GA <sub>3</sub> (w6+w12+w18)	208.89 ± 0.04 <sup>a</sup>	6.82 ± 0.09 <sup>a</sup>	7.60 ± 0.13 <sup>a</sup>	8.08 ± 0.05 <sup>a</sup>	660.19 ± 1.68 <sup>a</sup>

The average numbers in the column with different letters are meaningful at  $p \leq 0.05$ .

synthase (SSS), thereby reducing the sucrose and starch content in the tuber (Li et al., 2021) and increasing the activity of the enzyme responsible for starch hydrolysis, similar to the study by Siega et al. (2025). Moreover, when sugar and starch levels decreased, flavonoid levels in both stages increased, slightly increasing in the early stages, at 30 ppm GA<sub>3</sub>, sharply increasing in the 6-week stage, at 40–50 ppm GA<sub>3</sub>, suggesting that carbon reserves were mobilized for new bulblet growth in the early stages and promoted rapid consumption of carbohydrate reserves to provide energy and substrates for organ initiation (Shu et al., 2024), but at a later stage, the metamorphosis shifts towards the biosynthesis of secondary compounds, increasing the content of flavonoids and polyphenols. The effect of GA on flavonoids is due to gibberellin activating phenylalanine aminotransferase activity and promoting the phenylpropanoid metabolism pathway, thereby increasing TFC. Similar to many crops, GA<sub>3</sub> promotes the synthesis and accumulation of total phenolic compounds and specific flavonoids (Xu et al., 2025). This is in contrast with some reports that gibberellin signaling suppresses flavonoid biosynthesis in *Medicago truncatula* or apple (Sun et al., 2021), showing that the effect of GA is species- and stage-dependent.

In the repeated treatment experiments, the combination of 30 ppm GA<sub>3</sub> at planting and 40 ppm at later stages (6–18 weeks) resulted in the highest bulb fresh weight, dry weight, bulblet yield, and flavonoid and PCs. This indicates that multiple applications at different developmental stages provide cumulative effects, optimizing both propagation and medicinal quality of bulbs. This once again confirms that the timing of gibberellin application is very important for bulblet bulbose initiation and enlargement in *E. bulbosa*, similar to the results of the study in lily (Anisah et al., 2023). Thus, GA<sub>3</sub> not only regulates bulblet initiation and biomass accumulation but also promotes the accumulation of flavonoid and polyphenol through carbon mobilization, enhanced respiration, and activation of the phenylpropanoid pathway.

## 6. CONCLUSION

At the mother bulb stage, GA<sub>3</sub> at 30 ppm was optimal for increasing bulblet number, fresh and dry weight, bulb length, and respiration intensity. At the 6-week stage, GA<sub>3</sub> at 40 ppm enhanced biomass accumulation and secondary metabolite levels, whereas 50 ppm inhibited growth.

Flavonoid and polyphenol contents (9PCs) increased with GA<sub>3</sub>, reaching the highest levels at 40–50 ppm, depending on growth stage.

Sequential treatments combining 30 ppm at planting with 40 ppm at later stages yielded the greatest improvements in bulb yield and in flavonoid and polyphenol accumulation.

## ORCID

Tran Thi Thanh Hien 0009-0002-0213-3440  
 Vang Thi My Tu 0009-0006-6552-8609  
 Nguyen Kieu Uyen Vi 0009-0008-2043-1824  
 Vo Thi Bach Mai 0009-0006-6508-0234

## ACKNOWLEDGEMENTS

This research was supported by the Plant Physiology Laboratory, Department of Plant Physiology, Faculty of Biology-Biotechnology, University of Sciences, Vietnam National University in Ho Chi Minh City (VNU-HCM).

## ETHICAL CONSIDERATIONS

Not relevant (the writers pledge to take full responsibility for this).

## AUTHORS CONTRIBUTIONS

All trials were carried out by Nguyen Kieu Uyen Vy, Vang Thi My Tu, and Tran Thi Thanh Hien.

Tran Thi Thanh Hien wrote the paper and came up with the idea. The final manuscript was edited and authorized by Vo Thi Bach Mai.

## CONFLICTS OF INTEREST

There are no conflicts of interest for the authors.

## FUNDING

There is no funding for this study.

## REFERENCES

- Anisah, S.N., Ratnadewi, D.I.A.H., Supena, E.D.J., 2023. Exogenous gibberellic acid stimulates bulb dormancy breaking and the role of paclobutrazol in maintaining the size of harvested bulb of lily (*Lilium* sp.) cv. Tisento. *Sains Malaysiana* 52(7), 1967–1976. <https://doi.org/10.17576/jsm-2023-5207-06>
- Atanassova, M., Georgieva, S., Ivancheva, K., 2011. Total phenolic and total flavonoid contents, antioxidant capacity and biological contaminants in medicinal herbs. *Journal of the University of Chemical Technology and Metallurgy* 46, 81–88.
- Atif, M.J., Ahanger, M.A., Amin, B., Ghani, M.I., Ali, M., Cheng, Z., 2020. Mechanism of allium crops bulb enlargement in response

- to photoperiod: a review. *International Journal of Molecular Sciences* 21(4), 1325. <https://doi.org/10.3390/ijms21041325>
- Borges, E.S., Galucio, N.C. da R., Veiga, A.S.S., Busman, D.V., Lins, A.L.F. de A., Bahia, M. de O., Rissino, J.D., Correa, R.M. dos S., Burbano, R.M.R., Marinho, A.M.R., Casique, J.V., Percário, S., Dolabela, M.F., 2020. Botanical studies, antimicrobial activity and cytotoxicity of *Eleutherine bulbosa* (Mill.) Urb. *Research, Society and Development* 9(11), e3369119992. <https://doi.org/10.33448/rsd-v9i11.9992>
- Coombs, J., Hind, G., Leegood, G.C., Tieszen, L.L., Vonshak, A., 1987. Measurement of starch and sucrose in leaves. In: Coombs, J., Hall, D.O., Long, S.P., Scurlock, J.M.O. (Eds.), *Techniques in Bioproductivity and Photosynthesis*. Pergamon Press, Oxford, UK, pp. 219–228.
- Da Silva, R.M.G., Alves, C.P., Barbosa, F.C., Santos, H.H., Adão, K.M., Granero, F.O., Figueiredo, C.C.M., Nicolau-Junior, N., Silva, L.P., 2024. Antioxidant, antitumoral, antimetastatic effect and inhibition of collagenase enzyme activity of *Eleutherine bulbosa* (Dayak onion) extract: In vitro, in vivo and in silico approaches. *Journal of Ethnopharmacology* 318, 117005. <https://doi.org/10.1016/j.jep.2023.117005>
- Harlita, T.D., Oedjijono, Asnani, A., 2018. The antibacterial activity of Dayak onion (*Eleutherine palmifolia* (L.) Merr) towards pathogenic bacteria. *Tropical Life Sciences Research* 29(2), 39–52. <https://doi.org/10.21315/tlsr2018.29.2.4>
- Herman, H., Ibrahim, A., Junaidin, J., Arifuddin, M., Hikmawan, B.D., Siska, S., Bariroh, T., Purwoko, R.Y., Febrina, L., Faisal, M., Iswahyudi, I., Angelina, M., Samsul, E., Rijai, L., Ahmad, I., 2024. Pharmacognostic profile and anti-diabetic activity of *Eleutherine bulbosa* Mills. bulbs from East Kalimantan, Indonesia. *Pharmacognosy Journal* 16(1), 118–125. <https://doi.org/10.5530/pj.2024.16.16>
- González-Vázquez, M., Calderón-Domínguez, G., Mora-Escobedo, R., Salgado-Cruz, M.P., Arreguín-Centeno, J.H., Monterrubio-López, R., 2022. Polysaccharides of nutritional interest in jicama (*Pachyrhizus erosus*) during root development. *Food Science & Nutrition* 10(4), 1146–1158. <https://doi.org/10.1002/fsn3.2746>
- Jiang, H., Man, W.-J., Hou, A.-J., Yang, L., Xing, X.-D., Yan, M.-L., Guo, X.-Y., Yang, L., 2020. The chemical constituents from the active fractions of *Eleutherine bulbosa* with their antimicrobial activity. *Natural Product Research* 34(12), 1743–1749. <https://doi.org/10.1080/14786419.2018.1530229>
- Kamarudin, A.A., Esa, N.M., Saad, N., Sayuti, N.H., Razak, N.A.A., 2020. Heat assisted extraction of phenolic compounds from *Eleutherine bulbosa* (Mill.) bulb and its bioactive profiles using response surface methodology. *Industrial Crops & Products* 144, 112064. <https://doi.org/10.1016/j.indcrop.2019.112064>
- Khokhar, K.M., 2023. Bulb development in garlic – a review. *The Journal of Horticultural Science and Biotechnology* 98(4), 432–442. <https://doi.org/10.1080/14620316.2022.2150326>
- Kou, E., Huang, X., Zhu, Y., Su, W., Liu, H., Sun, G., Chen, R., Hao, Y., Song, S., 2021. Crosstalk between auxin and gibberellin during stalk elongation in flowering Chinese cabbage. *Scientific Reports* 11, 3976. <https://doi.org/10.1038/s41598-021-83519-z>
- Li, J., Seng, S., Li, D., Zhang, F., Liu, Y., Yao, T., Liang, J., Yi, M., Wu, J., 2021. Antagonism between abscisic acid and gibberellin regulates starch synthesis and corm development in *Gladiolus hybridus*. *Horticulture Research* 8, 155. <https://doi.org/10.1038/s41438-021-00589-w>
- Liao, Y., Zeng, Z., Lin, K., Jiang, W., Wang, J., Duan, L., Liang, X., Huang, Y., Han, Z., Hu, H., Xu, Z.-F., Ni, J., 2025. Gibberellin promotes xylem expansion and cell lignification by regulating sugar accumulation and the expression of JcMYB43 and JcMYB63 in the woody plant *Jatropha curcas*. *International Journal of Biological Macromolecules* 294, 139434. <https://doi.org/10.1016/j.ijbiomac.2024.139434>
- Liu, H., Wen, Y., Cui, M., Qi, X., Deng, R., Gao, J., Cheng, Z., 2020. Histological, physiological and transcriptomic analysis reveal gibberellin-induced axillary meristem formation in garlic (*Allium sativum*). *Plants* 9(8), 970. <https://doi.org/10.3390/plants9080970>
- Liu, H., Deng, R., Huang, C., Cheng, Z., Meng, H., 2019. Exogenous gibberellins alter morphology and nutritional traits of garlic (*Allium sativum* L.) bulb. *Scientia Horticulturae* 246, 298–306. <https://doi.org/10.1016/j.scienta.2018.11.027>
- Naspiyah, N., Iskandar, Y., Moelyono, M.W., 2014. Artikel Ulasan: Bawang Tiwai (*Eleutherine americana* Merr.), Tanaman Multiguna. *Indonesian Journal of Applied Sciences* 4(2), 18–30.
- Panyachariwat, N., Jimtaisong, A., Saewan, N., 2024. Antioxidative potentials of *Eleutherine bulbosa* bulb and its utilization in tropical cosmetic emulsion. *Cosmetics* 11(4), 111. <https://doi.org/10.3390/cosmetics11040111>
- Plunkert, M.L., Martínez-Gómez, J., Madrigal, Y., Hernández, A.I., Tribble, C.M., 2024. Tuber, or not tuber: Molecular and morphological basis of underground storage organ development. *Current Opinion in Plant Biology* 80, 102544. <https://doi.org/10.1016/j.pbi.2024.102544>
- Ritonga, F.N., Zhou, D., Zhang, Y., Song, R., Li, C., Li, J., Gao, J., 2023. The roles of gibberellins in regulating leaf development. *Plants* 12(6), 1243. <https://doi.org/10.3390/plants12061243>
- Sade, N., Galkin, E., Moshelion, M., 2015. Measuring Arabidopsis, tomato and barley leaf relative water content. *Bio-protocol* 5(7), e1451. <https://doi.org/10.21769/BioProtoc.1451>
- Sari, Ö., 2024. Determination of the effect of GA<sub>3</sub> applications on plant development, nutrient content change and analysis of root architectural features using ML artificial neural network modelling in *Tulipa saxatilis*. *Folia Horticulturae* 36(3), 381–398. <https://doi.org/10.2478/fhort-2024-0024>
- Shi, P., Du, W., Wang, Y., Teng, X., Chen, X., Ye, L., 2019. Total phenolic, flavonoid content and antioxidant activity of bulbs, leaves and flowers made from *Eleutherine bulbosa* (Mill.) Urb. *Food Science & Nutrition* 7(1), 148–154. <https://doi.org/10.1002/fsn3.834>
- Siega, Y.P., Coelho, C.M.M., Silva, M.B.P.D., Albuquerque, M.R.M., Padilha, M.S., Mendes, G.C., 2025. Impact of gibberellic acid on seedling growth and enzymatic activity in bean cultivars with contrasting seed vigor. *Acta Scientiarum. Agronomy* 47, e72032. <https://doi.org/10.4025/actasciagron.v47i1.72032>
- Shu, F., Wang, D., Sarsaiya, S., Jin, L., Liu, K., Zhao, M., Wang, X., Yao, Z., Chen, G., Chen, J., 2024. Bulbil initiation: A comprehensive review on resources, development, and utilisation, with emphasis on molecular mechanisms, advanced technologies, and future prospects. *Frontiers in Plant Science* 15, 1343222. <https://doi.org/10.3389/fpls.2024.1343222>
- Sun, H., Cui, H., Zhang, J., Kang, J., Wang, Z., Li, M., Yi, F., Yang, Q., Long, R., 2021. Gibberellins inhibit flavonoid biosynthesis and promote nitrogen metabolism in *Medicago truncatula*. *International Journal of Molecular Sciences* 22(17), 9291. <https://doi.org/10.3390/ijms22179291>

- Sun, Z., Li, Q., Zhang, J., Sun, Z., Wu, H., Liang, H., Ma, G., 2024. Phenolic constituents with antibacterial activity from *Eleutherine bulbosa*. *Fitoterapia* 172, 105787. <https://doi.org/10.1016/j.fitote.2023.105787>
- Tribble, C.M., Martínez-Gómez, J., Howard, C.C., Males, J., Sosa, V., Sessa, E.B., Cellinese, N., Specht, C.D., 2021. Get the shovel: morphological and evolutionary complexities of belowground organs in geophytes. *American Journal of Botany* 108(3), 372–387. <https://doi.org/10.1002/ajb2.1623>
- Xu, X., Zhang, X., Wang, J., Liu, X., Zheng, R., Wu, Y., 2025. Effects of exogenous additives on the nutrient composition and metabolites of buckwheat. *Revista Brasileira de Engenharia Agrícola e Ambiental* 29(7), e285526. <https://doi.org/10.1590/1807-1929/agriambi.v29n7e285526>