


Original Research

View Article online



Received 31 October 2025
Revised 23 December 2025
Accepted 07 January 2026
Available Online 17 March 2026

Edited by Kannan RR Rengasamy

KEYWORDS:

Anticancer activity
Antioxidant
Thirunethira Chooranam

<https://doi.org/10.53365/nrfhh/216449>
eISSN: 2583-1194
Copyright © 2026 Visagaa Publishing House

In Vitro Antioxidant and Cytotoxicity, and *In Vivo* Anticancer Evaluation of the Siddha Formulation *Thirunethira Chooranam*

Deepa Natarajan^{1*}, Sudha Revathy Senthilkumar², Meenashree Balakrishnan³, Kayalvizhi Duraisamy³

¹Dean & Professor, Faculty of Pharmacy, SBMCH Campus, Bharath Institute of Higher Education and Research (BIHER), Chennai-600044, Tamil Nadu, India

²Assistant Professor, Department of Gunapadam, National Institute of Siddha, India

³Asthagiri Herbal Research Foundation, Perungudi, Chennai-600096, Tamil Nadu, India

ABSTRACT: Cancer remains a major global health worry, particularly in developing nations, where economic limitations necessitate affordable therapeutic alternatives. The Siddha medicine, a traditional South Indian system of medicine, incorporates several formulations with potential anticancer properties, including *Thirunethira Chooranam* (TNC). To evaluate the *in vitro* antioxidant and cytotoxic, and *in vivo* anticancer activity of TNC using the Dalton's lymphoma ascites (DLA) cells-induced tumor model in Swiss albino mice. Antioxidant activity was assessed using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay. Cytotoxic potential was examined against estrogen-responsive breast cancer cell line (MCF-7), lung adenocarcinoma epithelial cell line (A549), hepatocellular carcinoma cell line (HEP-G2), human cervical cancer cell line (HeLa), and African green monkey kidney (Vero) cell line via 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay. *In vivo* efficacy was evaluated in DLA-bearing mice by analyzing hematological, biochemical, cancer cell count, survival, and histopathological parameters. TNC demonstrated dose-dependent antioxidant activity (half-maximal inhibitory concentration [IC₅₀] = 7 mg/mL). The MTT assay revealed selective cytotoxicity, with the strongest effect against MCF-7 cells (IC₅₀ = 14.39 mg/mL) while sparing normal Vero cells. *In vivo* studies showed significant restoration of hematological and biochemical indices, an 80% increase in life span, reduction in tumor cell count, and partial restoration of hepatic and renal architecture. The results validated that TNC demonstrated its *in vitro* antioxidant and cytotoxic effects, and *in vivo* anticancer efficacy as a cost-effective and potential drug for cancer treatment.

1. INTRODUCTION

Cancer is one of the leading causes of morbidity and mortality all over the world. According to GLOBOCAN 2022 (an online database providing current, comprehensive

cancer incidence), there were 19.96 million new patients and 9.74 million deaths globally, and this number is projected to rise to 35 million new patients annually by 2050 (Bray et al., 2024; World Health Organization (WHO), 2024). In India, cancer accounted for 1.41 million new patients and 916,827

* Corresponding author.

E-mail address: sudharevathy_s@yahoo.com (Sudha Revathy Senthilkumar)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

deaths in 2022, reflecting the country's growing cancer burden. The most common cancers were breast (13.6%), lip and oral cavity (10.2%), cervix uteri (9%), and lung (5.8%). Projections suggest that India would have more than 2.1 million new patients every year by 2040, further intensifying the strain on healthcare systems if effective strategies are not implemented (International Agency for Research on Cancer [IARC], 2022).

Access to recently developed cancer drugs remains highly restricted in many low- and middle-income countries (LMICs). Of the 568 cancer drugs introduced between 1990 and 2022, around 35% never became available outside the single country where they were first approved, and fewer than 30% reached >10 countries. Most nations waited 1½–3 years after the initial launch prior to gaining access, with the longest delays occurring in LMICs. As a result, high-income regions secured earlier and broader availability, widening the existing treatment inequalities (Li et al., 2024).

Even when new cancer drugs reach LMICs, they often create a heavy financial burden on patients and their families. Cancer care largely remains out of reach, so households frequently borrow, sell assets, or stop treatment, leading to delays and poorer outcomes (Mohanty et al., 2024).

These challenges highlight the urgent need for cost-effective and accessible therapeutic approaches to reduce both clinical and economic burden of cancer. In this context, herbal medicines rooted in traditional systems have gained global attention. Emerging clinical evidence suggests that adjunctive use of herbal medicines can improve quality of life and enhance survival in cancer patients, with acceptable cost-effective profiles in certain healthcare systems (Tang et al., 2024).

Natural sources have been fundamental to the discovery of anticancer agents. Analyses of the US Food and Drug Administration (FDA) approvals from 1981 to 2019 show that nearly two-thirds of small-molecule anticancer drugs originated from natural products, their derivatives, or synthetic compounds modelled on natural structures (Newman & Cragg, 2020). Several widely used chemotherapeutic agents are vincristine and vinblastine from *Catharanthus roseus* and paclitaxel from *Taxus brevifolia* (Cragg & Pezzuto, 2016; Cragg et al., 2012). These landmark discoveries highlight the structural diversity and therapeutic potential of natural products in oncology.

Beyond single molecules, polyherbal formulations offer distinct advantages by combining multiple botanicals that act on diverse molecular targets, producing synergistic effects, improving bioavailability, and reducing toxicity (Wagner & Ulrich-Merzenich, 2009). This principle forms the basis of traditional medical systems, such as Ayurveda, Siddha, and traditional Chinese medicine, where complex formulations are preferred over isolated constituents.

The Siddha medicine, one of the oldest medical systems of South India, is rooted in Tamil culture and history. Its foundation is attributed to the *Siddhars*, ancient Tamil sages, and its origins can be traced back to the Sangam period (200 BCE–200 CE) (Kanagarathinam & Lourdasamy, 2023; Subbarayappa, 1997). The Siddha medical system is known for its extensive use of polyherbal formulations, many of which are indicated for chronic and degenerative diseases, including conditions resembling cancer. Among them, *Thirunethira Chooranam* (TNC) is a classical formulation comprising *vellarugu* (*Enicostemma axillare*), *shivanarvembu* (*Indigofera aspalathoides*), and *sangan kuppi* (*Clerodendrum inerme*). Although these plants have been individually studied for their pharmacological potential, the combined efficacy of TNC has not been validated scientifically. The present study aims to evaluate the *in vitro* antioxidant and cytotoxic, and *in vivo* anticancer activities of TNC, thereby bridging the Siddha medicine with modern pharmacological evidence.

2. MATERIALS AND METHODS

2.1. Preparation of test drug

Thirunethira Chooranam was prepared in accordance with classical Siddha literature (*Agasthiyar Vaithiya Vallathi – 600*). Raw materials for TNC were purchased from a licensed vendor in Chennai and authenticated by a botanist. After cleaning, shade-drying, and purification as per Siddha texts, the ingredients were pulverized, mixed in equal proportions, sieved to a fine powder, and stored in airtight containers. The ingredients used in the preparation of TNC are listed in Table 1 and the final products with ingredients are depicted in Figure 1.

2.2. *In Vitro* evaluation of antioxidant potential and cytotoxic effects

2.2.1. *In vitro* antioxidant activity evaluation using 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging assay

The antioxidant activity of TNC was determined by the DPPH radical scavenging assay with slight modifications

Table 1

Ingredients of *Thirunethira Chooranam*.

S. No.	Tamil name	Botanical name	Family
1.	<i>Vellarugu</i>	<i>Enicostemma axillare</i>	Gentianaceae
2.	<i>Shivanarvembu</i>	<i>Indigofera aspalathoides</i>	Fabaceae
3.	<i>Sanganakuppi</i>	<i>Clerodendrum inerme</i>	Lamiaceae




Figure 1. Ingredients of *Thirunethira Chooranam*.

(Koleva et al., 2002). The DPPH solution was freshly prepared in methanol at a strength of 0.2 mM. Various concentrations of TNC extract (0.5, 1, 2, 4, 8, and 16 mg/mL) were prepared in methanol, and 1 mL of each sample solution was mixed with 5 mL of DPPH reagent. The solution was vortexed and incubated at room temperature for 30 minutes in darkness. Methanol was used as a blank, while ascorbic acid (10, 20, 30, 40, 50, and 60 µg/mL) served as a positive control. After incubation, the reaction mixture was measured at 517-nm wavelength using a UV-visible spectrophotometer. The proportion of DPPH radical scavenging was evaluated using the following equation:

$$\text{DPPH radical scavenging effect (\%)} = \frac{A_0 - A_1}{A_0} \times 100,$$

where A_0 is the absorbance of the control, and A_1 is the absorbance of the test sample.

All experiments were conducted in triplicate, and the results were expressed as mean values. The half-maximal inhibitory concentration (IC_{50}) values for both TNC and ascorbic acid were also documented.

2.2.2. In vitro cytotoxicity evaluation using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay

By using Dulbecco's modified eagle medium (DMEM) having 10% fetal bovine serum (FBS), the cell counts of lung adenocarcinoma epithelial cell line (A549), estrogen-responsive breast cancer cell line (MCF-7), human cervical cancer cell line (HeLa), hepatocellular carcinoma cell line (HEP-G2), and African green monkey kidney (Vero) cell line were adjusted to 1.0×10^5 cells/mL. Approximately 100 µL of diluted cell suspension was added to each well of a 96-well microtiter plate. After 24 hours, cells were

centrifuged, and pellets were resuspended with 100 µL of test sample concentrations (31.25–250 µg/mL). Plates were incubated at 37°C for 48 hours in a 5% CO_2 atmosphere, with microscopic examination at 24-hour intervals. After 48 hours, solutions were centrifuged, and pellets were resuspended with 20 µL of MTT (2 mg/mL) in phenol red-free medium (MEM-PR). Further, the plates were incubated in a 5% CO_2 atmosphere for 2 hours at 37°C. To the plates 100 µL of dimethyl sulfoxide (DMSO) was added slowly and mixed gently to solubilize the formed formazan. Absorbance was read at 540 nm with a microplate reader (Mosmann, 1983). The proportion of cell viability was calculated using the following formula, and the concentration of test samples required to inhibit cell growth by 50% (IC_{50}) was derived from dose–response curves:

% Cell viability =

$$\frac{\text{Mean optical density [OD] of individual test group}}{\text{mean OD of control group}} \times 100.$$

2.3. In vivo anticancer evaluation using Dalton's lymphoma ascites (DLA) cells

2.3.1. Induction of cancer using DLA cells

The DLA model was employed to assess anticancer activity, as widely reported in earlier studies (Thavamani et al., 2014). DLA cells were obtained from Amala Cancer Research Center, Kerala, India. The DLA cells were maintained in Swiss albino mice (*in vivo*) through intraperitoneal (i.p.) transplantation. The DLA cells were aspirated from the peritoneal cavity of previously inoculated mice with saline solution to induce tumor. The cell count was adjusted to 1×10^6 cells/mL, and the diluted suspension was injected intraperitoneally. Tumor development was allowed for 7 days prior to the initiation of treatment.

2.3.2. Animals

A total of 24 male Swiss albino mice were placed in micro nylon boxes, maintained at a temperature of $25 \pm 2^\circ C$ and a 12-hour light–dark cycle. The mice were quarantined for 15 days prior to commencing the experiment. A healthy, standard laboratory diet was provided with water *ad libitum*. The animal house was maintained in a hygienic environment. The study was approved by the Institutional Animal Ethical Committee (IAEC approval No. IAEC/MEENASHREE BALAKRISHNAN/Ph.D/KMCP/92/2020).

The animals were divided into four groups ($n = 6$ per group):

- Group 1 (G1, normal control): received only a standard diet and water *ad libitum* without any intervention.

- Group 2 (G2, cancer control): injected intraperitoneally with DLA cells and left untreated to monitor progression of tumor.
- Group 3 (G3, positive control): treated with 5-fluorouracil (20 mg/kg, i.p.), a standard chemotherapeutic agent.
- Group 4 (G4, Siddha treatment I): administered TNC (100 mg/kg, by mouth/orally [p.o.]).
Treatment of tumor continued for 14 consecutive days, starting 24 hours after inoculation of tumor.

2.3.3. Hematological and biochemical analyses

At the end of the study period, the Swiss albino mice from all groups were euthanized, and blood samples were collected through retro-orbital plexus puncture for further examination. Hematological indices, such as hemoglobin (Hb) concentration, red blood cell (RBC) count, total white blood cell (WBC) count, packed cell volume, and platelet count, were quantified using a COBAS MICROS OT 18 cell analyzer (Roche Diagnostics, Basel, Switzerland). Serum was separated by centrifugation and subjected to biochemical and lipid profile assessment. Other significant parameters, such as alanine aminotransferase (ALT), total cholesterol (TC), aspartate aminotransferase (AST), triglycerides (TG), and alkaline phosphatase (ALP), were measured.

2.3.4. Life span measurement

In order to evaluate the efficacy of Siddha formulations in prolonging survival, the percentage ILS (%ILS) was determined. The average life span of animals was recorded for each group, and the %ILS was calculated for treatment groups, compared to the cancer control group:

$$\%ILS = \frac{\text{Life span of treated group}}{\text{Life span of control group}} - 1 \times 100$$

2.3.5. Body weight measurement

All mice were weighed at the start of the study and monitored at regular intervals until day 15. The final body weight of each group was recorded, and the average increase or decrease in body weight was determined to assess progression of tumor and response to treatment.

2.3.6. Cancer cell count

To assess tumor burden, 100 μ L of peritoneal fluid was collected from each animal using a sterile syringe. The collected sample was diluted with 0.8 mL of sterile phosphate buffer solution (PBS) to ensure proper cell dispersion. For viability assessment, 0.1 mL of trypan blue (0.1 mg/mL) was added to the sample. With the help of a hemocytometer, the number of viable cells was counted using the microscope:

$$\text{Cell count} = \frac{\text{Number of cells} \times \text{dilution factor}}{\text{Area} \times \text{thickness of liquid film}}$$

2.3.7. Histopathology procedure

The excised kidney and liver tissue samples were fixed in 10% neutral buffered formalin to preserve cellular morphology and structural integrity. Following fixation, the tissues were dehydrated through a graded ethanol series to remove water content, and subsequently cleared with a suitable clearing agent to enhance paraffin infiltration. The tissues were embedded in paraffin blocks by wax impregnation, which facilitated thin sectioning. With the help of a rotary microtome, thin sections of 5- μ m thickness were cut and stained using hematoxylin and eosin (H&E) for microscopic examination.

2.3.8. Statistical methods

All results were expressed as mean \pm standard deviation (SD). Statistical comparisons between groups were performed using one-way analysis of variance (ANOVA), followed by appropriate *post hoc* tests where applicable; $p < 0.05$ was considered statistically significant.

3. RESULTS

3.1. Antioxidant Activity Using DPPH Radical Scavenging Assay

The antioxidant activity of TNC was determined using the DPPH assay, and the percentage inhibition values are recorded in Table 2. The antioxidant effect of TNC increased with higher concentrations (0.5, 1, 2, 4, 8, and 16 mg/mL). The lowest concentration (0.5 mg/mL) showed 9.67% inhibition, which increased to 73.55% inhibition at a concentration of 16 mg/mL. The IC₅₀ value for TNC was 7 mg/mL. Ascorbic acid, which was used as a standard, also showed dose-dependent activity, ranging from 46.13% at 10 μ g/mL to 89.64% at 60 μ g/mL, with an IC₅₀ value of 17.9 μ g/mL. The dose–response curves for both TNC and ascorbic acid are shown in Figures 2 and 3, respectively.

3.2. Cytotoxicity evaluation using MTT assay

The cytotoxic activity of TNC was assessed on MCF-7, A549, HEP-G2, HeLa, and Vero cell lines using the MTT assay. TNC exhibited the highest cytotoxic effect against MCF-7 cells (IC₅₀ = 14.39 mg/mL), followed by A549 (IC₅₀ = 26.70 mg/mL) and HEP-G2 (IC₅₀ = 32.30 mg/mL). HeLa cells showed moderate sensitivity (IC₅₀ = 86.08 mg/mL),

whereas Vero cells demonstrated the least sensitivity (IC_{50} = 183.35 mg/mL). These IC_{50} values of TNC against the tested cell lines are given in Table 3. The results indicate that TNC possesses selective cytotoxic potential, particularly against breast (MCF-7) and lung (A549) cancer cell lines, while showing relatively lower toxicity on normal Vero cells.

Table 2
Percentage DPPH radical scavenging activity of *Thirunethira Chooranam* and ascorbic acid at different concentrations.

Concentration of ascorbic acid (µg/mL)	% Activity (ascorbic acid)	Concentration of TNC (mg/mL)	% Activity (TNC)
10	46.13	0.5	9.67
20	53.54	1	21.05
30	63.51	2	29.27
40	71.92	4	39.31
50	84.61	8	52.5
60	89.64	16	73.55

Notes: IC_{50} values: ascorbic acid = 17.9 µg/mL; TNC = 7 mg/mL (mean ± SD, n = 3).
TNC: *Thirunethira Chooranam*.

3.3. Anticancer activity in DLA-induced tumor-bearing Swiss albino mice

3.3.1. Hematological parameters

The effect of TNC on hematological indices in tumor-induced mice is shown in Table 4. In the cancer control group (G2), a significant increase in total white blood corpuscles (WBC) count ($13.52 \pm 2.60 \times 10^3$ cells/mL) and a reduction in red blood cell (RBC) count ($2.28 \pm 0.44 \times 10^6$ /mL), hemoglobin content (7.45 ± 0.92 g/dL), and platelet count ($1.72 \pm$

Table 3
 IC_{50} values of *Thirunethira Chooranam* against human cancer and normal cell lines determined by MTT assay.

Sl No.	Cell line	IC_{50} of TNC (mg/mL)
1.	MCF-7	14.39
2.	A549	26.70
3.	HEP-G2	32.30
4.	HeLa	86.08
5.	Vero	183.36

Note: TNC: *Thirunethira Chooranam*.




Figure 2. Dose-dependent DPPH radical scavenging activity of *Thirunethira Chooranam*.




Figure 3. Dose-dependent DPPH radical scavenging activity of ascorbic acid (as a standard).

Table 4Effect of *Thirunethira Chooranam* on hematological parameters in DLA-induced tumor-bearing mice.

Group	Total WBC ($\times 10^3$ cells/mL)	RBC ($\times 10^6$ /mL)	Hb (g/dL)	PCV (%)	Platelets (lakhs/mm ³)
G1 (normal control)	9.70 \pm 1.54	4.25 \pm 0.90	12.88 \pm 1.36	14.44 \pm 2.50	3.45 \pm 0.88
G2 (cancer control)	13.52 \pm 2.60*	2.28 \pm 0.44*	7.45 \pm 0.92*	30.80 \pm 3.30*	1.72 \pm 0.62*
G3 (Fluorouracil, 20 mg/kg, i.p.)	11.65 \pm 1.80**	4.04 \pm 0.84**	11.35 \pm 1.30**	18.35 \pm 1.55**	2.80 \pm 0.80**
G4 (TNC, 100 mg/kg, p.o.)	12.05 \pm 1.84**	3.63 \pm 0.74**	10.75 \pm 1.19**	22.05 \pm 1.68**	2.64 \pm 0.73**

Notes: TNC: *Thirunethira Chooranam*; i.p.: intraperitoneal; p.o.: by mouth/orally; PCV: packed cell volume.Values are mean \pm standard error of mean (SEM) (n = 6). *Significantly different from normal control (G1) at p < 0.01. **Significantly different from cancer control (G2) at p < 0.01.

0.62 lakhs/mm³) were observed, compared to the normal control (G1, p < 0.01). Treatment with TNC (G4) significantly restored these parameters to normal levels, compared with the positive control (G3), indicating a protective effect against tumor-induced hematological alterations.

3.3.2. Biochemical parameters

The biochemical profile of experimental groups is shown in Table 5. The cancer control group (G2) exhibited elevated serum cholesterol (143.90 \pm 4.68 mg/dL), triglycerides (224.38 \pm 4.82 mg/dL), AST (85.40 \pm 2.70 U/L), ALT (60.50 \pm 2.70 U/L), and ALP (242.50 \pm 4.45 U/L), compared to the normal control (G1, p < 0.01). Administration of TNC (G4) significantly reduced these elevated markers, with values comparable to the positive control (G3), thereby suggesting hepatoprotective and metabolic regulatory effects of TNC.

3.3.3. Life span and body weight

As shown in Table 6, the cancer control group (G2) demonstrated a reduced life span and a marked increase in body weight (7.80 \pm 0.95 g) because of tumor progression. In contrast, TNC-treated mice (G4) exhibited a substantial improvement in survival (80% ILS) and a significant reduction in body weight (4.14 \pm 0.80 g, p < 0.01 vs. G2). These findings were consistent with the effects of the positive

control group (G3), which recorded the highest survival benefit (93% ILS).

3.3.4. Cancer cell count

Tumor burden, measured by peritoneal cancer cell count, was markedly elevated in the cancer control group (G2; 2.68 \pm 0.42 \times 10⁶/mL), compared to the normal control (G1). TNC-treated mice (G4; 1.77 \pm 0.22 \times 10⁶/mL) showed significantly reduced viable cancer cell numbers (p < 0.01), compared to cancer control group (G2), and the effect was comparable to that of the positive control group (G3; 1.43 \pm 0.22 \times 10⁶/mL). These findings are presented in Table 6 and it confirms the *in vivo* cytotoxic efficacy of TNC against tumor-induced cancer cells.

3.3.5. Histopathology

Histopathological examination of liver sections presented in Figure 4 revealed distinct changes among experimental groups. Normal control (G1) animals displayed preserved hepatic architecture with well-arranged hepatocytes and normal sinusoids. Cancer control (G2) animals exhibited severe pathological changes, including hepatic congestion at sinusoids and portal vessels, diffused necrosis, and increased Kupffer cell proliferation and mononuclear infiltration, indicating significant liver damage. Positive control (G3) animals

Table 5Effect of *Thirunethira Chooranam* on biochemical parameters in DLA-induced tumor-bearing mice.

Group	Cholesterol (mg/dL)	TGL (mg/dL)	AST (U/L)	ALT (U/L)	ALP (U/L)
G1 (normal control)	118.32 \pm 3.75	142.60 \pm 2.35	42.75 \pm 1.40	35.60 \pm 1.64	130.45 \pm 2.20
G2 (cancer control)	143.90 \pm 4.68*	224.38 \pm 4.82*	85.40 \pm 2.70*	60.50 \pm 2.70*	242.50 \pm 4.45*
G3 (fluorouracil, 20 mg/kg, i.p.)	122.75 \pm 3.86**	166.75 \pm 2.54**	55.20 \pm 1.65**	40.45 \pm 1.60**	158.48 \pm 2.43**
G4 (TNC, 100 mg/kg, p.o.)	123.62 \pm 4.02**	169.20 \pm 2.80**	63.15 \pm 1.76**	47.85 \pm 2.10**	188.20 \pm 2.84**

Notes: TNC: *Thirunethira Chooranam*; i.p.: intraperitoneal; p.o.: by mouth/orally; TGL: triglycerides; AST: aspartate aminotransferase; ALT: alanine aminotransferase; ALP: alkaline phosphatase.Values are mean \pm standard error of mean (SEM) (n = 6). *Significantly different from normal control (G1) at p < 0.01. **Significantly different from cancer control (G2) at p < 0.01.

Table 6Effect of *Thirunethira Chooranam* on life span, body weight, and cancer cell count in DLA-induced tumor-bearing mice.

Group	% ILS (life span)	Increase in body weight (g)	Cancer cell count ($\times 10^6/\text{mL}$)
G1 (normal control)	NA	2.30 \pm 0.60	–
G2 (cancer control)	NA	7.80 \pm 0.95*	2.68 \pm 0.42*
G3 (fluorouracil, 20 mg/kg, i.p.)	93%	3.84 \pm 0.62**	1.43 \pm 0.22**
G4 (TNC, 100 mg/kg, p.o.)	80%	4.14 \pm 0.80**	1.77 \pm 0.22**


Notes. TNC: *Thirunethira Chooranam*; ILS: increase in life span; i.p.: intraperitoneal; p.o.: by mouth/orally.Values are mean \pm standard error of mean (SEM) (n = 6). *Significantly different from normal control (G1) at $p < 0.01$. **Significantly different from cancer control (G2) at $p < 0.01$.

Figure 4. Histopathological sections of liver tissues in different experimental groups. (A) G1: normal control showing intact hepatocytes arranged in cords with normal sinusoids and central vein. (B) G2: cancer control exhibiting hepatic congestion, micro-steatosis, increased Kupffer cell proliferation, hepatocyte necrosis, and mononuclear infiltration. (C) G3: positive control (fluorouracil, 20 mg/kg, i.p.) showing mild hepatic congestion with reduced necrosis and minimal mononuclear infiltration. (D) G4: *Thirunethira Chooranam*-treated group (100 mg/kg, p.o.) showing moderate hepatic congestion with reduced pericenter globular microsteatosis, less Kupffer cell proliferation, and partial restoration of hepatic architecture.

treated with fluorouracil showed partial restoration of hepatic tissue with only mild congestion at sinusoids and portal vessels, and no Kupffer cell proliferation and reduced necrosis. Animals treated with TNC (G4) demonstrated moderate improvement, with reduced hepatocyte necrosis, less Kupffer cell proliferation, and an overall improvement in tissue architecture, compared to the tumor control group.

Histopathological examination of kidney sections presented in Figure 5 revealed that normal control (G1) animals

displayed preserved renal histoarchitecture with well-arranged glomeruli and proximal convoluted tubules. Cancer control (G2) animals exhibited severe renal damage, characterized by glomerular disruption, folding of Bowman's capsules, necrotic changes, and degeneration of the renal parenchyma. Positive control (G3) animals treated with fluorouracil showed largely intact glomerular structures, accompanied by only mild inflammatory changes, indicating partial protection. Animals treated with TNC (G4) demonstrated moderate




Figure 5. Histopathological sections of kidney tissue in different experimental groups. (A) G1: normal control showing intact glomeruli (GM) and proximal convoluted tubules with preserved renal architecture. (B) G2: cancer control exhibiting glomerular damage, folding of Bowman's capsules, necrotic changes, and disrupted renal parenchyma. (C) G3: positive control (fluorouracil, 20 mg/kg, i.p.) showing intact glomeruli with only mild inflammatory changes. (D) G4: *Thirunethira Chooranam*-treated group (100 mg/kg, p.o.) showing mild glomerular inflammation with lymphocyte infiltration and minimal degenerative changes in the renal tubules, indicating partial restoration of renal structure.

improvement, evidenced by mild glomerular inflammation, reduced tubular degeneration, and fewer necrotic changes, suggesting partial restoration of renal tissue architecture, compared to the tumor control group.

4. DISCUSSION

The present study provides evidence for the antioxidant activity of TNC, a Siddha polyherbal formulation. The DPPH assay confirmed its free radical scavenging effect ($IC_{50} = 7$ mg/mL), supporting its potential to mitigate oxidative stress, a major contributor to deoxyribonucleic acid (DNA) damage and carcinogenesis. The activity can be attributed to its individual ingredients: *Vellarugu* exhibited potent antioxidant effects in 2,2'-Azino-bis 3-ethylbenzothiazoline-6-sulfonic acid (ABTS), deoxyribose, nitric oxide (NO), and lipid peroxidation assays (Vaijanathappa et al., 2008); *vellarugu* further demonstrated significant phenolic and flavonoid content with DPPH, ABTS, and ferric reducing antioxidant power (FRAP) activity (Sanmugarajah et al., 2018); *shivanarvembu* showed strong radical scavenging in DPPH, ABTS, hydroxyl radical,

and NO assays (Philips et al., 2010), with indigocarpan further confirming activity in ABTS, DPPH, thiobarbituric acid reactive substances (TBARS), and reactive oxygen species (ROS) assays (Mahajan et al., 2016); and *sangan kuppi* exerted free radical scavenging, reducing power, and lipid peroxidation inhibition because of its phenolic and flavonoid constituents (Gurudeeban et al., 2010; Khan et al., 2013). These findings collectively corroborate the antioxidant efficacy of TNC.

Thirunethira Chooranam demonstrated selective cytotoxicity *in vitro*, with strongest effects against MCF-7 breast cancer cells, followed by A549 and HEP-G2, while sparing normal Vero cells. This suggests that its phytochemical constituents may preferentially target malignant pathways. The *in vivo* DLA model further reinforced these observations, as TNC treatment restored altered hematological indices, reduced viable tumor cell counts, prolonged survival, and protected hepatic and renal tissues. Each component of the formulation potentially contributes via distinct pathways. *Vellarugu* contains swertiamarin, lupeol, and other flavonoids and terpenoids, which are shown to modulate apoptosis and inhibit tumor growth. The ethanolic extract suppressed 7,12-dimethylbenz[a]anthracene (DMBA)-induced tumor

formation and restored antioxidant enzyme levels *in vivo*, indicating chemopreventive action via reduction of oxidative stress and lipid peroxidation (Rajasekaran et al., 2015). The isolated *swertiamarin* from *vellaragu* exhibits anticancer activity *in vitro* by inducing apoptosis and cell cycle arrest, possibly via ROS generation and mitochondrial pathway modulation (Muhamad Fadzil et al., 2021). In addition, methanolic extract of *vellaragu* has immunomodulatory effects, enhancing both humoral and cellular immune functions while reducing pro-inflammatory cytokines (Saravanan et al., 2012).

Shivanarvembu has documented anti-tumor effects in animal models, as using its aqueous extract reduced fibrosarcoma progression induced by 20-MCA in rats (Sivagnanam et al., 2012). The ethanol extract demonstrated significant antitumor activity *in vivo* against Ehrlich ascites carcinoma in mice. It also exhibited cytotoxicity *in vitro*, supporting its role in cancer inhibition (Raj Kapoor et al., 2004). *Sangan kuppi* contributes by inhibiting cancer cell motility, invasion, and adhesion by using the methanolic leaf extract, which suppressed migration, invasion, and adhesion of A549 lung cancer cells at sub-cytotoxic doses (Tayeh et al., 2020).

The combined evidence indicates that TNC's anticancer effects are potentially mediated through multiple, complementary mechanisms, including ROS-mediated mitochondrial apoptosis, antioxidant defense, and immunomodulation. Such polyherbal synergy is consistent with the Siddha principle of balancing cytotoxicity with systemic protection. Future studies focusing on *in vitro* molecular pathway validation and clinical trials for specific cancer are necessary to strengthen its translational potential.

5. CONCLUSION

Thirunethira Chooranam demonstrated significant *in vitro* antioxidant and cytotoxic, and *in vivo* anticancer activities. These findings support its traditional use in the Siddha medicine and suggest its potential in cancer management. Further studies are needed to confirm its efficacy and safety through detailed molecular and clinical investigations.

AUTHOR CONTRIBUTIONS

Sudha Revathy Senthilkumar: research concept and design, collection and/or assembly of data, critical revision of the article, and final approval of the article; Meenashree Balakrishnan: collection and/or assembly of data; Kayalvizhi Duraisami: data analysis and interpretation, and writing of the article; and Deepa Natarajan: Research concept and design.

CONFLICT OF INTEREST

The authors declared no conflict of interest regarding the publication of this paper.

ACKNOWLEDGEMENT

The authors thanked the management of Asthagiri Herbal Research Foundation, Chennai for providing laboratory facilities and support to carry out this research work.

FUNDING

This study was supported by the Ministry of AYUSH, Government of India, under the Extramural Research (EMR) scheme.

ORCID

Deepa Natarajan	0000-0003-1446-4028
Sudha Revathy Senthilkumar	0009-0003-7496-0792
Meenashree Balakrishnan	0009-0002-4183-4880
Kayalvizhi Duraisamy	0009-0005-6021-7994

REFERENCES

- Bray, F., Laversanne, M., Sung, H., Ferlay, J., Siegel, R.L., Soerjomataram, I., Jemal, A., 2024. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA: A Cancer Journal for Clinicians. 74(3), 229–263. <https://doi.org/10.3322/caac.21834>
- Cragg, G.M., Kingston, D.G.I., Newman, D.J., 2012. Anticancer Agents from Natural Products, 2nd ed. CRC Press, Boca Raton, FL.
- Cragg, G.M., Pezzuto, J.M., 2016. Natural products as a vital source for the discovery of cancer chemotherapeutic and chemopreventive agents. Medical Principles and Practice. 25(Suppl 2), 41–59. <https://doi.org/10.1159/000443404>
- Gurudeeban, S., Satyavani, K., Ramanathan, T., Umamaheswari, G., Shanmugapriya, R., 2010. Antioxidant and radical scavenging effect of *Clerodendrum inerme* (L.). World Journal of Fish and Marine Sciences. 2(1), 66–69.
- International Agency for Research on Cancer (IARC), 2022. India Fact Sheet 2022 (GLOBOCAN Estimates). IARC, Lyon, France. <https://gco.iarc.who.int/media/globocan/factsheets/populations/356-india-fact-sheet.pdf> (Accessed: 15 November 2025).
- Kanagarathinam, D.V., Lourdasamy, J.B., 2023. Rise of Siddha medicine: causes and constructions in the Madras Presidency (1920–1930s). Medical History. 67(1), 42–56. <https://doi.org/10.1017/mdh.2023.10>
- Khan, S.A., Rasool, N., Riaz, M., Nadeem, R., Rashid, U., Rizwan, K., Zubair, M., Bukhari, I.H., Gulzar, T., 2013. Evaluation of

- antioxidant and cytotoxicity studies of *Clerodendrum inerme*. Asian Journal of Chemistry. 25(11), 6431–6436.
- Koleva, I.I., van Beek, T.A., Linssen, J.P., de Groot, A. and Evstatieva, L.N., 2002. Screening of plant extracts for antioxidant activity: a comparative study on three testing methods. Phytochemical Analysis. 13(1), 8–17. <https://doi.org/10.1002/pca.611>
- Li, M., Ka, D., Chen, Q., 2024. Disparities in availability of new cancer drugs worldwide: 1990–2022. BMJ Global Health. 9(9), e015700. <https://doi.org/10.1136/bmjgh-2024-015700>
- Mahajan, P., Gnanaraj, R., Jachak, S.M., Bharate, S.B., Chaudhuri, B., 2016. Antioxidant and antiproliferative activity of indigocarpan, a pterocarpan from *Indigofera aspalathoides*. Journal of Pharmacy and Pharmacology. 68(12), 1581–1590. <https://doi.org/10.1111/jphp.12609>
- Mohanty, S.K., Wadasadawala, T., Sen, S., et al., 2024. Catastrophic health expenditure and distress financing of breast cancer treatment in India: evidence from a longitudinal cohort study. International Journal for Equity in Health. 23, 145. <https://doi.org/10.1186/s12939-024-02215-2>
- Mosmann, T., 1983. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. Journal of Immunological Methods. 65(1–2), 55–63. [https://doi.org/10.1016/0022-1759\(83\)90303-4](https://doi.org/10.1016/0022-1759(83)90303-4)
- Muhamad Fadzil, N.S., Sekar, M., Gan, S.H., Bonam, S.R., Wu, Y.S., Vaijanathappa, J., Ravi, S., Lum, P.T., Dhadde, S.B., 2021. Chemistry, pharmacology and therapeutic potential of swertiamarin – a promising natural lead for new drug discovery and development. Drug Design, Development and Therapy. 15, 2721–2746. <https://doi.org/10.2147/DDDT.S299753>
- Newman, D.J., Cragg, G.M., 2020. Natural products as sources of new drugs over the nearly four decades from 01/1981 to 09/2019. Journal of Natural Products. 83(3), 770–803. <https://doi.org/10.1021/acs.jnatprod.9b01285>
- Philips, A., Philip, S., Arul, V., Padmakeerthiga, B., Renju, V., Santha, S., Sethupathy, S., 2010. Free radical scavenging activity of leaf extracts of *Indigofera aspalathoides* – an *in vitro* analysis. Journal of Pharmaceutical Sciences and Research. 2(6), 322–328.
- Rajasekaran, D., Manoharan, S., Prabhakar, M., Manimaran, A., 2015. *Enicostemma littorale* prevents tumor formation in 7,12-dimethylbenz(a)anthracene-induced hamster buccal pouch carcinogenesis. Human & Experimental Toxicology. 34(9), 911–921. <https://doi.org/10.1177/0960327114562033>
- Rajkapoor, B., Jayakar, B., Muruges, N., 2004. Antitumor activity of *Indigofera aspalathoides* on Ehrlich ascites carcinoma in mice. Indian Journal of Pharmacology. 36(1), 38–40.
- Sanmugarajah, V., Thabrew, I., Sivapalan, S.R., 2018. *In vitro* antioxidant activities of aqueous and methanol extracts of *Enicostemma littorale* Blume. RA Journal of Applied Research. 4(4), 1649–1655. <https://doi.org/10.31142/rajar/v4i4.14>
- Saravanan, S., Prakash Babu, N., Pandikumar, P., Karunai Raj, M., Gabriel Paulraj, M., Ignacimuthu, S., 2012. Immunomodulatory potential of *Enicostema axillare* (Lam.) A. Raynal, a traditional medicinal plant. Journal of Ethnopharmacology. 140(2), 239–246. <https://doi.org/10.1016/j.jep.2012.01.010>
- Sivagnanam, S.K., Rao, M.R., Balasubramanian, M.P., 2012. Chemotherapeutic efficacy of *Indigofera aspalathoides* on 20-methylcholanthrene-induced fibrosarcoma in rats. ISRN Pharmacology. 2012, 134356. <https://doi.org/10.5402/2012/134356>
- Subbarayappa, B.V., 1997. Siddha medicine: an overview. The Lancet. 350(9094), 1841–1844. [https://doi.org/10.1016/s0140-6736\(97\)04223-2](https://doi.org/10.1016/s0140-6736(97)04223-2)
- Tang, M-B., Kuo, W-Y., Kung, P-T, Tsai, W-C., 2024. The survival and cost-effectiveness analysis of adjunctive Chinese medicine therapy for patients with non-small cell lung cancer: a nationwide cohort study in Taiwan. Frontiers in Pharmacology. 15, 1378483. <https://doi.org/10.3389/fphar.2024.1378483>
- Tayeh, M., Hiransai, P., Kommen, H., Watanapokasin, R., 2020. Anti-migration and anti-invasion abilities of methanolic leaves extract of *Clerodendrum inerme* on lung cancer cells. Pharmacognosy Journal. 12(5), 1024–1031.
- Thavamani, B.S., Mathew, M., Palaniswamy, D.S., 2014. Anticancer activity of *Cocculus hirsutus* against Dalton's lymphoma ascites (DLA) cells in mice. Pharmaceutical Biology, 52(7), 867–872. <https://doi.org/10.3109/13880209.2013.871642>
- Vaijanathappa, J., Badami, S., Bhojraj, S., 2008. *In vitro* antioxidant activity of *Enicostemma axillare*. Journal of Health Science. 54(5), 524–528. <https://doi.org/10.1248/jhs.54.524>
- Wagner, H., Ulrich-Merzenich, G., 2009. Synergy research: approaching a new generation of phytopharmaceuticals. Phytomedicine. 16(2–3), 97–110. <https://doi.org/10.1016/j.phymed.2008.12.018>
- World Health Organization (WHO), 2024. Global Cancer Burden Growing, amidst Mounting Need for Services. WHO, Geneva, Switzerland. <https://www.who.int/news/item/01-02-2024-global-cancer-burden-growing--amidst-mounting-need-for-services> (Accessed: 15 November 2025).