

Original Article

View Article online



Received 27 October 2025

Revised 30 October 2025

Accepted 28 January 2026

Available Online 13 April 2026

Edited by Kannan RR Rengasamy

KEYWORDS:

Anti-inflammatory

Moringa-based cytotoxic

Anti-methanogen

Anti-mastitis

ADME traits

<https://doi.org/10.53365/nrfhh/217470>

eISSN: 2583-1194

Copyright © 2026 Visagaa Publishing House

Eco-functional phytogetic compounds: in silico and biological assessment toward host health and environmental microbial modulation

Srisan Phupaboon¹, Farah J. Hashim², Pongpat Kiatprasert³, Nattawadee Kanpipit⁴, Sakornchon Mattariganont⁴, Sukrita Punyauppa-Path³

¹Department of Biology, Faculty of Science, Maharakham University, Maharakham, Thailand

²Department of Biology, College of Science, Baghdad University, Baghdad, Iraq

³Department of Mathematics and Science, Faculty of Agriculture and Technology, Rajamangala University of Technology Isan Surin Campus, Surin, Thailand

⁴Faculty of Pharmaceutical Sciences, Khon Kaen University, Khon Kaen, Thailand


ABSTRACT: This study established the bioavailability of cytotoxic compounds, including polyphenolics, flavonoids, and antioxidants, derived from *Moringa oleifera* leaf extract through in silico analysis, focusing on their antimethanogen, antimicrobial mastitis, and anti-inflammatory properties to predict active-site protein targets. The analysis involved 40 compounds utilizing LC-QTOF/MS, molecular docking, and ADME property assessments. These compounds were evaluated against the methyl-coenzyme M reductase (MRC) receptor (PDB: 1MRO), Mycoplasma bovis protein (PDB: 7E2P), and interleukin-37 receptor (PDB: 5HN1) using the SwissDock server via the AutoDock Vina platform. The molecular docking results identified three optimal ligands: apigenin (−7.23 kcal/mol), tirandamycin-A (−8.70 kcal/mol), and 2''-O-acetylrutin (−7.94 kcal/mol), each demonstrating significant binding affinity with different target proteins. In conclusion, apigenin and tirandamycin-A compounds demonstrate significant potential for the development of food-feed additives or pharmaceuticals to catalyze host health and address mastitis-causing pathogens in mammalian glands. This potential aligns with Lipinski's rule of five and encompasses ADME properties, including physicochemical, pharmacokinetic, and drug-likeness characteristics.

* Corresponding author.

E-mail address: phira.bo@rmuti.ac.th (Sukrita Punyauppa-Path)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

GRAPHICAL ABSTRACT



1. INTRODUCTION

The demand for animal protein, particularly meat and milk products from ruminant production, is projected to rise in accordance with the growing global population over the next 25 years, from the current 6.9 billion to more than 9.7 billion in 2050 (FAO, 2009; Wanapat et al., 2024a). The influence of this message on public discourse regarding hunger and malnutrition has been significant and continues to

be so. The FAO's emphasis on enhancing livestock production significantly contributes to global greenhouse gas (GHG) emissions, primarily methane (CH_4). Enteric fermentation, the source of these emissions, accounts for 90% of global GHG emissions from microbial methanogenesis. Ruminal methane production impacts the environment and represents an energy loss for animals, reducing growth rate and milk quality (Wanapat et al., 2024a). For ruminal methanogenesis process, methanogenic archaea including bacteria, protozoa,

and anaerobic fungi convert H₂ and CO₂ through the key enzyme of methyl-coenzyme M reductase (MRC) for CH₄ production (Ginovska et al., 2025; Patra & Saxena, 2010). Subsequently, animal sickness caused by pathogen infection such as *Mycoplasma* *bovis*-causing severe mastitis disease and cytokine inflammatory activation by interleukin-37 would affect animal productivity and their health (Ellisdon et al., 2017; Gelgie et al., 2024).

Various methane mitigation and risk-reduction strategies for animal sickness have been implemented globally, including the use of feed additives and the application of tropical/marine plant-based phytonutrients (Phupaboon et al., 2025a; Wanapat et al., 2024b,c). Recently, in vitro anti-methanogenesis and in vitro antimicrobial trials have shown that the phytochemicals (e.g., polyphenols, flavonoids, and antioxidant reagents) derived from plant extracts including garlic oil (Phupaboon et al., 2025b), lemongrass oil (Prachumchai et al., 2024), mangosteen peel (Sommai et al., 2025), rambutan/red dragon fruit peels (Suriyapha et al., 2025), banana leaf (Dagaew et al., 2024), hemp/marijuana leaf (Phupaboon et al., 2025c), and *Mitragyna* leaf (Matra et al., 2025). Especially, *Moringa oleifera* leaf provides several ethnopharmacological applications owing to their potent constituents, including carotenoids, polyphenols, flavonoids, essential amino acids, and phenolic acids, as well as its superior biological efficacy demonstrated in antioxidative reactions through in vitro and in vivo studies (Hashim et al., 2021a,b). Numerous innovative tools and in silico methodologies are being created to enhance the drug discovery process from plant-derived compounds, including molecular docking (e.g., AutoDock Vina, FAFDrugs3, PROTOX, and SwissDock) and pharmacokinetic design (e.g., SwissADME) (Arokiyaraj et al., 2019; Phupaboon et al., 2025a; Stefaniu & Pirvu, 2022). Additionally, the advantages and advancements in science facilitating docking analysis enable scientists to digitally examine chemical databases and select the most potent binders using various scoring algorithms. This study elucidates the binding interaction between two molecules, specifically a ligand (compounds) and a receptor protein of interest, along with friendly safety and pharmacokinetics (Punyauppa-Path et al., 2025).

According to a previous research study, Thai-Moringa leaf extract had the highest potency to decrease in vitro superoxide radical antioxidant capacity and in vivo neuroprotection against oxidative-stress-induced cytotoxicity in SHSY5Y neuroblastoma cells, as well as inhibition of tyrosinase activity (Hashim et al., 2021a,b), while the mode of mechanism was not reported in the effect of ruminal anti-methanogenesis (MRC receptor), anti-mastitis pathogen (*Mycoplasma* *bovis* of enolase protein), and anti-inflammatory (interleukin-37 protein). Therefore, this study hypothesized the effects of the

selected 40 Moringa-based cytotoxic components on in silico docking analysis with three target proteins.

2. MATERIALS AND METHODS

2.1. Plant collection and extraction

M. oleifera leaves were obtained from a local market in Khon Kaen, Thailand. Clean plant materials were dried in a hot-air oven at 65 °C and subsequently ground. The powdered *M. oleifera* leaves were subsequently mixed with 70% ethanol at a 1:4 (w/v) ratio and extracted using microwave extraction (Hashim et al., 2021a). The result was a *Moringa*-based ethanol extract powder for further use.

2.2. Phytochemicals analysis

The phytochemical compositions of *Moringa*-based ethanol extract were analyzed in terms of antioxidant activities, including total polyphenolic content (TPC) and total flavonoid content (TFC). TPC is reported as mg GAE/g DE, while TFC is expressed as mg QUE/g DE. Following the procedures outlined by Phupaboon et al. (2022), several plant extracts were subjected to the DPPH, ABTS, and FRAP assays, with results presented as mg TROE/g DE. The evaluation of antioxidant activity (expressed as mg TROE/g DE) was performed with three separate methodologies: the DPPH reagent measured at 517 nm, the ABTS reagent measured at 734 nm, and the completed FRAP reagent measured at 593 nm. All analyses were performed in duplicate.

2.3. Biological activity

A part of *Moringa*-based ethanol extract was tested for its antibacterial efficacy in vitro study because of its powerful antioxidant capability. The minimal inhibitory concentration (MIC) against pathogenic bacteria, including Gram(-): *E. coli* TISTR 073, *Ent. aerogenes* TISTR 1540, *S. typhimurium* TISTR 292, and Gram(+): *S. aureus* TISTR 029, *B. cereus* TISTR 678, was tested at concentrations of 31.25, 62.5, 125, 250, and 500 µg/mL using a two-fold dilution technique, as per the methodology given by Phupaboon et al. (2025a).

2.4. Moringa-based phytochemicals using LC-MS analysis

Based on earlier studies that showed different compounds in *Moringa*-based ethanol extract, it selected 40 compounds

for this study obtained from a liquid chromatograph-quadrupole time-of-flight mass spectrometer (LC-QTOF/MS) analysis, along with their names, molecular formulas, various retention times, mass spectra, and similarity indexes, which are listed in Table 1.

2.5. Preparation of ligand molecules and target protein structure

All 40 compounds (Table 1) were chosen as ligand molecules for preliminary screening and to predict the binding site of the energy-minimized target protein structure using

Table 1

Catalog of *Moringa*-based phytochemical compounds.

No.	Compound names	Molecular formula	Retention time (min)	[M+H] ⁺ Mass (m/z)	Similarity (%)
1	Isorhamnetin	C ₁₆ H ₁₂ O ₇	83.499	315.05	99.0
2	Quercetin	C ₁₅ H ₁₀ O ₇	48.551	303.04	46.0
3	Apigenin	C ₁₅ H ₁₀ O ₅	79.602	269.04	20.2
4	2 ^o -O-acetylrutin	C ₂₉ H ₃₂ O ₁₇	37.044	651.16	99.0
5	Hesperetin	C ₁₆ H ₁₄ O ₆	79.820	301.07	40.1
6	Hydroxytyrosol	C ₈ H ₁₀ O ₃	13.423	315.10	92.1
7	Kaempferol	C ₁₅ H ₁₀ O ₆	35.262	287.05	100.0
8	Hydroquinone	C ₆ H ₆ O ₂	14.005	110.04	99.0
9	Chlorogenic acid	C ₁₆ H ₁₈ O ₉	18.834	354.90	99.8
10	Ferulic acid	C ₁₀ H ₁₀ O ₄	27.61	194.06	61.0
11	Geniposide	C ₁₇ H ₂₄ O ₁₀	36.056	387.13	52.2
12	Inosine	C ₁₀ H ₁₂ N ₄ O ₅	8.017	268.08	90.5
13	Quinic acid	C ₇ H ₁₂ O ₆	18.884	192.06	99.0
14	Rutin	C ₂₇ H ₃₀ O ₁₆	35.299	610.15	100.0
15	L-threonic acid	C ₄ H ₈ O ₅	2.05	135.04	98.0
16	Tirandamycin A	C ₂₂ H ₂₇ NO ₇	18.508	417.18	99.0
17	Salidroside	C ₁₄ H ₂₀ O	20.98	300.12	67.2
18	Folic acid	C ₁₉ H ₁₉ N ₇ O ₆	64.017	441.14	46.1
19	Chlorogenic acid	C ₁₆ H ₁₈ O ₉	19.106	354.09	98.9
20	Piperic acid	C ₁₂ H ₁₀ O ₄	46.987	218.06	99.0
21	Lobeline	C ₂₂ H ₂₇ NO ₂	84.935	337.20	37.1
22	Granisetron	C ₁₈ H ₂₄ N ₄ O	30.764	312.20	99.0
23	Ramipril glucuronide	C ₂₉ H ₄₀ N ₂ O ₁₁	37.755	592.26	99.0
24	Dicrocin	C ₃₂ H ₄₄ O ₁₄	59.614	652.27	99.0
25	cilostazol	C ₂₀ H ₂₇ N ₅ O ₂	29.338	369.22	99.0
26	Aucubin	C ₁₅ H ₂₂ O ₉	16.334	346.13	76.2
27	3-Hydroxydecanoic acid	C ₁₀ H ₂₀ O ₃	26.432	188.14	99.1
28	Astragaln	C ₂₁ H ₂₀ O ₁₁	52.464	448.10	99.0
29	Protoporphyrinogen IX	C ₃₄ H ₄₀ N ₄ O ₄	45.690	568.31	99.0
30	Salicylanilide	C ₁₃ H ₁₁ NO ₂	64.054	213.08	97.9
31	Succinoadenosine	C ₁₄ H ₁₇ N ₅ O ₈	17.058	383.11	99.0
32	Quercitrin	C ₂₁ H ₂₀ O ₁₁	52.426	448.10	65.0
33	Isovitexin	C ₂₁ H ₂₀ O ₁₀	47.171	432.11	99.0
34	Dephospho Coenzyme A	C ₂₁ H ₃₅ N ₇ O ₁₃ P ₂ S	13.365	687.15	99.0
35	Nimodipine	C ₂₁ H ₂₆ N ₂ O ₇	41.972	418.17	98.9
36	Deoxynivalenol	C ₁₅ H ₂₀ O ₆	35.069	296.13	99.0
37	Esculin	C ₁₅ H ₁₆ O ₉	21.798	340.08	61.1
38	<i>Cis</i> -Cinnamic acid	C ₉ H ₈ O ₂	3.619	148.05	98.9
39	Penicillamine disulfide	C ₁₀ H ₂₀ N ₂ O ₄ S ₂	13.03	341.08	44.0
40	Gingerol	C ₁₇ H ₂₆ O ₄	88.746	294.18	99.0

the SwissDock server within the SwissDrugDesign platform with AutoDock Vina (<https://www.swissdock.ch/>). The methods and parameter setup were slightly modified from those described by Phupaboon et al. (2025) and Punyappa-Path et al. (2025). Ligand molecules were generated via the SwissParam website (<http://swissparam.ch/>) utilizing the simplified molecular input line entry system (SMILES) derived from the PubChem database.

Subsequently, three active site protein targets were selected: methyl-coenzyme M reductase (MRC) of *Methanothermobacter marburgensis* str. *marburg* (PDB: 1MRO), *Mycoplasmopsis bovis* (PDB: 7E2P), and interleukin-37 (PDB: 5HN1). These proteins were retrieved from the RCSB-PDB database, and their three-dimensional (3D) structures and grid box were generated using SWISS-MODEL (Figure 1), according to the procedure of Bugnon et al. (2024).

2.6. Molecular docking prediction

This study assessed the efficacy of 40 selected ligands derived from *Moringa*-based cytotoxic compounds in




Figure 1. The 3D structure of three active site protein targets: (A) *Methanothermobacter marburgensis* str. *marburg* (PDB: 1MRO), (B) *Mycoplasmopsis bovis* (PDB: 7E2P), and (C) interleukin-37 (PDB: 5HN1).

inhibiting methanogenesis, combating mastitis pathogens, and reducing inflammation. Each ligand molecule displays unique SMILES structures, as shown in Table 2. This study employed protein active site targets, as illustrated in Figure 1. Molecular docking predictions were performed by evaluating ligand interactions with the active sites of protein target structures using the SwissDock server (<https://www.swissdock.ch/>) through the AutoDock Vina platform (Bugnon et al., 2024; Punyappa-Path et al., 2025; Phupaboon et al., 2025a).

2.7. In silico ADME analysis

The in-silico evaluation involved assessing the physicochemical and pharmacokinetic properties of the potent ligands by inputting each SMILES structure into the SwissADME web tool (<http://www.swissadme.ch/>). This analysis examined physicochemical properties, including lipophilicity (Consensus Log P_{ow}), water solubility (Log S), pharmacokinetics, and druglikeness in relation to Lipinski's violations (Phupaboon et al., 2025a).

3. RESULTS

3.1. Phytochemical composition and biological activity

The phytochemical compositions of *Moringa*-based ethanol extract were 35.2 mg GAE/g DE, 9.2 mg QUE/g DE, as well as 7.0, 31.6, and 62.7 (mg TROX/g DE) in terms of TPC, TFC, DPPH, ABTS, and FRAP, respectively. Importantly, the antimicrobial activity of this extract presented the minimal inhibition concentration (MIC) at 31.25 $\mu\text{g/mL}$ against *E. coli*, *Ent. aerogenes*, and *S. typhimurium*, while the MIC of *B. cereus* was exhibited at 62.5 $\mu\text{g/mL}$, as shown in Table 2.

3.2. Molecular docking of ligand binding target protein

Table 3 illustrates the binding affinity values of the selected 40 ligands binding with three types of target protein, including methyl-coenzyme M reductase (MRC) on subunit alpha protein target of *Methanothermobacter marburgensis* str. *marburg* (PDB: 1MRO), *Mycoplasmopsis bovis* (PDB: 7E2P), and interleukin-37 (PDB: 5HN1) receptors. Interestingly, results showed maximum binding affinity and 3D structure of apigenin ligand with the MRC receptor (PDB: 1MRO) at -7.23 kcal/mol (as shown in Figure 2A,B), along with kaempferol at -7.10 kcal/mol. In terms of tirandamycin-A inhibited the protein of *Mycoplasmopsis bovis* (PDB: 7E2P) (3D structure in Figure 2C,D), which exhibited the highest affinity value

Table 2

Phytochemical composition and antimicrobial activity of *Moringa*-based ethanol extract

Phytochemical assay	Values	Antimicrobial activity	MIC ($\mu\text{g/mL}$)
TPC (mg GAE/g DE)	35.2 \pm 3.4	<i>E. coli</i> TISTR 073	31.25
TFC (mg QUE/g DE)	9.2 \pm 0.1	<i>Ent. aerogenes</i> TISTR 1540	31.25
DPPH (mg TROX/g DE)	7.0 \pm 0.3	<i>S. typhimurium</i> TISTR 292	31.25
ABTS (mg TROX/g DE)	31.6 \pm 0.4	<i>S. aureus</i> TISTR 029	31.25
FRAP (mg TROX/g DE)	62.7 \pm 0.4	<i>B. cereus</i> TISTR 678	62.5
		Norflloxacin	6.25

of -8.70 kcal/mol, while cilostazol, lobeline, folic acid, and granisetron were -8.39 , -8.23 , -8.11 , and -8.00 kcal/mol, respectively. In addition, the ligand of 2''-O-acetylrutin binding with the protein of interleukin-37 (PDB: 5HN1) receptor (3D structure in Figure 2E,F) showed the greater affinity value of -7.94 kcal/mol and -7.18 and -7.05 kcal/mol obtained from folic acid and dicrocic as ligand compounds.

3.3. Physicochemical and ADME properties of selected ligands

Table 4 presents the selection of three ligands—apigenin, tirandamycin-A, and 2''-O-acetylrutin—based on their highest binding affinity, for the evaluation of physicochemical and in silico ADME properties in accordance with Lipinski's rule of five. The physicochemical and ADME results are presented through various parameters, including molecular weight, the count of hydrogen bond acceptors and donors, the number of rotatable bonds, lipophilicity ($\text{Log } P_{ow}$), water solubility ($\text{Log } S$), pharmacokinetics such as gastrointestinal (GI) absorption, blood-brain barrier (BBB) permeability, P-glycoprotein (P-gp) substrate, and drug-likeness (Lipinski violations and bioavailability score). The results for apigenin and tirandamycin-A ligands were accepted, demonstrating excellent bioavailability in accordance with Lipinski's rule of five. In contrast, the 2''-O-acetylrutin ligand was not accepted due to a molecular weight exceeding 500g/mol, as well as having more than 10 hydrogen acceptors and more than 5 hydrogen donors.

4. DISCUSSION

As a plant with therapeutic potential and many health benefits, *Moringa* leaf has been the subject of numerous investigations into its biological activities, including anticancer, antipathogen, anti-inflammatory, and neuroprotective activities. The plant is rich in bioactive phenolic compounds

Table 3

Molecular docking of selected LC-MS compounds binding with methyl-coenzyme M reductase (MRC) on subunit alpha protein target of *Methanothermobacter marburgensis* str. *marburg* (PDB: 1MRO), *Mycoplasmopsis bovis* (PDB: 7E2P), and interleukin-37 (PDB: 5HN1).

No.	Ligand names	SMILES IDs	Calculated affinity (kcal/mol)		
			1MRO	7E2P	5HN1
1	Isorhamnetin	5281654	-6.59	-6.65	-6.19
2	Quercetin	5280343	-6.42	-7.36	6.66
3	Apigenin	5280443	-7.23	-7.70	-6.59
4	2''-O-Acetylrutin	9809744	-6.14	-3.07	-7.94
5	Hesperetin	72281	-4.37	-7.49	-6.21
6	Hydroxytyrosol	82755	-	-	-
7	Kaempferol	5280863	-7.10	-7.63	-7.637
8	Hydroquinone	785	-3.90	-5.70	-5.12
9	Chlorogenic acid	1794427	-5.13	-7.84	-6.04
10	Ferulic acid	445858	-4.70	-5.72	-5.20
11	Geniposide	107848	-5.68	-7.51	-6.49
12	Inosine	135398641	-5.80	-6.52	-5.79
13	Quinic acid	37439	-4.33	-5.37	-5.06
14	Rutin	5280805	-5.97	-3.44	-6.98
15	L-threonine acid	5460407	-3.38	-4.42	-3.91
16	Tirandamycin A	54717225	-6.35	-8.70	-6.93
17	Salidroside	159278	-5.41	-6.82	-5.68
18	Folic acid	135398658	-5.77	-8.11	-7.18
19	Chlorogenic acid	1794427	-5.04	-7.84	-6.26
20	Piperic acid	5370536	-4.97	-6.34	-5.61
21	Lobeline	101616	-5.26	-8.23	-6.45
22	Granisetron	5284566	-5.76	-8.00	-6.35
23	Ramipril glucuronide	76973982	-5.45	-6.45	-6.61
24	Dicrocic	25244294	-6.06	-3.56	-7.05
25	Cilostazol	2754	-5.23	-8.39	-5.74
26	Aucubin	91458	-5.99	-7.00	-6.27
27	3-Hydroxydecanoic acid	26612	-4.26	-4.86	-4.72
28	Astragalin	5282102	-6.80	-7.96	-6.48
29	Protoporphyrinogen IX	121893	-5.49	-4.79	-6.32
30	Salicylanilide	6872	-6.01	-7.04	-6.62
31	Succinoadenosine	20849086	-5.45	-7.49	-5.55
32	Quercitrin	5280459	-6.89	-7.20	-6.43
33	Isovitexin	162350	-5.78	-4.07	-6.68
34	Dephospho Co-A	444485	-1.33	-4.36	-5.95
35	Nimodipine	4497	-5.01	-6.40	-5.37
36	Deoxyvalenol	40024	-5.19	-5.72	-5.85
37	Esculin	5281417	-5.90	-7.72	-6.37
38	Cis-Cinnamic acid	5372954	-4.14	-5.63	-5.78
39	Penicillamine disulfide	258527	-4.35	-5.78	-5.36
40	Gingerol	442793	-4.57	-6.37	-5.39




Figure 2. The 3D molecular docking and the surface-protein binding site structures of apigenin with MRC (A, B), tirandamycin A with *Mycoplasma bovis*-enolase (C, D), and 2''-O-acetylrutin with interleukin-37 (E, F) protein targets.

and is widely acknowledged as an essential resource (Hashim et al., 2021a; Shalaby et al., 2022; Said-Al Ahl et al., 2017). The current results confirm the acceptance of apigenin and kaempferol, particularly apigenin, in accordance with Lipinski's rule. Khusro et al. (2020), who used in silico tools to target MCR receptors to predict the anti-methanogenic attributes of *Moringa*-based phytochemicals, concur with our findings. Among all the tested phytochemicals, 3,5-bis(1,1-dimethylethyl)-phenol, kaempferol, moringyne, niazimisin, and tetradecanoic acid. The review shows that tetradecanoic acid had the strongest link to the MCR receptor, with a binding energy of -142.98 KJ/mol, followed by niazimicin at -133.98 KJ/mol, kaempferol at -110.36 KJ/mol, 3,5-bis(1,1-dimethylethyl)-phenol at -93.72 KJ/mol, and moringyne at -92.62 KJ/mol. In addition, Dinakarkumar et al. (2021) investigated the application of an in silico molecular docking

approach to assess the antimethanogenic properties of phytochemicals derived from *Cymbopogon citratus*, *Origanum vulgare*, *Lavandula officinalis*, *Cinnamomum zeylanicum*, *Piper betle*, *Cuminum cyminum*, *Ocimum gratissimum*, *Salvia sclarea*, *Allium sativum*, *Rosmarinus officinalis*, and *Thymus vulgaris*. Under this review, 25 analyzed compounds were assessed for their interaction with MCR protein utilizing the AutoDock 4.0 software; specifically, five compounds are identified: rosmarinic acid (-10.71 kcal/mol), biotin (-9.38 kcal/mol), α -cadinol (-8.16 kcal/mol), and (3R,3aS,6R,6aR)-3-(2H-1,3-benzodioxol-4-yl)-6-(2H-1,3-benzodioxol-5-yl)-hexahydrofuro[3,4-c]furan-1-one (-12.21 kcal/mol), and 2,4,7,9-tetramethyl-5-decyn-4,7-diol (-9.02 kcal/mol) exhibited greater binding energy to the MCR protein.

The result of this study indicated that tirandamycin-A, cilostazol, lobeline, folic acid, and granisetron have a greater

Table 4

Physicochemical properties and SwissADME profiles of the selected compounds apigenin and kaempferol.

Categories	Ligands/compounds		
	Apigenin	Tirandamycin A	2''-O-Acetylrutin
Physicochemical properties			
Molecular weight (g/mol)	270.24	417.45	610.52
Num. H-bond acceptors	5	7	16
Num. H-bond donors	3	2	10
Num. rotatable bonds	1	4	7
Lipophilicity			
Log P _{ow} (Consensus/WLOGP)	2.11	1.13	-1.44
Water solubility			
Log S (ESOL)	-3.94	-	-3.36
Solubility class	Soluble	-	Soluble
Pharmacokinetics			
GI absorption	High	-	Low
BBB permeant	No	-	No
P-gp substrate	No	-	Yes
Drug-likeness			
Lipinski violation	Accepted	Accepted	No
Oral bioavailability	Good	Good	Good

binding affinity, and tirandamycin-A was accepted in pharmacokinetics and drug-likeness. In accordance with Zahran et al. (2020), it was reported that the selected compounds, specifically tirandamycin derived from marine natural products, exhibited potential inhibitory effects on SARS-CoV-2 targets through molecular docking studies against SARS-CoV-2 RNA-dependent RNA polymerase (RdRp) and methyltransferase (nsp16). Another research study by Adamu et al. (2024) conducted research examining the efficacy of *Moringa*-based phytochemicals as inhibitors of the Newcastle disease virus (NDV), utilizing molecular docking analysis to assess their anti-pathogenic properties. The review highlights that certain their bioactive compounds strongly attach to a part of the NDV fusion protein, which is crucial for the virus to infect host cells. The review indicated that specific phytochemicals, such as β -sitosterol, catechins, and kaempferol, exhibit significant binding affinity scores of -8.5 kcal/mol, -8.2 kcal/mol, and -7.9 kcal/mol. Additionally, Phupaboon et al. (2025a) demonstrated that the binding affinity of the allicin ligand from microencapsulated garlic oil extract docked with five PDB-protein targets of mastitis pathogens—*E. coli*, *S. aureus*, *Ent. aerogenes*, *S. typhimurium*, and *B. cereus*—were -14.99 , -14.71 , -16.12 , -11.14 , and

-18.89 kcal/mol of AC score by SwissDock server through the attracting cavity platform.

Moreover, this research found that the 2''-O-Acetylrutin ligand strongly reduces the levels of the pro-inflammatory cytokine interleukin-37, along with folic acid and dicrocin. In reviewing the literature, no information was found on how the 2''-O-Acetylrutin ligand interacts with interleukin-37 in terms of anti-inflammatory activity, using molecular docking and in silico ADME properties. Some publications related to medical plants, including *Centipeda minima*, *Ipomoea pes-caprae*, and *Dodonaea viscosa*, specified phytochemicals aliphatic acid, betulinic acid, malabaric acid, and hispidulin reduced proinflammatory cytokines and cyclooxygenase enzymes (COX-1, 2), with IC₅₀ values ranging from 11.5 to 46.9 μ M (Yeshe et al., 2022). Elsewhere, Ali et al. (2023) aimed to identify novel compounds; picrocrocin showed the highest binding affinity of -8.1 kcal/mol when docked against the COX-2 protein, which is associated with various diseases, including inflammation. Another research study by Punyappa-Path et al. (2025) related to alternative plant-based protein from quinoa natto with Thai herbs used to predict the efficacy of L-leucine ligand docking with different target proteins; type 1 angiotensin II receptor and parathyroid hormone-related protein of Caco-2 cell was -7.65 and -9.15 kcal/mol for AC score.

5. CONCLUSION

To the best of our understanding, *Moringa*-based polyphenols and/or cytotoxic compounds, which were selected from the 40 components, were potentially used to evaluate in silico, including molecular docking and ADME properties. The main discovery from the molecular docking study showed that the stronger compounds apigenin, tirandamycin-A, and 2''-O-acetylrutin could help reduce various risks associated with the MRC protein receptor involved in methane production, block the *Mycoplasma bovis* protein associated with bacteria that cause mastitis, and lower the inflammatory cytokine interleukin-37 linked to mastitis. Additionally, regarding the ADME profile, it was indicated that the potency of the compounds may be acceptable for developing anti-multifunctional drugs for livestock production, particularly in organic meat and milk products in the ruminant field.

FUNDING

This research project was financially supported by Mahasarakham University.

ACKNOWLEDGMENTS

The authors express gratitude to the Department of Biological, Faculty of Science, Mahasarakham University, Thailand, for their support of laboratory facilities and assistance.

CONFLICT OF INTEREST

The authors declare no conflict of interest with any organization about the topics discussed in this manuscript.

DATA AVAILABILITY

The datasets generated and analyzed during the present study are available within this article.

ETHICS STATEMENT

This research has no implications for animal ethics.

AUTHORSHIP CONTRIBUTION

S. Phupaboon and S. Punyauppa-Path: conceived and designed the experiment. FJ. Hashim, N. Kanpipit, and S. Mattariganont: performed the study, supported the materials, and conducted lab analyses. S. Phupaboon and P. Kialprasert: performed statistical analyses of experimental data, supervised and coordinated the experiments. S. Phupaboon, FJ. Hashim, and S. Punyauppa-Path: prepared the manuscript format, reviewed, and revised the manuscript. All authors critically revised the manuscript and approved the final version.

GENERATIVE AI STATEMENT

The authors declare that no Gen AI/DeepSeek was used in the writing/creation of this manuscript.

ORCID

Srisan Phupaboon	0000-0001-7719-4067
Farah J. Hashim	0009-0004-4057-038X
Sakornchon Mattariganont	0009-0003-5145-7449
Pongpat Kiatprasert	0009-0000-9669-8834

Nattawadee Kanpipit	0009-0003-2019-1254
Sukrita Punyauppa-Path	0009-0008-2271-5206

REFERENCES

- Adamu, U.A., Dhakar, R., Teli, P.K., Muhammad, S.S., & Gumel, I.A., 2024. In-silico evaluation of *Moringa oleifera* phytochemicals as inhibitors against Newcastle disease virus in birds. *International Journal for Multidisciplinary Research* 6(5), 1–33.
- Ali, A., Wani, A.B., Malla, B.A., Poyya, J., Dar, N.J., Ali, F., Ahmad, S.B., Rehman, M.U., Nadeem, A., 2023. Network pharmacology integrated molecular docking and dynamics to elucidate saffron compounds targeting human COX-2 protein. *Medicina* 59(12), 2058. <https://doi.org/10.3390/medicina59122058>
- Arokiyaraj, S., Stalin, A., Shin, H., 2019. Anti-methanogenic effect of rhubarb (*Rheum* spp.)—an in silico docking studies on methyl-coenzyme M reductase (MCR). *Saudi Journal of Biological Sciences* 26(7), 1458–1462. <https://doi.org/10.1016/j.sjbs.2019.06.008>
- Bugnon M, Röhrig UF, Goullieux M, Perez MAS, Daina A, Michielin O, Zoete V., 2024. SwissDock 2024: major enhancements for small-molecule docking with attracting cavities and AutoDock Vina. *Nucleic Acids Research* 52(W1), W324–W332. <https://doi.org/10.1093/nar/gkae300>
- Dagaew, G., Kunhareang, S., Phupaboon, S., Matra, M., Sommai, S., Suriyapha, C., Prachumchai, R., Wanapat, M., 2024. The use of microencapsulated banana flower powder pellet on in vitro ruminal fermentation, digestibility, microbial diversity, and methane production. *Tropical Animal Health and Production* 56(8), 263. <https://doi.org/10.1007/s11250-024-04101-z>
- Dinakarkumar, Y., Rajabathar, J.R., Arokiyaraj, S., Jeyaraj, I., Anjaneyulu, S.R., Sandeep, S., ... Wilson, L.D., 2021. Anti-methanogenic effect of phytochemicals on methyl-coenzyme M reductase—potential: in silico and molecular docking studies for environmental protection. *Micromachines* 12(11), 1425. <https://doi.org/10.3390/mi12111425>
- Ellisdon, A.M., Nold-Petry, C.A., D'Andrea, L., Cho, S.X., Lao, J.C., Rudloff, I., ... Nold, M.F., 2017. Homodimerization attenuates the anti-inflammatory activity of interleukin-37. *Science Immunology* 2(8), eaaj1548. <https://doi.org/10.1126/sciimmunol.aaj1548>
- FAO, 2009. Global agriculture towards 2050, high-level expert forum, how to feed the world 2050, Rome 12–13 October 2009. Food and Agriculture Organization of United Nations FAO.
- Gelgie, A.E., Schneider, P., Citti, C., Dordet-Frisoni, E., Gillespie, B.E., Almeida, R.A., ... Lysnyansky, I., 2024. *Mycoplasma bovis* 5'-nucleotidase is a virulence factor conferring mammary fitness in bovine mastitis. *PLoS Pathogens* 20(11), e1012628. <https://doi.org/10.1371/journal.ppat.1012628>
- Ginovska, B., Raugei, S., Ragsdale, S.W., Ohmer, C., Sarangi, R., 2025. Structural and mechanistic advances in the chemistry of methyl-coenzyme M reductase (MCR). *Accounts of Chemical Research* 58(6), 824–833. <https://doi.org/10.1021/acs.accounts.4c00730>
- Hashim, FJ., Vichitphan, S., Han, J., Vichitphan, K., 2021b. Alternative approach for specific tyrosinase inhibitor screening: Uncompetitive inhibition of tyrosinase by *Moringa oleifera*. *Molecules* 26(15), 4576. <https://doi.org/10.3390/molecules26154576>
- Hashim, FJ., Vichitphan, S., Boonsiri, P., Vichitphan, K., 2021a. Neuroprotective assessment of *Moringa oleifera* leaves extract against

- oxidative-stress-induced cytotoxicity in SHSY5Y neuroblastoma cells. *Plants* 10(5), 889. <https://doi.org/10.3390/plants10050889>
- Khusro, A., Aarti, C., Salem, A.Z., Pliego, A.B., Rivas-Caceres, R.R., 2020. Methyl-coenzyme M reductase (MCR) receptor as potential drug target for inhibiting methanogenesis in horses using *Moringa oleifera* L.: an in silico docking study. *Journal of Equine Veterinary Science* 88, 102949. <https://doi.org/10.1016/j.jevs.2020.102949>
- Matra, M., Phupaboon, S., Totakul, P., Prommachart, R., Wanapat, M., 2025. Microencapsulated-bioactive compounds from medicinal leaf extracts used as feed supplements: Effects on in vitro rumen fermentation, microbial population and methane mitigation. *International Journal of Agriculture and Biosciences* 14(2), 258–264. <https://doi.org/10.47278/journal.ijab/2025.001>
- Patra, A.K., Saxena, J., 2010. A new perspective on the use of plant secondary metabolites to inhibit methanogenesis in the rumen. *Phytochemistry* 71(11-12), 1198–1222. <https://doi.org/10.1016/j.phytochem.2010.05.010>
- Phupaboon, S., Matra, M., Prommachart, R., Totakul, P., Wanapat, M., 2025c. In vitro fermentation studies through synergistic effects of antioxidant phytonutrients derived from encapsulated medical plants. *International Journal of Agriculture and Biosciences* 14(6), 1113–1121. <https://doi.org/10.47278/journal.ijab/2025.xxx>
- Phupaboon, S., Hashim, F.J., Punyappa-Path, S., Kanpipit, N., klinsukon, C., Saowakoon, S., 2025. Formulation, characterization and effectiveness of microencapsulated phytochemicals-derived essential oil against animal-pathogenic bacteria: in vitro and in silico studies. *International Journal of Veterinary Science* 14(5), 998–1007. <https://doi.org/10.47278/journal.ijvs/2025.057>
- Phupaboon, S., Matra, M., Prommachart, R., Totakul, P., Supapong, C., Wanapat, M., 2022. Extraction, characterization, and chitosan microencapsulation of bioactive compounds from *Cannabis sativa* L., *Cannabis indica* L., and *Mitragyna speciosa* K. *Antioxidants* 11(11), 2103. <https://doi.org/10.3390/antiox11112103>
- Phupaboon, S., Muslykhah, U., Suriyapha, C., Sommai, S., Matra, M., Dagaew, G., ... Wanapat, M., 2025. Encapsulated phytochemicals enhance in vitro rumen fermentation and reduce methane emissions. *BMC Veterinary Research* 21(1), 1–11. <https://doi.org/10.1186/s12917-025-04812-x>
- Prachumchai, R., Suriyapha, C., Dagaew, G., Sommai, S., Matra, M., Phupaboon, S., ... Wanapat, M., 2024. Microencapsulation of lemongrass and mangosteen peel as phytochemicals to gas kinetics, fermentation, degradability, methane production, and microbial population using in vitro gas technique. *Plos One* 19(6), e0304282. <https://doi.org/10.1371/journal.pone.0304282>
- Punyaappa-Path, S., Phupaboon, S., Kiatprasert, P., Toomthong, P., Piyadeatsoontorn, S., Thonsri, U. ... Rattanachaikunsopon, P., 2025. Protein digestibility, protein patterns, and antioxidant activities of quinoa natto supplemented with Thai traditional herbs. *Natural Resources for Human Health* 5(2), 217–226. <https://doi.org/10.53365/nrfhh/200471>
- Said-Al Ahl, H.A., Hikal, W.M., Mahmoud, A.A., 2017. Biological activity of *Moringa peregrina*: A review. *American Journal of Food Science and Health* 3(4), 83–87.
- Shalaby, E.A., Shanab, S.M., El-Raheem, W.M.A., Hanafy, E.A., 2022. Biological activities and antioxidant potential of different biosynthesized nanoparticles of *Moringa oleifera*. *Scientific Reports* 12(1), 18400. <https://doi.org/10.1038/s41598-022-23164-2>
- Somma, S., Suntara, C., Matra, M., Phupaboon, S., Dagaew, G., Suriyapha, C., ... Wanapat, M., 2025. Effects of feed supplementation with fruit peel pellet microencapsulated phytonutrients on rumen fermentation efficiency, in vitro degradability, methane production and microbial diversity. *Animal Bioscience* 38(7), 1435–1445. <https://doi.org/10.5713/ab.24.0660>
- Stefaniu, A., Pirvu, L.C., 2022. In silico study approach on a series of 50 polyphenolic compounds in plants: A comparison on the bioavailability and bioactivity data. *Molecules* 27(4), 1413. <https://doi.org/10.3390/molecules27041413>
- Suriyapha, C., Pongsub, S., Sommai, S., Phupaboon, S., Dagaew, G., Muslykhah, U., ... Wanapat, M., 2025. In vitro fermentation characteristics, microbial changes and gas production of microencapsulated phytonutrient pellets at varying dietary crude protein levels. *Scientific Reports* 15(1), 11214. <https://doi.org/10.1038/s41598-025-95748-7>
- Wanapat, M., Prachumchai, R., Dagaew, G., Matra, M., Phupaboon, S., Sommai, S., Suriyapha, C., 2024. Potential use of seaweed as a dietary supplement to mitigate enteric methane emission in ruminants. *Science of The Total Environment* 931, 173015. <https://doi.org/10.1016/j.scitotenv.2024.173015>
- Wanapat, M., Suriyapha, C., Dagaew, G., Matra, M., Phupaboon, S., Sommai, S., ... Muslykhah, U., 2024. Sustainable livestock production systems are key to ensuring food security resilience in response to climate change. *Agriculture and Natural Resources* 58(4), 537–546. <https://doi.org/10.34044/j.anres.2024.58.4.13>
- Wanapat, M., Suriyapha, C., Dagaew, G., Prachumchai, R., Phupaboon, S., Sommai, S., Matra, M., 2024. The recycling of tropical fruit peel waste-products applied in feed additive for ruminants: food manufacturing industries, phytonutrient properties, mechanisms, and future applications. *Journal of Agriculture and Food Research* 17, 101234. <https://doi.org/10.1016/j.jafr.2024.101234>
- Yeshi, K., Turpin, G., Jantsho, T., Wangchuk, P., 2022. Indigenous uses, phytochemical analysis, and anti-inflammatory properties of Australian tropical medicinal plants. *Molecules* 27(12), 3849. <https://doi.org/10.3390/molecules27123849>
- Zahran, E.M., Albohy, A., Khalil, A., Ibrahim, A.H., Ahmed, H.A., El-Hossary, E.M., ... Abdelmohsen, U.R., 2020. Bioactivity potential of marine natural products from Scleractinia-associated microbes and in silico anti-SARS-COV-2 evaluation. *Marine Drugs* 18(12), 645. <https://doi.org/10.3390/md18120645>