

Original Research

View Article online



Received 9 December 2025
Revised 12 January 2026
Accepted 29 January 2026
Available Online 17 April 2026

Edited by Balamuralikrishnan
Balasubramanian

KEYWORDS:

Herbal medicine
Basella alba var. *scandens*
Crop-wild variety
GC-MS profiling
Antidiabetic and hypolipidemic properties

Natr Resour Human Health 2026; 6 (2): 450–456
<https://doi.org/10.53365/nrhh/217530>
eISSN: 2583-1194
Copyright © 2026 Visagaa Publishing House

Role of crop-wild variety *Basella alba* var. *scandens* (Basellaceae) as a dietary supplement in controlling diabetes and hyperlipidemia

Sagnik Das¹, Saubhik Das^{2*}

¹Department of Microbiology, St. Xavier's College, Kolkata – 700016, West Bengal, India

²Department of Botany, Bidhannagar College (Affiliated to West Bengal State University), Salt Lake, Kolkata – 700064, West Bengal, India


ABSTRACT: *Basella alba* L. (Basellaceae) is a popular leafy vegetable in the hot humid tropics of India. A lesser-known crop-wild variety, *B. alba* var. *scandens*, is equally palatable and is grown in rural areas in India and consumed locally. Besides nutritive values, *B. alba* has been reported to have many health-promoting properties such as—antidiabetic, hypolipidemic, and antioxidant. Methanolic leaf extracts of both cultivated and crop-wild variants of *B. alba* were subjected to GC-MS analysis, which revealed the presence of a plethora of bioactive phytochemicals, including secondary metabolites, some of which are believed to contribute to these properties. A total of 230 phytochemicals (including fatty acids and fatty acid derivatives, phenolics, terpenes and terpenoids, phytosterols, steroids, alkanes etc.) were identified, of which 72 were found to be present in both. A large number of phytochemical compounds were identified, with a wide range of pharmaceutical, medicinal, and nutritive properties. The therapeutic potential of *B. alba*, a herbal medicine, used to manage hyperglycemia and hyperlipidemia, was correlated with the phytochemical components present in its methanolic leaf extract, a correlation that has not yet been reported. Crop-wild variety appeared very competitive, even better than its cultivated relative, in terms of pharmaceutical and medicinal values, especially as anti-diabetic and antihyperlipidemic agents.

* Corresponding author.

E-mail address: sbhk_das@yahoo.com (Saubhik Das)

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

GRAPHICAL ABSTRACT



1. INTRODUCTION

Leafy vegetables are among the palatable ingredients of the daily diet in the Indian subcontinent due to their taste and nutritional value. They are the source of nutrients such as carbohydrates, dietary protein, vitamins, minerals, and dietary fiber, as well as a storehouse of a broad spectrum of bioactive components, mostly secondary metabolites (Kadam et al., 2017). These bioactive components are attributed to their antioxidant, antidiabetic, hypocholesterolemic, hypolipidemic, and antimicrobial properties, which help to reduce the risk of developing chronic diseases and cardiovascular diseases (Genkinger et al., 2004). Chronic disorders like hyperlipidemia, diabetes, and hypercholesterolemia are direct or indirect causes of cardiovascular diseases (Rana et al., 2007).

Basella alba L. (Basellaceae), commonly known as vine spinach, is one of the favorite leafy vegetables in the hot humid tropics of India as well as in tropical Asia and Africa (Tindall, 1983). The variability in morphology and habit in *B. alba* has recently been addressed through the introduction of three varieties—var. *alba*, var. *rubra*, and var. *scandens*, of which the first two are cultivated varieties and *B. alba* var. *scandens* is crop-wild variety, new for science (Das, 2025).

The lesser-known, underutilized, more palatable crop-wild variety *B. alba* var. *scandens* has been identified from the lower Gangetic plain of West Bengal and is morphologically distinguishable from the other two cultivated varieties. According to ethnopharmacological reports, extracts or formulations of *Basella* have been used in various countries to treat various ailments as traditional medicine (Kumar et al., 2013).

In this study, attempts have been made to identify the bioactive phytochemicals in the leaves of popular cultivated and crop-wild varieties of *B. alba*, which confer various health-promoting benefits, using GC-MS analysis.

2. MATERIALS AND METHODS

2.1. Materials

Two varieties of popular leafy vegetable *B. alba* L. (*B. alba* L. var. *alba* and *B. alba* L. var. *scandens* Saubhik Das) were taken into consideration, and their taxonomic identity was ratified from Central National Herbarium, Acharya Jagadish Chandra Bose Indian Botanic Garden, Sibpur, Howrah, West Bengal, India. All the chemicals used were of analytical grade.

2.2. Methods

2.2.1. Sample preparation

Fresh leaf samples of each plant specimen were dried in the shade and ground in powder. Leaf powder of 10 g from each sample was extracted with 50 mL of methanol for 48 hours with continuous stirring. The crude organic extracts were partially purified by passing through a Silica gel column with an intercalated layer of activated charcoal. The samples were then concentrated using a rotary evaporator and stored at 4°C.

2.2.2. GC-MS analysis

The GC-MS of methanolic leaf extracts was performed on a Shimadzu (Model GCMS-QP2020) series of GC-MS system (Japan) with auto-sampler. SH-Rtx-5 fused silica capillary column (length 30 m × 0.25 mm ID × 0.25 µm film thickness) was used, and pure helium gas was applied as a carrier gas at a constant flow rate of ±1 mL/min and pressure 100 Kpa. The column temperature was programmed from 50°C to 290°C at 5°C increase/min, then held isothermal for 10 minutes. The injector temperature was set at 250°C, and an autosampler injection volume of 1 µL was injected with splitless mode. The ionization energy was 70 eV with a scan time of 0.3 seconds and a mass range of 45–500 AMU.

The compounds were identified using two methods. Specific bioactive compounds represented by certain peaks in the chromatogram were identified by peak area percentage, based on their retention times comparing with those reported in the literature for SH-Rtx-5 columns. The mass spectrum of GC-MS was interpreted using the National Institute of Standards and Technology (NIST) and Wiley spectral libraries. The chemical identity of the bioactive compounds was confirmed by comparison of the mass spectral fragmentation pattern of an unknown compound with the spectrum of the known component stored in the Wiley spectral library and NIST (National Institute of Standards and Technologies, Mass Spectra Libraries) database.

3. RESULTS AND DISCUSSION

3.1. Morphological features

The crop-wild variety (*B. alba* var. *scandens*) is morphologically very distinct from the two popular cultivated varieties (*B. alba* var. *alba* and *B. alba* var. *rubra*). The popular cultivated varieties are non-twining, trailing, or procumbent with thick fleshy stems and large oval leaf, while the crop-wild relative is always twining with a thin cord-like stem and smaller cordate leaf. The crop-wild variety is generally grown in the kitchen garden, consumed locally mainly by growers due to

its palatability and nutritive value, but it has yet to achieve wide acceptance.

3.2. GC-MS of methanolic leaf extract

Phytochemical profiling of organic leaf extracts of both cultivated and crop-wild variety (Figure 1) revealed presence of 230 bioactive phytochemicals belonging different chemical classes such as fatty acids and fatty acid derivatives, alkanes, phenolic compounds including flavonoids, different types of terpenes and triterpenoids, phytosterols, steroids, ketones, and organic acid esters (Table 1), of which 72 were found to be present in both the variants. A large number of identified phytochemicals are reported to have antidiabetic or hypoglycemic (Table 2) and hypolipidemic or hypocholesterolemic (Table 3) potentials.


In terpenes and terpenoid content, the crop-wild variety is far enriched (16.04%) than its cultivated relative (6.79%), while it is very much competitive (26.72%) with its cultivated relative (30.75%) in terms of fatty acids and fatty acid derivatives. In terms of phenolics and phytosterols, the cultivated variety (7.93% and 1.19%) showed little superiority over crop-wild variety (3.52% and 1.11%).

3.3. Phytochemicals with hypoglycemic hypolipidemic effect

Diabetes mellitus is one of the common health problems in the modern world. Synthetic drugs are commonly applied to combat hyperglycemia, though prolonged use of such drugs may have side effects. Plant-derived bioactive phytochemicals have significant potential for the management of diabetes and hyperglycemia (Balbaa et al., 2021). Ethnobotanical studies have identified nearly 1200 rare plants worldwide that may have antidiabetic potential (Arumugam et al., 2013).

Several terpenoids and diterpenes have been reported as antidiabetic agents, and many have advanced to clinical trials (Mohammed et al., 2022; Panigrahy et al., 2021). Dietary phenolics, especially flavonoids, have demonstrated excellent antihyperglycemic effects (Keshari et al., 2016), inhibiting α -amylase and α -glucosidase enzymes and glucose transporters (Pascual et al., 2015). Terpenoids can inhibit enzymes responsible for the development of insulin resistance and glucose metabolism. Triterpenes appeared promising in the management of diabetes and related problems such as retinopathy, neuropathy, nephropathy, and impaired wound healing (Putta et al., 2016; Mabhida et al., 2018).

Flavonoids have been shown to be therapeutically active in reducing total cholesterol, triglycerides, LDL-cholesterol, and apolipoprotein levels, while increasing HDL-cholesterol, bile acid secretion, and lipid catabolism



herbal remedies with no side effects for managing chronic diseases such as diabetes, hypercholesterolemia, and hyperlipidemia. Proper identification and adequate exploitation of traditional herbs and phytotherapy have been encouraged by World Health Organization (WHO) since 1980.

Table 1

Distribution of phytochemical groups in cultivated and crop-wild variety of *Basella alba* L.

Broad groups	Phytochemical groups	Cultivated variety	Crop-wild variety
		Pick area %	Pick area %
Phenolics	Phenolic compound	6.27	3.28
	Flavonoids	1.66	0.24
Terpenes and Terpenoids	Monoterpene and Monoterpenoid	1.05	1.57
	Monoterpenoid derivative	1.54	2.13
	Diterpene and Diterpenoids	0.56	0.65
	Diterpene derivatives	2.40	1.60
	Triterpenoid	0.10	0.85
	Sesquiterpene and Sesquiterpenoid	0.85	2.78
	Sesterterpenoid	0.29	6.46
Sterols	Phytosterols	1.19	1.11
	Steroid	0.52	1.31
	Alkane	15	12.02
	Fatty acid and Fatty acid derivative	30.57	26.72

B. alba is traditionally used to control blood sugar level. The contributory factors for this hypoglycemic effect have yet to be identified. Mucilage present in the plant was considered responsible for that hypoglycemic effect (Palanuvej et al., 2009). Previous studies have also reported its antidiabetic (Nirmala et al., 2009; Bamidele et al., 2014), hypocholesterolemic, and anti-atherosclerotic properties (Bhaskaran et al., 2015). Aqueous extract of *B. rubra* (400mg/kg body weight for 30 days) was found to significantly reduce the blood sugar level (Nirmala et al., 2011). Appreciable presence of dietary phenolics, terpenes and terpenoids, sterols, fatty acids and fatty acid derivatives, besides mucilage may be contributory factors in lowering blood sugar level as well as the hypolipidemic effect of *B. alba*. The crop-wild variety appeared more competitive, particularly in terms of phytochemical components.

4. CONCLUSION

The study established the crop-wild variety *B. alba* var. *scandens* not only as a palatable substitute for cultivated *Basella* but also as an herbal medicine. It appeared very competitive, even better than its cultivated popular variety, in terms of phytochemical components—terpenes, terpenoids, and phenolics, including flavonoids, and sterols. Significant presence of such bioactive components may be responsible

Table 2

List of bioactive phytochemicals with hypoglycemic or antidiabetic effects identified in the organic leaf extracts of cultivated and crop-wild variety of *B. alba* L.

Sl. No	Bioactive phytochemical	Chemical nature	Mol. formula	Cultivated variety		Crop-wild variety	
				Rt	Area %	Rt	Area %
1	α/γ – Butyrolactone	Butanolide	C ₄ H ₆ O ₂	5.240	0.13	–	–
2	4H-Pyran-4-one,2,3-dihydro-3,5-dihydroxy-6-methyl-	Flavonoids	C ₆ H ₈ O ₄	10.970	1.66	–	–
3	Tetradecanoic acid or Myristic acid	Fatty acid	C ₁₄ H ₂₈ O ₂	26.695	0.37	26.720	0.65
4	Neophytadiene	Sesquiterpenoid	C ₂₀ H ₃₈	28.335	0.07	–	–
5	14- β -Pregna OR 14B-Pregname	Steroid	C ₂₁ H ₃₆	30.085	0.52	30.095	1.22
6	Hexadecanoic acid	Saturated fatty acid	C ₁₆ H ₃₂ O ₂	30.970	5.90	32.145	0.66
7	9-Octadecanoic acid (Z)-	Unsaturated fatty acid	C ₁₈ H ₃₄ O ₂	32.285	0.73	–	–
8	Phytol	Diterpene alcohol	C ₂₀ H ₄₀ O	33.855	2.40	33.865	1.60
8	1-Iododotriacontane	Alkyl iodide	C ₃₂ H ₆₅ I	–	–	35.545	0.99
9	α -Tocospiro B	Terpenoids	C ₂₉ H ₅₀ O ₄	–	–	46.025	2.75
10	α -Tocospiro B	Terpenoids	C ₂₉ H ₅₀ O ₄	46.320	0.29	46.335	3.71
11	Dihydrobrassicasterol OR (S) – Campesterol	Phytosterol	C ₂₈ H ₄₈ O	–	–	51.960	0.18
12	Stigmasterol	Phytosterol	C ₂₉ H ₄₈ O	52.605	0.31	52.615	0.45
13	γ -Sitosterol OR Clionasterol	Phytosterol	C ₂₉ H ₅₀ O	53.850	0.70	53.850	0.36
14	β -Amyrin OR α -Amyrin	Triterpenoid	C ₃₀ H ₅₀ O	54.750	0.10	–	–
15	Lupeol OR Fagarasterol	Triterpenoid	C ₃₀ H ₅₀ O	–	–	55.915	0.53

Table 3

List of bioactive phytochemicals with hypolipidemic effect identified in the organic leaf extract of cultivated *B. alba* L. and its crop-wild variant.

Sl	Bioactive phytochemical	Chemical nature	Mol. formula	Cultivated variety		Crop-wild variety	
				Rt	Area %	Rt	Area %
1	Dodecanoic acid or Lauric acid	Fatty acid	C ₁₂ H ₂₄ O ₂	22.110	0.13	22.095	0.07
2	Tetradecanoic acid or Myristic acid	Fatty acid	C ₁₄ H ₂₈ O ₂	26.695	0.37	26.720	0.65
3	14-β-Pregna OR 14B-Pregnane	Steroid	C ₂₁ H ₃₆	30.085	0.52	30.095	1.22
4	Hexadecanoic acid	Saturated fatty acid	C ₁₆ H ₃₂ O ₂	30.970	5.90	–	–
5	Linoleic acid, methyl ester	Fatty acid derivative	C ₁₉ H ₃₄ O ₂	33.510	0.12	33.515	0.12
6	Linolenic acid methyl ester	Fatty acid methyl ester	C ₁₉ H ₃₂ O ₂	33.640	0.19	–	–
7	Linoleic acid Or Linoelaidic acid	Unsaturated fatty acid	C ₁₈ H ₃₂ O ₂	34.240	0.25	–	–
8	Octadecanoic acid OR Stearic acid	Saturated fatty acid	C ₁₈ H ₃₆ O ₂	34.765	0.98	34.795	1.08
9	Octacosanol	Fatty alcohol	C ₂₈ H ₅₈ O	41.995	1.15	42.000	0.93
10	Dihydrobrassicasterol OR (S) – Campesterol	Phytosterol	C ₂₈ H ₄₈ O	–	–	51.960	0.18
11	Stigmasterol	Phytosterol	C ₂₉ H ₄₈ O	52.605	0.31	52.615	0.45
12	γ-Sitosterol OR Clionasterol	Phytosterol	C ₂₉ H ₅₀ O	53.850	0.70	53.850	0.36
13	β-Amyrin OR α-Amyrin	Triterpenoid	C ₃₀ H ₅₀ O	54.750	0.10	–	–
14	Lupeol OR Fagarasterol	Triterpenoid	C ₃₀ H ₅₀ O	–	–	55.915	0.53
15	Simiarenol	Triterpenoid	C ₃₀ H ₅₀ O	–	–	56.185	0.32

for its specific medicinal properties in managing hyperglycemia, increased level of blood cholesterol, triglycerides, and LDL-cholesterol.

ACKNOWLEDGEMENTS

The authors express their sincere gratitude to Mr. Sandip Chowdhury, Sr. Technical Officer, Indian Institute of Chemical Biology, Kolkata, for his assistance and cooperation in GC-MS analysis, also to the Director, CSIR-Indian Institute of Chemical Biology, Kolkata, for necessary assistance in the study.

AUTHORS' CONTRIBUTIONS

Saubhik Das: Research concept and designing of research plan, collection of plant materials, data analysis, and preparation of manuscript.

Sagnik Das: Data analysis, review of references, and compilation of data.

CONFLICT OF INTEREST

The authors have no relevant conflict of interest in the present work.

FUNDING

This work was not funded by any funding agency.

ORCID

Sagnik Das [0009-0001-5609-5980](https://orcid.org/0009-0001-5609-5980)
Saubhik Das [0000-0002-0056-7766](https://orcid.org/0000-0002-0056-7766)

REFERENCES

- Arumugam, G., Manjula, P., Paari, N., 2013. A review: anti diabetic medicinal plants used for diabetes mellitus. *Journal of Acute Disease* 2(3), 196–200. [https://doi.org/10.1016/s2221-6189\(13\)60126-2](https://doi.org/10.1016/s2221-6189(13)60126-2).
- Bahr, T., Butler, G., Rock, C., Welburn, K., Allred, K., Rodriguez, D., 2021. Cholesterol-lowering activity of natural mono- and sesquiterpenoid compounds in essential oils: a review and investigation of mechanisms using in silico protein–ligand docking. *Phytotherapy Research* 35(8), 4215–4245. <https://doi.org/10.1002/ptr.7083>.
- Balbaa, M., El-Zeftawy, M., Abdulmalek, S.A., 2021. Therapeutic screening of herbal remedies for the management of diabetes. *Molecules* 26(22), 1–18. <https://doi.org/10.3390/molecules26226836>.
- Bamidele, O., Arokoyo, D.S., Akinnuga, A.M., Oluwarole, A.O., 2014. Antidiabetic effect of aqueous extract of *Basella alba* leaves and metformin in alloxan-induced diabetic albino rats. *African Journal of Biotechnology* 13(24), 2455–2458. <http://www.academicjournals.org/AJB>.

- Bhaskaran, G., Salvamani, S., Azlan, A., Ahmad, S.A., Yeap, S.K., Shukor, M.Y., 2015. Hypocholesterolemic and Anti-atherosclerotic potential of *Basella alba* leaf extract in hypercholesterolemia induced rabbits. Evidence-Based Complementary and Alternative Medicine 9, 509–517. <https://doi.org/10.1155/2015/751714>.
- Chen, G., Wang, H., Zhang, X., Yang, S.-T., 2014. Nutraceuticals and functional foods in the management of hyperlipidemia. Critical Reviews in Food Science and Nutrition 54(9), 1180–1201. <https://doi.org/10.1080/10408398.2011.629354>.
- Das, S., 2025. Notes about *Basella alba* (Basellaceae) with description of a new variety from West Bengal (India). Phytotaxa 731(1), 78–86. <https://www.mapress.com/pt/>.
- Genkinger, J.M., Platz, E.A., Hoffman, S.C., Comstock, G.W., Helzlsouer, K.J., 2004. Fruit, vegetable, and antioxidant intake and all-cause, cancer, and cardiovascular disease mortality in a community-dwelling population in Washington County, Maryland. American Journal of Epidemiology 160(12), 1223–1233. <https://doi.org/10.1093/aje/kwh339>.
- Kadam, D.M., Monika, S., Devinder, K., 2017. Vegetables Process and Bioactive Compounds. Studium Press (India) Pvt. Ltd., New Delhi.
- Keshari, A.K., Kumar, G., Kushwaha, P.S., Bhardwaj, M., Kumar, P., Rawat, A., Kumar, D., Prakash, A., Ghosh, B., Saha, S., 2016. Isolated flavonoids from *Ficus racemosa* stem bark possess anti-diabetic, hypolipidemic and protective effects in albino Wistar rats. Journal of Ethnopharmacology 181, 252–262. <https://doi.org/10.1016/j.jep.2016.02.004>.
- Kumar, S., Prasad, A.K., Iyer, S.V., Vaidya, S.K., 2013. Systematic, Pharmacognostical, phytochemical and pharmacological review on an ethnomedicinal plant, *Basella alba* L. Journal of Pharmacognosy and Phytotherapy 5(4), 53–58.
- Mabhida, S.E., Dlodla, P.V., Johnson, R., Ndlovu, M., Louw, J., Opoku, A.R., Mosa, R.A., 2018. Protective effect of triterpenes against diabetes-induced β -cell damage: An overview of in vitro and in vivo studies. Pharmacological Research 137, 179–192. <https://doi.org/10.1016/j.phrs.2018.10.004>.
- Mamgain, A., Dhruv, S., 2025. Usage of dietary supplements among diabetic patients in Vadodara, India: A cross-sectional study. Nutrition Health 21, 2601060251343975. <https://doi.org/10.1177/02601060251343975>.
- Martin, J., Wang, Z.Q., Zhang, X.H., Wachtel, D., Volaufova, J., Mathews, D.E., Cefalu, W.T., 2006. Chromium picolinate supplementation attenuates body weight gain and Increases insulin sensitivity in subjects with type 2 diabetes. Diabetes Care 29(8), 1826–1832. <https://doi.org/10.2337/dc06-0254>.
- Mohammed, A., Tajuddeen, N., Ibrahim, M.A., Isah, M.B., Aliyu, A.B., Islam, M.S., 2022. Potential of diterpenes as anti-diabetic agents: evidence from clinical and pre-clinical studies. Pharmacological Research 179, 106158. <https://doi.org/10.1016/j.phrs.2022.106158>.
- Nguyen, T.T., 1999. The cholesterol-lowering action of plant stanol esters. The Journal of Nutrition 129(12), 2109–2112. <https://doi.org/10.1093/jn/129.12.2109>.
- Nirmala, A., Saroja, S., Vasanthi, H.R., Lalitha, G., 2009. Hypoglycemic effect of *Basella rubra* in streptozotocin-induced diabetic albino rat. Journal of Pharmacognosy and Phytotherapy 1(2), 25–30.
- Nirmala, A., Saroja, S., Gayathri, Devi G., 2011. Antidiabetic activity of *Basella rubra* and its relationship with the antioxidant property. Biotechnology Journal International 1(1), 1–9. <https://doi.org/10.9734/BBJ/2011/223>.
- Palanuvej, C., Hokputsa, S., Tunsaringkarn, T., Ruangrunsi, N., 2009. In vitro glucose entrapment and alpha-glucosidase inhibition of mucilaginous substances from selected Thai medicinal plant. Scientia Pharmaceutica 77(4), 837–850.
- Panigrahy, S.K., Bhatt, R., Kumar, A., 2021. Targeting type II diabetes with plant terpenes: the new and promising antidiabetic therapeutics. Biologia 76(1), 241–254. <https://doi.org/10.2478/s11756-020-00575-y>.
- Pascual, J.M., Wang, D., De Vivo, D.C., 2015. Glucose transporter type I deficiency and other glucose flux disorders, In: Rosener, R.N., Pascual, J.M. (Eds.), Rosenberg's Molecular and Genetic Basis of Neurological and Psychiatric Disease 5th Edn. Academic Press, Boston, MA, USA, pp. 649–662.
- Putta, S., Yarla, N.S., Kilari, E.K., Surekha, C., Aliev, G., Divakara, M.B., Santosh, M.S., Ramu, R., Zameer, F., Mn, N.P. et al. 2016. Therapeutic Potentials of Triterpenes in Diabetes and its Associated Complications. Current Topics in Medicinal Chemistry 16(23), 2532–2542. <https://doi.org/10.2174/1568026616666160414123343>.
- Rana, J.S., Nieuwdorp, M., Jukema, J.W., Kastelein, J.J.P., 2007. Cardiovascular metabolic syndrome—an interplay of obesity, inflammation, diabetes and coronary heart disease. Diabetes, Obesity and Metabolism 9(3), 218–232. <https://doi.org/10.1111/j.1463-1326.2006.00594.x>.
- Sknepnek, A., Miletić, D., Stupar, A., Salević-Jelić, A., Nedović, V., Cvetanović Kljakić, A., 2025. Natural solutions for diabetes: the therapeutic potential of plants and mushrooms. Frontiers in Nutrition. 12, 1511049. <https://doi.org/10.3389/fnut.2025.1511049>.
- Tindall, H.D., 1983. Vegetables in the Tropics. Macmillan Press Ltd., London. <https://doi.org/10.1007/978-1-349-17223-8>.
- Yeh, G.Y., Eisenberg, D.M., Kaptchuk, T.J., Phillips, R.S., 2003. Systematic review of herbs and dietary supplements for glycemic control in diabetes. Diabetes Care 26(4), 1277–1294.
- Zeka, K., Ruparelia, K., Arroo, R.R., Budriesi, R., Micucci, M., 2017. Flavonoids and their metabolites: prevention in cardiovascular diseases and diabetes. Diseases 5(3), 19. <https://doi.org/10.3390/diseases5030019>.